



AGROCLIMATIC ZONES OF CUTTACK DISTRICT:



STANDARD PRECIPITATION INDEX AND NDVI MAP





LEGEND(Medium range block level rainfall forecast)	
	Very light to light rainfall
	Light to moderate rainfall
	Moderate to heavy rainfall
	Heavy rainfall
	Very heavy rainfall
	Extremely heavy rainfall



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com



Bulletin No.-140
Date- 10/11/2020



Extended range outlook for subsequent 1 week (6th to 12th days)
(11th to 17th November 2020)

Met Sub division	Departure (%)	Category
Odisha	-99.74	Below Normal

Dr Sujata Sethy
(Nodal officer)

















Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427

Weather(Past and Forecasted) based Disease and pest in Major Crops and vegetables






SI No	Crop	Stage and condition of the crop	Anticipated Disease and pest	
1	Rice	Panicle initiation to Flowering stage		Leaf folder
				BPH
				Gundhi bug
				Gall midge
				Hispa
				False smut
				Bacterial leaf Blight
2	Sugarcane	Grand growth stage to ripening		Ring spot
				Top borer
				Wooly Aphid
				Scale insect
				White fly
				Mealy bug
3	Vegetables(Brinjal and okra)	Vegetative to fruiting		Fruit and shoot borer



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com



Bulletin No.-140
Date- 10/11/2020

	Okra			YMV
	Chilli , Cowpea			Thrips in Chilli and Aphids in Cowpea
4	Fruit (Papaya)	Flowering stage to fruit formation stage		Foot rot
	Banana	Flowering stage to fruit formation stage		Sigatoka disease
	Cashew	Vegetative stage		Tea mosquito bug

Dr Sujata Sethy
(Nodal officer)



**Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area**



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com



Bulletin No.-140
Date- 10/11/2020

AGROMET ADVISORY BULLETIN FOR ATHAGAD

District Past week weather condition and Block past Rainfall condition

The block received 0 mm and district received 0 mm rainfall during the last week. The mean daily maximum and minimum temperature was 32°C and 18°C, respectively. Mean relative humidity of the district was around 70-66%. **According to NDVI map (28.10.2020 to 03.11.2020) the agriculture vigor was light to moderate over few places of Cuttack district whereas according to SPI value (cumulative 4 weeks-08october 2020 to 04 November 2020), moderately wet condition was there.**

Forecasted period: 11th November 2020 to 15th November 2020

Weather advisory: No rainfall is forecasted for upcoming 5 days (Today to Saturday) in the block. The day temperature will be 32-33°C and night temperature will be 19-21°C for upcoming 5 days. Maximum and minimum relative humidity will be 62-77% and 37-49% respectively. There will be mainly clear to partly cloudy sky for next 5 days.

Block	Athagad (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	11/11/2020	12/11/2020	13/11/2020	14/11/2020	15/11/2020
Rainfall (mm)	0	0	0	0	0
T-Max (°C)	32	32	33	33	33
T-Min (°C)	20	20	21	20	19
Cloud Cover (octa)	1	1	6	4	0
RH Max (%)	77	72	62	67	72
RH Min (%)	48	49	40	37	39
Wind Speed (kmph)	8	9	8	8	7
Wind Direction (deg)	63	45	60	45	342

Agromet Advisory

General Advisory: Farmers should go for harvesting of early matured paddy at 85% maturity of the panicles in sunny days. They should keep 5 cm water in paddy fields to maintain saturation during PI to grain filling stage. Carry out spraying of insecticides or pesticides in crops during morning and afternoon hours. Complete nursery sowing for winter vegetable crops. They should cover the nursery beds with polythene in night to protect it from cold temperature. Irrigate the vegetable nursery during morning and evening time as below normal rainfall likely to occur in Odisha in upcoming week. Go for sowing black gram and green gram in upland rice fallow condition by utilizing residual soil moisture.

Kharif Rice:

Paddy is at flowering to milking stage. In Early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop. Current weather favors the infestation of **Leaf Folder**. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days. If infestation of **BPH & WBPH** occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 80g Thiamethoxam or 60g Flonicamid per acre in alternate basis. To prevent crop from **Gundhi bug** apply 5 ml neem based insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. Present weather conditions are favourable for the incidence of **Hispa**. To control, go for spraying of chlorpyrifos @ 2.5ml (or) profenophos @ 2ml /l. In **Gall midge** affected areas apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre or Chlorpyrifos 20EC @ 500ml/acre to manage the pest. For prevention of **false smut** spray the field before panicle initiation or flowering with propiconazole 25%EC @ 2ml/liter of water. Present weather conditions are favourable for incidence of **Bacterial leaf blight**. To control, go for spraying of plantomycin @ 1g/L or Streptocyclin 0.1g/L + COC 3g/L.

PULSES:

This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

Mustard: Taking the advantage of available soil moisture due to recent rain, land preparation and sowing of the crop should be started to get optimum yield. Suitable varieties: Toria- Anuradha, Parbati, Sushree, Indian mustard- Mahyco bold, JD 6. Seed rate- 8-10 Kg/ha Spacing- 30 cm X 10 cm, Fertilizer dose: Toria @ 60:30:30 N-P2O5-K2O Kg/ha Mustard @ 80:40:40 N-P2O5-K2O Kg/ha. Apply 50%N and all P2O5 and K2O fertilizers should be applied as basal.

VEGETABLES:

Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. To control **Fruit and shoot borer** in Brinjal and Okra, Spray Neem based pesticide 1500ppm @ 600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45% SC @ 70ml/acre and Thiodicarb @ 300g/acre at 15 days interval in 200 litre of water. To control **YMV in Okra, thrips in Chilli and aphids in Cowpea** crop, spray Imidacloprid @ 5 ml in 10 litres of water. **Fruits:** **Banana:** Present weather conditions are favourable for the incidence and spread of **sigatoka leaf spot** disease. To curb the spread of this disease, go for spraying of propiconazole @ 0.1% (or) tebuconazole + trifloxystrobin @ 1ml per litre for 2-3 times in 25 days interval. **Papaya:** Present weather conditions are favourable for the incidence of **foot rot** in Papaya. To control go for soil drenching around the base of the stem with Metalaxyl @ 2g/L. **Cashewnut:** Present weather conditions are favourable for the incidence of **tea mosquito bug**. To control go for spraying of Imidacloprid @ 5 ml in 10 litres of water at the time of sprouting. At the time of flowering, go for spraying of lambda cyhalothrin @ 0.6ml/l of water.

Animal husbandry:

Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com



Bulletin No.-140
Date- 10/11/2020

AGROMET ADVISORY BULLETIN FOR BADAMBA

District Past week weather condition and Block past Rainfall condition

The block received 0 mm and district received 0 mm rainfall during the last week. The mean daily maximum and minimum temperature was 32°C and 18°C, respectively. Mean relative humidity of the district was around 70-66%. **According to NDMI map (28.10.2020 to 03.11.2020) the agriculture vigor was light to moderate over few places of Cuttack district whereas according to SPI value (cumulative 4 weeks-08 October 2020 to 04 November 2020), moderately wet condition was there.**

Forecasted period: 11th November 2020 to 15th November 2020

Weather advisory: No rainfall is forecasted for upcoming 5 days (Today to Saturday) in the block. The day temperature will be 31-33°C and night temperature will be 18-20°C for upcoming 5 days. Maximum and minimum relative humidity will be 63-81% and 36-49% respectively. There will be mainly clear to partly cloudy sky for next 5 days.

Block	Badamba (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	11/11/2020	12/11/2020	13/11/2020	14/11/2020	15/11/2020
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	31	32	33	33	32
T-Min(°C)	19	19	20	19	18
Cloud Cover(octa)	1	2	6	4	0
RH Max (%)	81	75	63	66	71
RH Min (%)	48	49	41	36	39
Wind Speed (Kmph)	7	7	7	7	7
Wind Direction (deg)	101	23	68	23	90

Agromet Advisory

General Advisory: Farmers should go for harvesting of early matured paddy at 85% maturity of the panicles in sunny days. They should keep 5 cm water in paddy fields to maintain saturation during PI to grain filling stage. Carry out spraying of insecticides or pesticides in crops during morning and afternoon hours. Complete nursery sowing for winter vegetable crops. They should cover the nursery beds with polythene in night to protect it from cold temperature. Irrigate the vegetable nursery during morning and evening time as below normal rainfall likely to occur in Odisha in upcoming week. Go for sowing black gram and green gram in upland rice fallow condition by utilizing residual soil moisture.

Kharif Rice:

Paddy is at flowering to milking stage. In Early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop. Current weather favors the infestation of **Leaf Folder**. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days. If infestation of **BPH & WBPH** occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 80g Thiamethoxam or 60g Flonicamid per acre in alternate basis. To prevent crop from **Gundhi bug** apply 5 ml neem based insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. Present weather conditions are favourable for the incidence of **Hispa**. To control, go for spraying of chloropyrifos @ 2.5ml (or) profenophos @ 2ml /l. In **Gall midge** affected areas apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre or Chlorpyrifos 20EC @ 500ml/acre to manage the pest. For prevention of **false smut** spray the field before panicle initiation or flowering with propiconazole 25%EC @ 2ml/liter of water. Present weather conditions are favourable for incidence of **Bacterial leaf blight**. To control, go for spraying of plantomycin @ 1g/L or Streptocyclin 0.1g/L + COC 3g/L.

PULSES:

This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

Mustard:

Taking the advantage of available soil moisture due to recent rain, land preparation and sowing of the crop should be started to get optimum yield. Suitable varieties: Toria- Anuradha, Parbati, Sushree, Indian mustard- Mahyco bold, JD 6. Seed rate- 8-10 Kg/ha Spacing- 30 cm X 10 cm, Fertilizer dose: Toria @ 60:30:30 N-P2O5-K2O Kg/ha Mustard @ 80:40:40 N-P2O5-K2O Kg/ha. Apply 50%N and all P2O5 and K2O fertilizers should be applied as basal.

VEGETABLES:

Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. To control **Fruit and shoot borer** in Brinjal and Okra, Spray Neem based pesticide 1500ppm @ 600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45% SC @ 70ml/acre and Thiodicarb @ 300g/acre at 15 days interval in 200 litre of water. To control **YMV in Okra, thrips in Chilli and aphids in Cowpea** crop, spray Imidacloprid @ 5 ml in 10 litres of water. **Fruits: Banana:** Present weather conditions are favourable for the incidence and spread of **sigatoka leaf spot** disease. To curb the spread of this disease, go for spraying of propiconazole @ 0.1% (or) tebuconazole + trifloxystrobin @ 1ml per litre for 2-3 times in 25 days interval. **Papaya:** Present weather conditions are favourable for the incidence of **foot rot** in Papaya. To control go for soil drenching around the base of the stem with Metalaxyl @ 2g/L. **Cashewnut:** Present weather conditions are favourable for the incidence of **tea mosquito bug**. To control go for spraying of Imidacloprid @ 5 ml in 10 litres of water at the time of sprouting. At the time of flowering, go for spraying of lambda cyhalothrin @ 0.6ml/l of water.

Animal husbandry:

Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com



Bulletin No.-140
Date- 10/11/2020

AGROMET ADVISORY BULLETIN FOR BANKI

District Past week weather condition and Block past Rainfall condition

The block received 0 mm and district received 0 mm rainfall during the last week. The mean daily maximum and minimum temperature was 32°C and 18°C, respectively. Mean relative humidity of the district was around 70-66%. **According to NDTV map (28.10.2020 to 03.11.2020) the agriculture vigor was light to moderate over few places of Cuttack district whereas according to SPI value (cumulative 4 weeks-08october 2020 to 04 November 2020), moderately wet condition was there.**

Forecasted period: 11th November 2020 to 15th November 2020

Weather advisory: No rainfall is forecasted for upcoming 5 days (Today to Saturday) in the block. The day temperature will be 32-33°C and night temperature will be 19-20°C for upcoming 5 days. Maximum and minimum relative humidity will be 63-80% and 37-49% respectively. There will be mainly clear to partly cloudy sky for next 5 days.

Block	Banki (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	11/11/2020	12/11/2020	13/11/2020	14/11/2020	15/11/2020
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	32	33	33	33	33
T-Min(°C)	19	20	20	19	19
Cloud Cover(octa)	1	2	6	4	0
RH Max (%)	80	74	63	67	71
RH Min (%)	49	49	40	37	39
Wind Speed (Kmph)	8	8	8	8	8
Wind Direction (deg)	68	30	60	60	90

Agromet Advisory

General Advisory: Farmers should go for harvesting of early matured paddy at 85% maturity of the panicles in sunny days. They should keep 5 cm water in paddy fields to maintain saturation during PI to grain filling stage. Carry out spraying of insecticides or pesticides in crops during morning and afternoon hours. Complete nursery sowing for winter vegetable crops. They should cover the nursery beds with polythene in night to protect it from cold temperature. Irrigate the vegetable nursery during morning and evening time as below normal rainfall likely to occur in Odisha in upcoming week. Go for sowing black gram and green gram in upland rice fallow condition by utilizing residual soil moisture.

Kharif Rice:

Paddy is at flowering to milking stage. In Early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop. Current weather favors the infestation of **Leaf Folder**. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days. If infestation of **BPH & WBPH** occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 80g Thiamethoxam or 60g Flonicamid per acre in alternate basis. To prevent crop from **Gundhi bug** apply 5 ml neem based insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. Present weather conditions are favourable for the incidence of **Hispa**. To control, go for spraying of chloropyrifos @ 2.5ml (or) profenophos @ 2ml /l. In **Gall midge** affected areas apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre or Chlorpyrifos 20EC @ 500ml/acre to manage the pest. For prevention of **false smut** spray the field before panicle initiation or flowering with propiconazole 25%EC @ 2ml/liter of water. Present weather conditions are favourable for incidence of **Bacterial leaf blight**. To control, go for spraying of plantomycin @ 1g/L or Streptocyclin 0.1g/L + COC 3g/L.

PULSES:

This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

Mustard:

Taking the advantage of available soil moisture due to recent rain, land preparation and sowing of the crop should be started to get optimum yield. Suitable varieties: Toria- Anuradha, Parbati, Sushree, Indian mustard- Mahyco bold, JD 6. Seed rate- 8-10 Kg/ha Spacing- 30 cm X 10 cm, Fertilizer dose: Toria @ 60:30:30 N-P2O5-K2O Kg/ha Mustard @ 80:40:40 N-P2O5-K2O Kg/ha. Apply 50%N and all P2O5 and K2O fertilizers should be applied as basal.

VEGETABLES:

Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. To control **Fruit and shoot borer** in Brinjal and Okra, Spray Neem based pesticide 1500ppm @ 600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45% SC @ 70ml/acre and Thiodicarb @ 300g/acre at 15 days interval in 200 litre of water. To control **YMV** in Okra, **thrips** in Chilli and **aphids** in Cowpea crop, spray Imidacloprid @ 5 ml in 10 litres of water. **Fruits:** **Banana:** Present weather conditions are favourable for the incidence and spread of **sigatoka leaf spot** disease. To curb the spread of this disease, go for spraying of propiconazole @ 0.1% (or) tebuconazole + trifoxystrobin @ 1ml per litre for 2-3 times in 25 days interval. **Papaya:** Present weather conditions are favourable for the incidence of **foot rot** in Papaya. To control go for soil drenching around the base of the stem with Metalaxyl @ 2g/L. **Cashewnut:** Present weather conditions are favourable for the incidence of **tea mosquito bug**. To control go for spraying of Imidacloprid @ 5 ml in 10 litres of water at the time of sprouting. At the time of flowering, go for spraying of lambda cyhalothrin @ 0.6ml/l of water.

Animal husbandry: Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com



Bulletin No.-140
Date- 10/11/2020

AGROMET ADVISORY BULLETIN FOR BANKI-DAMPARA

District Past week weather condition

The district received 0 mm rainfall during the last week. The mean daily maximum and minimum temperature was 32°C and 18°C, respectively. Mean relative humidity of the district was around 70-66%. **According to NDVI map (28.10.2020 to 03.11.2020) the agriculture vigor was light to moderate over few places of Cuttack district whereas according to SPI value (cumulative 4 weeks-08october 2020 to 04 November 2020), moderately wet condition was there.**

Forecasted period: 11th November 2020 to 15th November 2020

Weather Advisory: No rainfall is forecasted for upcoming 5 days (Today to Saturday) in the block. The day temperature will be 32-33°C and night temperature will be 19-20°C for upcoming 5 days. Maximum and minimum relative humidity will be 62-79% and 37-49% respectively. There will be mainly clear to partly cloudy sky for next 5 days.

Block	Banki-Dampara (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	11/11/2020	12/11/2020	13/11/2020	14/11/2020	15/11/2020
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	32	33	33	33	33
T-Min(°C)	19	20	20	19	19
Cloud Cover(octa)	1	2	6	4	0
RH Max (%)	79	74	62	67	71
RH Min (%)	49	49	40	37	39
Wind Speed (Kmph)	8	9	8	8	8
Wind Direction (deg)	68	30	60	60	90

Agromet Advisory

General Advisory: Farmers should go for harvesting of early matured paddy at 85% maturity of the panicles in sunny days. They should keep 5 cm water in paddy fields to maintain saturation during PI to grain filling stage. Carry out spraying of insecticides or pesticides in crops during morning and afternoon hours. Complete nursery sowing for winter vegetable crops. They should cover the nursery beds with polythene in night to protect it from cold temperature. Irrigate the vegetable nursery during morning and evening time as below normal rainfall likely to occur in Odisha in upcoming week. Go for sowing black gram and green gram in upland rice fallow condition by utilizing residual soil moisture.

Kharif Rice: Paddy is at flowering to milking stage. In Early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop. Current weather favors the infestation of **Leaf Folder**. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days. If infestation of **BPH & WBPH** occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 80g Thiamethoxam or 60g Flonicamid per acre in alternate basis. To prevent crop from **Gundhi bug** apply 5 ml neem based insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. Present weather conditions are favourable for the incidence of **Hispa**. To control, go for spraying of chloropyrifos @ 2.5ml (or) profenophos @ 2ml /l. In **Gall midge** affected areas apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre or Chlorpyrifos 20EC @ 500ml/acre to manage the pest. For prevention of **false smut** spray the field before panicle initiation or flowering with propiconazole 25%EC @ 2ml/liter of water. Present weather conditions are favourable for incidence of **Bacterial leaf blight**. To control, go for spraying of plantomycin @ 1g/L or Streptocyclin 0.1g/L + COC 3g/L.

PULSES:

This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

Mustard:

Taking the advantage of available soil moisture due to recent rain, land preparation and sowing of the crop should be started to get optimum yield. Suitable varieties: Toria- Anuradha, Parbati, Sushree, Indian mustard- Mahyco bold, JD 6. Seed rate- 8-10 Kg/ha Spacing- 30 cm X 10 cm, Fertilizer dose: Toria @ 60:30:30 N-P2O5-K2O Kg/ha Mustard @ 80:40:40 N-P2O5-K2O Kg/ha. Apply 50%N and all P2O5 and K2O fertilizers should be applied as basal.

VEGETABLES:

Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. To control **Fruit and shoot borer** in Brinjal and Okra, Spray Neem based pesticide 1500ppm @ 600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45% SC @ 70ml/acre and Thiodicarb @ 300g/acre at 15 days interval in 200 litre of water. To control **YMV in Okra, thrips in Chilli and aphids in Cowpea** crop, spray Imidacloprid @ 5 ml in 10 litres of water. **Fruits:** **Banana:** Present weather conditions are favourable for the incidence and spread of **sigatoka leaf spot** disease. To curb the spread of this disease, go for spraying of propiconazole @ 0.1% (or) tebuconazole + trifoxystrobin @ 1ml per litre for 2-3 times in 25 days interval. **Papaya:** Present weather conditions are favourable for the incidence of **foot rot** in Papaya. To control go for soil drenching around the base of the stem with Metalaxyl @ 2g/L. **Cashewnut:** Present weather conditions are favourable for the incidence of **tea mosquito bug**. To control go for spraying of Imidacloprid @ 5 ml in 10 litres of water at the time of sprouting. At the time of flowering, go for spraying of lambda cyhalothrin @ 0.6ml/l of water.

Animal husbandry:

Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com



Bulletin No.-140
Date- 10/11/2020

AGROMET ADVISORY BULLETIN FOR BARANGA

District Past week weather condition and Block past Rainfall condition

The block received 0 mm and district received 0 mm rainfall during the last week. The mean daily maximum and minimum temperature was 32°C and 18°C, respectively. Mean relative humidity of the district was around 70-66%. **According to NDMI map (28.10.2020 to 03.11.2020) the agriculture vigor was light to moderate over few places of Cuttack district whereas according to SPI value (cumulative 4 weeks-08 October 2020 to 04 November 2020), moderately wet condition was there.**

Forecasted period: 11th November 2020 to 15th November 2020

Weather advisory: No rainfall is forecasted for upcoming 5 days (Today to Saturday) in the block. The day temperature will be 32-33°C and night temperature will be 20-21°C for upcoming 5 days. Maximum and minimum relative humidity will be 60-75% and 38-50% respectively. There will be mainly clear to partly cloudy sky for next 5 days.

Block	Baranga (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	11/11/2020	12/11/2020	13/11/2020	14/11/2020	15/11/2020
Rainfall (mm)	0	0	0	0	0
T-Max (°C)	32	32	33	32	32
T-Min (°C)	21	21	21	21	20
Cloud Cover	0	1	7	4	1
RH Max (%)	75	70	60	66	70
RH Min (%)	50	50	41	38	40
Wind Speed (Kmph)	10	11	10	11	10
Wind Direction (deg)	36	45	45	32	23

Agromet Advisory

General Advisory: Farmers should go for harvesting of early matured paddy at 85% maturity of the panicles in sunny days. They should keep 5 cm water in paddy fields to maintain saturation during PI to grain filling stage. Carry out spraying of insecticides or pesticides in crops during morning and afternoon hours. Complete nursery sowing for winter vegetable crops. They should cover the nursery beds with polythene in night to protect it from cold temperature. Irrigate the vegetable nursery during morning and evening time as below normal rainfall likely to occur in Odisha in upcoming week. Go for sowing black gram and green gram in upland rice fallow condition by utilizing residual soil moisture.

Kharif Rice:

Paddy is at flowering to milking stage. In Early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop. Current weather favors the infestation of **Leaf Folder**. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days. If infestation of **BPH & WBPH** occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 80g Thiamethoxam or 60g Flonicamid per acre in alternate basis. To prevent crop from **Gundhi bug** apply 5 ml neem based insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. Present weather conditions are favourable for the incidence of **Hispa**. To control, go for spraying of chloropyrifos @ 2.5ml (or) profenophos @ 2ml /l. In **Gall midge** affected areas apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre or Chlorpyrifos 20EC @ 500ml/acre to manage the pest. For prevention of **false smut** spray the field before panicle initiation or flowering with propiconazole 25% EC @ 2ml/liter of water. Present weather conditions are favourable for incidence of **Bacterial leaf blight**. To control, go for spraying of plantomycin @ 1g/L or Streptocyclin 0.1g/L + COC 3g/L.

PULSES:

This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

Mustard:

Taking the advantage of available soil moisture due to recent rain, land preparation and sowing of the crop should be started to get optimum yield. Suitable varieties: Toria- Anuradha, Parbati, Sushree, Indian mustard- Mahyco bold, JD 6. Seed rate- 8-10 Kg/ha Spacing- 30 cm X 10 cm, Fertilizer dose: Toria @ 60:30:30 N-P2O5-K2O Kg/ha Mustard @ 80:40:40 N-P2O5-K2O Kg/ha. Apply 50%N and all P2O5 and K2O fertilizers should be applied as basal.

VEGETABLES:

Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. To control **Fruit and shoot borer** in Brinjal and Okra, Spray Neem based pesticide 1500ppm @ 600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45% SC @ 70ml/acre and Thiodicarb @ 300g/acre at 15 days interval in 200 litre of water. To control **YMV** in Okra, **thrips** in Chilli and **aphids** in Cowpea crop, spray Imidacloprid @ 5 ml in 10 litres of water. **Fruits:** **Banana:** Present weather conditions are favourable for the incidence and spread of **sigatoka leaf spot** disease. To curb the spread of this disease, go for spraying of propiconazole @ 0.1% (or) tebuconazole + trifloxystrobin @ 1ml per litre for 2-3 times in 25 days interval. **Papaya:** Present weather conditions are favourable for the incidence of **foot rot** in Papaya. To control go for soil drenching around the base of the stem with Metalaxyl @ 2g/L. **Cashewnut:** Present weather conditions are favourable for the incidence of **tea mosquito bug**. To control go for spraying of Imidacloprid @ 5 ml in 10 litres of water at the time of sprouting. At the time of flowering, go for spraying of lambda cyhalothrin @ 0.6ml/l of water.

Animal husbandry: Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com



Bulletin No.-140
Date- 10/11/2020

AGROMET ADVISORY BULLETIN FOR CUTTACK SADAR
District Past week weather condition and Block past Rainfall condition

The block received 0 mm and district received 0 mm rainfall during the last week. The mean daily maximum and minimum temperature was 32°C and 18°C, respectively. Mean relative humidity of the district was around 70-66%. **According to NDMI map (28.10.2020 to 03.11.2020) the agriculture vigor was light to moderate over few places of Cuttack district whereas according to SPI value (cumulative 4 weeks-08 October 2020 to 04 November 2020), moderately wet condition was there.**

Forecasted period: 11th November 2020 to 15th November 2020

Weather advisory: No rainfall is forecasted for upcoming 5 days (Today to Saturday) in the block. The day temperature will be 32-33°C and night temperature will be 20-21°C for upcoming 5 days. Maximum and minimum relative humidity will be 59-75% and 37-50% respectively. There will be mainly clear to partly cloudy sky for next 5 days.

Block	Cuttack sadar (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	11/11/2020	12/11/2020	13/11/2020	14/11/2020	15/11/2020
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	32	32	33	32	32
T-Min(°C)	21	21	21	21	20
Cloud Cover(octa)	0	1	6	4	0
RH Max (%)	75	71	59	65	71
RH Min (%)	49	50	40	37	40
Wind Speed (Kmph)	10	10	10	10	9
Wind Direction (deg)	36	45	45	45	30

Agromet Advisory

General Advisory: Farmers should go for harvesting of early matured paddy at 85% maturity of the panicles in sunny days. They should keep 5 cm water in paddy fields to maintain saturation during PI to grain filling stage. Carry out spraying of insecticides or pesticides in crops during morning and afternoon hours. Complete nursery sowing for winter vegetable crops. They should cover the nursery beds with polythene in night to protect it from cold temperature. Irrigate the vegetable nursery during morning and evening time as below normal rainfall likely to occur in Odisha in upcoming week. Go for sowing black gram and green gram in upland rice fallow condition by utilizing residual soil moisture.

Kharif Rice: Paddy is at flowering to milking stage. In Early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop. Current weather favors the infestation of **Leaf Folder**. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days. If infestation of **BPH & WBPH** occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 80g Thiamethoxam or 60g Flonicamid per acre in alternate basis. To prevent crop from **Gundhi bug** apply 5 ml neem based insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. Present weather conditions are favourable for the incidence of **Hispa**. To control, go for spraying of chloropyrifos @ 2.5ml (or) profenophos @ 2ml /l. In **Gall midge** affected areas apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre or Chlorpyrifos 20EC @ 500ml/acre to manage the pest. For prevention of **false smut** spray the field before panicle initiation or flowering with propiconazole 25%EC @ 2ml/liter of water. Present weather conditions are favourable for incidence of **Bacterial leaf blight**. To control, go for spraying of plantomycin @ 1g/L or Streptocyclin 0.1g/L + COC 3g/L.

Sugarcane:

Remove the borer affected tillers and late formed tillers, tie the cane shoot with two or three together with partially dried lower leaves. Present weather conditions are favourable for the incidence of **ring spot disease** in Sugarcane. To control go for spraying of copperoxychloride @ 3g/l (or) Carbendazim @ 1g/l (or) Mancozeb @ 3g/l for three times once in three weeks. To control **top shoot borers** spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water. For control of Sucking pest in sugarcane like **Woolly aphid, Whitefly, Scale insects, mealybug** in present weather conditions go for alternate spraying of thiomethoxam 25WG @ 0.25 gm/lit or imidacloprid 0.45 ml/lit of water or Dinotefuron 20% SG 3gm/lit of Water.

PULSES:

This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

VEGETABLES: Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. To control **Fruit and shoot borer** in Brinjal and Okra, Spray Neem based pesticide 1500ppm @ 600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45% SC @ 70ml/acre and Thiodicarb @ 300g/acre at 15 days interval in 200 litre of water. To control **YMV in Okra, thrips in Chilli and aphids in Cowpea** crop, spray Imidacloprid @ 5 ml in 10 litres of water. **Fruits:** **Banana:** Present weather conditions are favourable for the incidence and spread of **sigatoka leaf spot** disease. To curb the spread of this disease, go for spraying of propiconazole @ 0.1% (or) tebuconazole + trifoxystrobin @ 1ml per litre for 2-3 times in 25 days interval. **Papaya:** Present weather conditions are favourable for the incidence of **foot rot** in Papaya. To control go for soil drenching around the base of the stem with Metalaxyl @ 2g/L. **Cashewnut:** Present weather conditions are favourable for the incidence of **tea mosquito bug**. To control go for spraying of Imidacloprid @ 5 ml in 10 litres of water at the time of sprouting. At the time of flowering, go for spraying of lambda cyhalothrin @ 0.6ml/l of water.

Animal husbandry:

Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com



Bulletin No.-140
Date- 10/11/2020

AGROMET ADVISORY BULLETIN FOR KANTAPADA

District Past week weather condition and Block past Rainfall condition

The block received 0 mm and district received 0 mm rainfall during the last week. The mean daily maximum and minimum temperature was 32°C and 18°C, respectively. Mean relative humidity of the district was around 70-66%. **According to NDTV map (28.10.2020 to 03.11.2020) the agriculture vigor was light to moderate over few places of Cuttack district whereas according to SPI value (cumulative 4 weeks-08 October 2020 to 04 November 2020), moderately wet condition was there.**

Forecasted period: 11th November 2020 to 15th November 2020

Weather advisory: No rainfall is forecasted for upcoming 5 days (Today to Saturday) in the block. The day temperature will be 31-32°C and night temperature will be 20-22°C for upcoming 5 days. Maximum and minimum relative humidity will be 61-76% and 38-52% respectively. There will be mainly clear to partly cloudy sky for next 5 days.

Block	Kantapada (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	11/11/2020	12/11/2020	13/11/2020	14/11/2020	15/11/2020
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	31	32	32	32	32
T-Min(°C)	22	22	22	21	20
Cloud Cover(octa)	0	2	6	2	2
RH Max (%)	76	71	61	66	69
RH Min (%)	52	52	42	38	41
Wind Speed (Kmph)	11	11	11	12	11
Wind Direction (deg)	36	45	45	32	23

Agromet Advisory

General Advisory: Farmers should go for harvesting of early matured paddy at 85% maturity of the panicles in sunny days. They should keep 5 cm water in paddy fields to maintain saturation during PI to grain filling stage. Carry out spraying of insecticides or pesticides in crops during morning and afternoon hours. Complete nursery sowing for winter vegetable crops. They should cover the nursery beds with polythene in night to protect it from cold temperature. Irrigate the vegetable nursery during morning and evening time as below normal rainfall likely to occur in Odisha in upcoming week. Go for sowing black gram and green gram in upland rice fallow condition by utilizing residual soil moisture.

Kharif Rice: Paddy is at flowering to milking stage. In Early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop. Current weather favors the infestation of **Leaf Folder**. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days. If infestation of **BPH & WBPH** occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 80g Thiamethoxam or 60g Flonicamid per acre in alternate basis. To prevent crop from **Gundhi bug** apply 5 ml neem based insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. Present weather conditions are favourable for the incidence of **Hispa**. To control, go for spraying of chlorpyrifos @ 2.5ml (or) profenophos @ 2ml /l. In **Gall midge** affected areas apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre or Chlorpyrifos 20EC @ 500ml/acre to manage the pest. For prevention of **false smut** spray the field before panicle initiation or flowering with propiconazole 25%EC @ 2ml/litre of water. Present weather conditions are favourable for incidence of **Bacterial leaf blight**. To control, go for spraying of plantomycin @ 1g/L or Streptocyclin 0.1g/L + COC 3g/L.

Sugarcane:

Remove the borer affected tillers and late formed tillers, tie the cane shoot with two or three together with partially dried lower leaves. Present weather conditions are favourable for the incidence of **ring spot disease** in Sugarcane. To control go for spraying of copperoxychloride @ 3g/l (or) Carbendazim @ 1g/l (or) Mancozeb @ 3g/l for three times once in three weeks. To control **top shoot borers** spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water. For control of Sucking pest in sugarcane like **Woolly aphid, Whitefly, Scale insects, mealybug** in present weather conditions go for alternate spraying of thiomethoxam 25WG @ 0.25 gm/lit or imidacloprid 0.45 ml/lit of water or Dinotefuron 20% SG 3gm/lit of Water.

PULSES:

This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

VEGETABLES: Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. To control **Fruit and shoot borer** in Brinjal and Okra, Spray Neem based pesticide 1500ppm @ 600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45% SC @ 70ml/acre and Thiodicarb @ 300g/acre at 15 days interval in 200 litre of water. To control **YMV in Okra, thrips in Chilli and aphids in Cowpea** crop, spray Imidacloprid @ 5 ml in 10 litres of water. **Fruits:** **Banana:** Present weather conditions are favourable for the incidence and spread of **sigatoka leaf spot** disease. To curb the spread of this disease, go for spraying of propiconazole @ 0.1% (or) tebuconazole + trifoxystrobin @ 1ml per litre for 2-3 times in 25 days interval. **Papaya:** Present weather conditions are favourable for the incidence of **foot rot** in Papaya. To control go for soil drenching around the base of the stem with Metalaxyl @ 2g/L. **Cashewnut:** Present weather conditions are favourable for the incidence of **tea mosquito bug**. To control go for spraying of Imidacloprid @ 5 ml in 10 litres of water at the time of sprouting. At the time of flowering, go for spraying of lambda cyhalothrin @ 0.6ml/l of water.

Animal husbandry: Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com



Bulletin No.-140
Date- 10/11/2020

AGROMET ADVISORY BULLETIN FOR MAHANGA

District Past week weather condition and Block past Rainfall condition

The block received 0 mm and district received 0 mm rainfall during the last week. The mean daily maximum and minimum temperature was 32°C and 18°C, respectively. Mean relative humidity of the district was around 70-66%. **According to NDTV map (28.10.2020 to 03.11.2020) the agriculture vigor was light to moderate over few places of Cuttack district whereas according to SPI value (cumulative 4 weeks-08 October 2020 to 04 November 2020), moderately wet condition was there.**

Forecasted period: 11th November 2020 to 15th November 2020

Weather advisory: No rainfall is forecasted for upcoming 5 days (Today to Saturday) in the block. The day temperature will be 31-32°C and night temperature will be 20-22°C for upcoming 5 days. Maximum and minimum relative humidity will be 58-74% and 38-51% respectively. There will be mainly clear to partly cloudy sky for next 5 days.

Block	Mahanga (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	11/11/2020	12/11/2020	13/11/2020	14/11/2020	15/11/2020
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	31	32	32	32	32
T-Min(°C)	21	22	21	21	20
Cloud Cover(octal)	0	1	6	4	1
RH Max (%)	74	69	58	64	68
RH Min (%)	50	51	41	38	41
Wind Speed (Kmph)	11	11	11	12	10
Wind Direction (deg)	45	45	45	32	23

Agromet Advisory

General Advisory: Farmers should go for harvesting of early matured paddy at 85% maturity of the panicles in sunny days. They should keep 5 cm water in paddy fields to maintain saturation during PI to grain filling stage. Carry out spraying of insecticides or pesticides in crops during morning and afternoon hours. Complete nursery sowing for winter vegetable crops. They should cover the nursery beds with polythene in night to protect it from cold temperature. Irrigate the vegetable nursery during morning and evening time as below normal rainfall likely to occur in Odisha in upcoming week. Go for sowing black gram and green gram in upland rice fallow condition by utilizing residual soil moisture.

Kharif Rice: Paddy is at flowering to milking stage. In Early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop. Current weather favors the infestation of **Leaf Folder**. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days. If infestation of **BPH & WBPH** occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 80g Thiamethoxam or 60g Flonicamid per acre in alternate basis. To prevent crop from **Gundhi bug** apply 5 ml neem based insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. Present weather conditions are favourable for the incidence of **Hispa**. To control, go for spraying of chlorpyrifos @ 2.5ml (or) profenophos @ 2ml /l. In **Gall midge** affected areas apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre or Chlorpyrifos 20EC @ 500ml/acre to manage the pest. For prevention of **false smut** spray the field before panicle initiation or flowering with propiconazole 25%EC @ 2ml/litre of water. Present weather conditions are favourable for incidence of **Bacterial leaf blight**. To control, go for spraying of plantomycin @ 1g/L or Streptocyclin 0.1g/L + COC 3g/L.

Sugarcane:

Remove the borer affected tillers and late formed tillers, tie the cane shoot with two or three together with partially dried lower leaves. Present weather conditions are favourable for the incidence of **ring spot disease** in Sugarcane. To control go for spraying of copperoxychloride @ 3g/l (or) Carbendazim @ 1g/l (or) Mancozeb @ 3g/l for three times once in three weeks. To control **top shoot borers** spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water. For control of Sucking pest in sugarcane like **Woolly aphid, Whitefly, Scale insects, mealybug** in present weather conditions go for alternate spraying of thiomethoxam 25WG @ 0.25 gm/lit or imidacloprid 0.45 ml/lit of water or Dinotefuron 20% SG 3gm/lit of Water.

PULSES:

This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

VEGETABLES: Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. To control **Fruit and shoot borer** in Brinjal and Okra, Spray Neem based pesticide 1500ppm @ 600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45% SC @ 70ml/acre and Thiodicarb @ 300g/acre at 15 days interval in 200 litre of water. To control **YMV in Okra, thrips in Chilli and aphids in Cowpea** crop, spray Imidacloprid @ 5 ml in 10 litres of water. **Fruits:** **Banana:** Present weather conditions are favourable for the incidence and spread of **sigatoka leaf spot** disease. To curb the spread of this disease, go for spraying of propiconazole @ 0.1% (or) tebuconazole + trifloxystrobin @ 1ml per litre for 2-3 times in 25 days interval. **Papaya:** Present weather conditions are favourable for the incidence of **foot rot** in Papaya. To control go for soil drenching around the base of the stem with Metalaxyl @ 2g/L. **Cashewnut:** Present weather conditions are favourable for the incidence of **tea mosquito bug**. To control go for spraying of Imidacloprid @ 5 ml in 10 litres of water at the time of sprouting. At the time of flowering, go for spraying of lambda cyhalothrin @ 0.6ml/l of water.

Animal husbandry:

Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com



Bulletin No.-140
Date- 10/11/2020

AGROMET ADVISORY BULLETIN FOR NARASINGHPUR
District Past week weather condition and Block past Rainfall condition

The block received 0 mm and district received 0 mm rainfall during the last week. The mean daily maximum and minimum temperature was 32°C and 18°C, respectively. Mean relative humidity of the district was around 70-66%. **According to NDMI map (28.10.2020 to 03.11.2020) the agriculture vigor was light to moderate over few places of Cuttack district whereas according to SPI value (cumulative 4 weeks-08 October 2020 to 04 November 2020), moderately wet condition was there.**

Forecasted period: 11th November 2020 to 15th November 2020

Weather advisory: No rainfall is forecasted for upcoming 5 days (Today to Saturday) in the block. The day temperature will be 31-32°C and night temperature will be 18-19°C for upcoming 5 days. Maximum and minimum relative humidity will be 65-86% and 36-49% respectively. There will be mainly clear to partly cloudy sky for next 5 days.

Block	Narasinghpur (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	11/11/2020	12/11/2020	13/11/2020	14/11/2020	15/11/2020
Rainfall (mm)	0	0	0	0	0
T-Max (°C)	31	32	32	32	32
T-Min (°C)	18	18	19	18	18
Cloud Cover (/octa)	1	2	6	4	0
RH Max (%)	86	82	65	65	73
RH Min (%)	46	49	41	36	39
Wind Speed (Kmph)	5	6	5	5	5
Wind Direction (deg)	101	108	75	75	68

Agromet Advisory

General Advisory: Farmers should go for harvesting of early matured paddy at 85% maturity of the panicles in sunny days. They should keep 5 cm water in paddy fields to maintain saturation during PI to grain filling stage. Carry out spraying of insecticides or pesticides in crops during morning and afternoon hours. Complete nursery sowing for winter vegetable crops. They should cover the nursery beds with polythene in night to protect it from cold temperature. Irrigate the vegetable nursery during morning and evening time as below normal rainfall likely to occur in Odisha in upcoming week. Go for sowing black gram and green gram in upland rice fallow condition by utilizing residual soil moisture.

Kharif Rice:

Paddy is at flowering to milking stage. In Early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop. Current weather favors the infestation of **Leaf Folder**. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days. If infestation of **BPH & WBPH** occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 80g Thiamethoxam or 60g Flonicamid per acre in alternate basis. To prevent crop from **Gundhi bug** apply 5 ml neem based insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. Present weather conditions are favourable for the incidence of **Hispa**. To control, go for spraying of chloropyrifos @ 2.5ml (or) profenophos @ 2ml /l. In **Gall midge** affected areas apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre or Chlorpyrifos 20EC @ 500ml/acre to manage the pest. For prevention of **false smut** spray the field before panicle initiation or flowering with propiconazole 25% EC @ 2ml/liter of water. Present weather conditions are favourable for incidence of **Bacterial leaf blight**. To control, go for spraying of plantomycin @ 1g/L or Streptocyclin 0.1g/L + COC 3g/L.

PULSES:

This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

Mustard:

Taking the advantage of available soil moisture due to recent rain, land preparation and sowing of the crop should be started to get optimum yield. Suitable varieties: Toria- Anuradha, Parbati, Sushree, Indian mustard- Mahyco bold, JD 6. Seed rate- 8-10 Kg/ha Spacing- 30 cm X 10 cm, Fertilizer dose: Toria @ 60:30:30 N-P2O5-K2O Kg/ha Mustard @ 80:40:40 N-P2O5-K2O Kg/ha. Apply 50%N and all P2O5 and K2O fertilizers should be applied as basal.

VEGETABLES:

Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. To control **Fruit and shoot borer** in Brinjal and Okra, Spray Neem based pesticide 1500ppm @ 600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45% SC @ 70ml/acre and Thiodicarb @ 300g/acre at 15 days interval in 200 litre of water. To control **YMV in Okra, thrips in Chilli and aphids in Cowpea** crop, spray Imidacloprid @ 5 ml in 10 litres of water. **Fruits: Banana:** Present weather conditions are favourable for the incidence and spread of **sigatoka leaf spot** disease. To curb the spread of this disease, go for spraying of propiconazole @ 0.1% (or) tebuconazole + trifloxystrobin @ 1ml per litre for 2-3 times in 25 days interval. **Papaya:** Present weather conditions are favourable for the incidence of **foot rot** in Papaya. To control go for soil drenching around the base of the stem with Metalaxyl @ 2g/L. **Cashewnut:** Present weather conditions are favourable for the incidence of **tea mosquito bug**. To control go for spraying of Imidacloprid @ 5 ml in 10 litres of water at the time of sprouting. At the time of flowering, go for spraying of lambda cyhalothrin @ 0.6ml/l of water.

Animal husbandry: Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com



Bulletin No.-140
Date- 10/11/2020

AGROMET ADVISORY BULLETIN FOR NIALI

District Past week weather condition and Block past Rainfall condition

The block received 0 mm and district received 0 mm rainfall during the last week. The mean daily maximum and minimum temperature was 32°C and 18°C, respectively. Mean relative humidity of the district was around 70-66%. **According to NDVI map (28.10.2020 to 03.11.2020) the agriculture vigor was light to moderate over few places of Cuttack district whereas according to SPI value (cumulative 4 weeks-08 October 2020 to 04 November 2020), moderately wet condition was there.**

Forecasted period: 11th November 2020 to 15th November 2020

Weather advisory: No rainfall is forecasted for upcoming 5 days (Today to Saturday) in the block. The day temperature will be 31-33°C and night temperature will be 20-22°C for upcoming 5 days. Maximum and minimum relative humidity will be 63-78% and 39-54% respectively. There will be mainly clear to partly cloudy sky for next 5 days.

Block	Niali (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	11/11/2020	12/11/2020	13/11/2020	14/11/2020	15/11/2020
Rainfall(mm)	0	1	0	0	0
T-Max(°C)	31	32	33	32	32
T-Min(°C)	22	22	21	21	20
Cloud Cover(octa)	0	2	4	1	2
RH Max (%)	78	74	63	66	69
RH Min (%)	53	54	43	39	42
Wind Speed(Kmph)	12	12	12	13	12
Wind Direction(deg)	45	45	45	32	54

Agromet Advisory

General Advisory: Farmers should go for harvesting of early matured paddy at 85% maturity of the panicles in sunny days. They should keep 5 cm water in paddy fields to maintain saturation during PI to grain filling stage. Carry out spraying of insecticides or pesticides in crops during morning and afternoon hours. Complete nursery sowing for winter vegetable crops. They should cover the nursery beds with polythene in night to protect it from cold temperature. Irrigate the vegetable nursery during morning and evening time as below normal rainfall likely to occur in Odisha in upcoming week. Go for sowing black gram and green gram in upland rice fallow condition by utilizing residual soil moisture.

Kharif Rice: Paddy is at flowering to milking stage. In Early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop. Current weather favors the infestation of **Leaf Folder**. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days. If infestation of **BPH & WBPH** occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 80g Thiamethoxam or 60g Flonicamid per acre in alternate basis. To prevent crop from **Gundhi bug** apply 5 ml neem based insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. Present weather conditions are favourable for the incidence of **Hispa**. To control, go for spraying of chlorpyrifos @ 2.5ml (or) profenophos @ 2ml /l. In **Gall midge** affected areas apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre or Chlorpyrifos 20EC @ 500ml/acre to manage the pest. For prevention of **false smut** spray the field before panicle initiation or flowering with propiconazole 25%EC @ 2ml/liter of water. Present weather conditions are favourable for incidence of **Bacterial leaf blight**. To control, go for spraying of plantomycin @ 1g/L or Streptocyclin 0.1g/L + COC 3g/L.

Sugarcane:

Remove the borer affected tillers and late formed tillers, tie the cane shoot with two or three together with partially dried lower leaves. Present weather conditions are favourable for the incidence of **ring spot disease** in Sugarcane. To control go for spraying of copperoxychloride @ 3g/l (or) Carbenazim @ 1g/l (or) Mancozeb @ 3g/l for three times once in three weeks. To control **top shoot borers** spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water. For control of Sucking pest in sugarcane like **Woolly aphid, Whitefly, Scale insects, mealybug** in present weather conditions go for alternate spraying of thiomethoxam 25WG @ 0.25 gm/lit or imidacloprid 0.45 ml/lit of water or Dinotefuron 20% SG 3gm/lit of Water.

PULSES:

This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

VEGETABLES: Go for early planting of cole crops like radish, cabbage, cauliflower, beans, cluster bean and tomato. To control **Fruit and shoot borer** in Brinjal and Okra, Spray Neem based pesticide 1500ppm @ 600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45% SC @ 70ml/acre and Thiodicarb @ 300g/acre at 15 days interval in 200 litre of water. To control **YMV in Okra, thrips in Chilli and aphids in Cowpea** crop, spray Imidacloprid @ 5 ml in 10 litres of water.

Fruits: **Banana:** Present weather conditions are favourable for the incidence and spread of **sigatoka leaf spot** disease. To curb the spread of this disease, go for spraying of propiconazole @ 0.1% (or) tebuconazole + trifloxystrobin @ 1ml per litre for 2 -3 times in 25 days interval. **Papaya:** Present weather conditions are favourable for the incidence of **foot rot** in Papaya. To control go for soil drenching around the base of the stem with Metalaxyl @ 2g/L. **Cashewnut:** Present weather conditions are favourable for the incidence of **tea mosquito bug**. To control go for spraying of Imidacloprid @ 5 ml in 10 litres of water at the time of sprouting. At the time of flowering, go for spraying of lambda cyhalothrin @ 0.6ml/l of water.

Animal husbandry:

Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com



Bulletin No.-140
Date- 10/11/2020

AGROMET ADVISORY BULLETIN FOR NISCHINTAKOILI
District Past week weather condition and Block past Rainfall condition

The block received 0 mm and district received 0 mm rainfall during the last week. The mean daily maximum and minimum temperature was 32°C and 18°C, respectively. Mean relative humidity of the district was around 70-66%. **According to NDVI map (28.10.2020 to 03.11.2020) the agriculture vigor was light to moderate over few places of Cuttack district whereas according to SPI value (cumulative 4 weeks-08 October 2020 to 04 November 2020), moderately wet condition was there.**

Forecasted period: 11th November 2020 to 15th November 2020

Weather advisory: No rainfall is forecasted for upcoming 5 days (Today to Saturday) in the block. The day temperature will be 31-32°C and night temperature will be 20-22°C for upcoming 5 days. Maximum and minimum relative humidity will be 59-75% and 38-52% respectively. There will be mainly clear to partly cloudy sky for next 5 days.

Block	Nischintakoili (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	11/11/2020	12/11/2020	13/11/2020	14/11/2020	15/11/2020
Rainfall (mm)	0	0	0	0	0
T-Max (°C)	31	32	32	32	32
T-Min (°C)	22	22	21	21	20
Cloud Cover (octa)	0	1	6	4	2
RH Max (%)	75	69	59	65	69
RH Min (%)	51	52	41	38	41
Wind Speed (Kmph)	11	12	11	13	11
Wind Direction (deg)	45	45	45	32	23

Agromet Advisory

General Advisory: Farmers should go for harvesting of early matured paddy at 85% maturity of the panicles in sunny days. They should keep 5 cm water in paddy fields to maintain saturation during PI to grain filling stage. Carry out spraying of insecticides or pesticides in crops during morning and afternoon hours. Complete nursery sowing for winter vegetable crops. They should cover the nursery beds with polythene in night to protect it from cold temperature. Irrigate the vegetable nursery during morning and evening time as below normal rainfall likely to occur in Odisha in upcoming week. Go for sowing black gram and green gram in upland rice fallow condition by utilizing residual soil moisture.

Kharif Rice: Paddy is at flowering to milking stage. In Early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop. Current weather favors the infestation of **Leaf Folder**. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days. If infestation of **BPH & WBPH** occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 80g Thiamethoxam or 60g Flonicamid per acre in alternate basis. To prevent crop from **Gundhi bug** apply 5 ml neem based insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. Present weather conditions are favourable for the incidence of **Hispa**. To control, go for spraying of chlorpyrifos @ 2.5ml (or) profenophos @ 2ml /l. In **Gall midge** affected areas apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre or Chlorpyrifos 20EC @ 500ml/acre to manage the pest. For prevention of **false smut** spray the field before panicle initiation or flowering with propiconazole 25% EC @ 2ml/liter of water. Present weather conditions are favourable for incidence of **Bacterial leaf blight**. To control, go for spraying of plantomycin @ 1g/L or Streptocyclin 0.1g/L + COC 3g/L.

Sugarcane:

Remove the borer affected tillers and late formed tillers, tie the cane shoot with two or three together with partially dried lower leaves. Present weather conditions are favourable for the incidence of **ring spot disease** in Sugarcane. To control go for spraying of copperoxychloride @ 3g/l (or) Carbendazim @ 1g/l (or) Mancozeb @ 3g/l for three times once in three weeks. To control **top shoot borers** spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water. For control of Sucking pest in sugarcane like **Woolly aphid, Whitefly, Scale insects, mealybug** in present weather conditions go for alternate spraying of thiomethoxam 25WG @ 0.25 gm/lit or imidacloprid 0.45 ml/lit of water or Dinotefuron 20% SG 3gm/lit of Water.

PULSES:

This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

VEGETABLES: Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. To control **Fruit and shoot borer** in Brinjal and Okra, Spray Neem based pesticide 1500ppm @ 600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45% SC @ 70ml/acre and Thiodicarb @ 300g/acre at 15 days interval in 200 litre of water. To control **YMV in Okra, thrips in Chilli and aphids in Cowpea** crop, spray Imidacloprid @ 5 ml in 10 litres of water.

Fruits: **Banana:** Present weather conditions are favourable for the incidence and spread of **sigatoka leaf spot** disease. To curb the spread of this disease, go for spraying of propiconazole @ 0.1% (or) tebuconazole + trifoxystrobin @ 1ml per litre for 2 -3 times in 25 days interval. **Papaya:** Present weather conditions are favourable for the incidence of **foot rot** in Papaya. To control go for soil drenching around the base of the stem with Metalaxyl @ 2g/L. **Cashewnut:** Present weather conditions are favourable for the incidence of **tea mosquito bug**. To control go for spraying of Imidacloprid @ 5 ml in 10 litres of water at the time of sprouting. At the time of flowering, go for spraying of lambda cyhalothrin @ 0.6ml/l of water.

Animal husbandry:

Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com



Bulletin No.-140
Date- 10/11/2020

AGROMET ADVISORY BULLETIN FOR SALEPUR

District Past week weather condition and Block past Rainfall condition

The block received 0 mm and district received 0 mm rainfall during the last week. The mean daily maximum and minimum temperature was 32°C and 18°C, respectively. Mean relative humidity of the district was around 70-66%. According to NDVI map (28.10.2020 to 03.11.2020) the agriculture vigor was light to moderate over few places of Cuttack district whereas according to SPI value (cumulative 4 weeks-08 October 2020 to 04 November 2020), moderately wet condition was there.

Forecasted period: 11th November 2020 to 15th November 2020

Weather advisory: No rainfall is forecasted for upcoming 5 days (Today to Saturday) in the block. The day temperature will be 31-32°C and night temperature will be 20-22°C for upcoming 5 days. Maximum and minimum relative humidity will be 59-74% and 39-52% respectively. There will be mainly clear to partly cloudy sky for next 5 days.

Block	Salepur (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	11/11/2020	12/11/2020	13/11/2020	14/11/2020	15/11/2020
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	31	32	32	32	32
T-Min(°C)	22	22	21	21	20
Cloud Cover(octa)	0	1	6	4	1
RH Max (%)	74	69	59	65	69
RH Min (%)	50	52	41	39	41
Wind Speed (Kmph)	11	11	11	12	10
Wind Direction (deg)	45	45	45	32	23

Agromet Advisory

General Advisory: Farmers should go for harvesting of early matured paddy at 85% maturity of the panicles in sunny days. They should keep 5 cm water in paddy fields to maintain saturation during PI to grain filling stage. Carry out spraying of insecticides or pesticides in crops during morning and afternoon hours. Complete nursery sowing for winter vegetable crops. They should cover the nursery beds with polythene in night to protect it from cold temperature. Irrigate the vegetable nursery during morning and evening time as below normal rainfall likely to occur in Odisha in upcoming week. Go for sowing black gram and green gram in upland rice fallow condition by utilizing residual soil moisture.

Kharif Rice:

Paddy is at flowering to milking stage. In Early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop. Current weather favors the infestation of **Leaf Folder**. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days. If infestation of **BPH & WBPH** occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 80g Thiamethoxam or 60g Flonicamid per acre in alternate basis. To prevent crop from **Gundhi bug** apply 5 ml neem based insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. Present weather conditions are favourable for the incidence of **Hispa**. To control, go for spraying of chloropyrifos @ 2.5ml (or) profenophos @ 2ml /l. In **Gall midge** affected areas apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre or Chlorpyrifos 20EC @ 500ml/acre to manage the pest. For prevention of **false smut** spray the field before panicle initiation or flowering with propiconazole 25%EC @ 2ml/litre of water. Present weather conditions are favourable for incidence of **Bacterial leaf blight**. To control, go for spraying of plantomycin @ 1g/L or Streptocyclin 0.1g/L + COC 3g/L.

PULSES:

This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

Mustard: Taking the advantage of available soil moisture due to recent rain, land preparation and sowing of the crop should be started to get optimum yield. Suitable varieties: Toria- Anuradha, Parbati, Sushree, Indian mustard- Mahyco bold, JD 6. Seed rate- 8-10 Kg/ha Spacing- 30 cm X 10 cm, Fertilizer dose: Toria @ 60:30:30 N-P2O5-K2O Kg/ha Mustard @ 80:40:40 N-P2O5-K2O Kg/ha. Apply 50%N and all P2O5 and K2O fertilizers should be applied as basal.

VEGETABLES:

Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. To control **Fruit and shoot borer** in Brinjal and Okra, Spray Neem based pesticide 1500ppm @ 600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45% SC @ 70ml/acre and Thiodicarb @ 300g/acre at 15 days interval in 200 litre of water. To control **YMV in Okra, thrips in Chilli and aphids in Cowpea** crop, spray Imidacloprid @ 5 ml in 10 litres of water. **Fruits:** **Banana:** Present weather conditions are favourable for the incidence and spread of **sigatoka leaf spot** disease. To curb the spread of this disease, go for spraying of propiconazole @ 0.1% (or) tebuconazole + trifloxystrobin @ 1ml per litre for 2 -3 times in 25 days interval. **Papaya:** Present weather conditions are favourable for the incidence of **foot rot** in Papaya. To control go for soil drenching around the base of the stem with Metalaxyl @ 2g/L. **Cashewnut:** Present weather conditions are favourable for the incidence of **tea mosquito bug**. To control go for spraying of Imidacloprid @ 5 ml in 10 litres of water at the time of sprouting. At the time of flowering, go for spraying of lambda cyhalothrin @ 0.6ml/l of water.

Animal husbandry:

Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com



Bulletin No.-140
Date- 10/11/2020

AGROMET ADVISORY BULLETIN FOR TANGI-CHOUDWAR
District Past week weather condition and Block past Rainfall condition

The block received 0 mm and district received 0 mm rainfall during the last week. The mean daily maximum and minimum temperature was 32°C and 18°C, respectively. Mean relative humidity of the district was around 70-66%. **According to NDVI map (28.10.2020 to 03.11.2020) the agriculture vigor was light to moderate over few places of Cuttack district whereas according to SPI value (cumulative 4 weeks-08 October 2020 to 04 November 2020), moderately wet condition was there.**

Forecasted period: 11th November 2020 to 15th November 2020

Weather advisory: No rainfall is forecasted for upcoming 5 days (Today to Saturday) in the block. The day temperature will be 30-31°C and night temperature will be 22-24°C for upcoming 5 days. Maximum and minimum relative humidity will be 64-78% and 42-59% respectively. There will be mainly clear to partly cloudy sky for next 5 days.

Block	Tangi-Choudwar (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	11/11/2020	12/11/2020	13/11/2020	14/11/2020	15/11/2020
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	30	31	31	31	31
T-Min(°C)	24	24	23	23	22
Cloud Cover(octa)	0	2	2	1	2
RH Max (%)	78	71	64	66	69
RH Min (%)	57	59	47	42	46
Wind Speed (Kmph)	14	14	14	16	14
Wind Direction (deg)	58	58	32	27	54

Agromet Advisory

General Advisory: Farmers should go for harvesting of early matured paddy at 85% maturity of the panicles in sunny days. They should keep 5 cm water in paddy fields to maintain saturation during PI to grain filling stage. Carry out spraying of insecticides or pesticides in crops during morning and afternoon hours. Complete nursery sowing for winter vegetable crops. They should cover the nursery beds with polythene in night to protect it from cold temperature. Irrigate the vegetable nursery during morning and evening time as below normal rainfall likely to occur in Odisha in upcoming week. Go for sowing black gram and green gram in upland rice fallow condition by utilizing residual soil moisture.

Kharif Rice:

Paddy is at flowering to milking stage. In Early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop. Current weather favors the infestation of **Leaf Folder**. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days. If infestation of **BPH & WBPH** occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 80g Thiamethoxam or 60g Flonicamid per acre in alternate basis. To prevent crop from **Gundhi bug** apply 5 ml neem based insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. Present weather conditions are favourable for the incidence of **Hispa**. To control, go for spraying of chloropyrifos @ 2.5ml (or) profenophos @ 2ml /l. In **Gall midge** affected areas apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre or Chlorpyrifos 20EC @ 500ml/acre to manage the pest. For prevention of **false smut** spray the field before panicle initiation or flowering with propiconazole 25%EC @ 2ml/liter of water. Present weather conditions are favourable for incidence of **Bacterial leaf blight**. To control, go for spraying of plantomycin @ 1g/L or Streptocyclin 0.1g/L + COC 3g/L.

PULSES:

This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

Mustard:

Taking the advantage of available soil moisture due to recent rain, land preparation and sowing of the crop should be started to get optimum yield. Suitable varieties: Toria- Anuradha, Parbati, Sushree, Indian mustard- Mahyco bold, JD 6. Seed rate- 8-10 Kg/ha Spacing- 30 cm X 10 cm, Fertilizer dose: Toria @ 60:30:30 N-P2O5-K2O Kg/ha Mustard @ 80:40:40 N-P2O5-K2O Kg/ha. Apply 50%N and all P2O5 and K2O fertilizers should be applied as basal.

VEGETABLES:

Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. To control **Fruit and shoot borer** in Brinjal and Okra, Spray Neem based pesticide 1500ppm @ 600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45% SC @ 70ml/acre and Thiodicarb @ 300g/acre at 15 days interval in 200 litre of water. To control **YMV in Okra, thrips in Chilli and aphids in Cowpea** crop, spray Imidacloprid @ 5 ml in 10 litres of water. **Fruits:** **Banana:** Present weather conditions are favourable for the incidence and spread of **sigatoka leaf spot** disease. To curb the spread of this disease, go for spraying of propiconazole @ 0.1% (or) tebuconazole + trifloxystrobin @ 1ml per litre for 2 -3 times in 25 days interval. **Papaya:** Present weather conditions are favourable for the incidence of **foot rot** in Papaya. To control go for soil drenching around the base of the stem with Metalaxyl @ 2g/L. **Cashewnut:** Present weather conditions are favourable for the incidence of **tea mosquito bug**. To control go for spraying of Imidacloprid @ 5 ml in 10 litres of water at the time of sprouting. At the time of flowering, go for spraying of lambda cyhalothrin @ 0.6ml/l of water.

Animal husbandry:

Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com



Bulletin No.-140
Date- 10/11/2020

AGROMET ADVISORY BULLETIN FOR TIGIRIA

District Past week weather condition and Block past Rainfall condition

The block received 0 mm and district received 0 mm rainfall during the last week. The mean daily maximum and minimum temperature was 32°C and 18°C, respectively. Mean relative humidity of the district was around 70-66%. **According to NDVI map (28.10.2020 to 03.11.2020) the agriculture vigor was light to moderate over few places of Cuttack district whereas according to SPI value (cumulative 4 weeks-08 October 2020 to 04 November 2020), moderately wet condition was there.**

Forecasted period: 11th November 2020 to 15th November 2020

Weather advisory: No rainfall is forecasted for upcoming 5 days (Today to Saturday) in the block. The day temperature will be 32-33°C and night temperature will be 19-20°C for upcoming 5 days. Maximum and minimum relative humidity will be 63-79% and 37-49% respectively. There will be mainly clear to partly cloudy sky for next 5 days.

Block	Tigiria (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	11/11/2020	12/11/2020	13/11/2020	14/11/2020	15/11/2020
Rainfall(mm)	0	0	0	0	0
00T-Max(°C)	32	32	33	33	33
T-Min(°C)	19	20	20	19	19
Cloud Cover(octa)	1	2	6	4	0
RH Max (%)	79	72	63	68	73
RH Min (%)	49	49	40	37	39
Wind Speed (Kmph)	7	8	7	7	7
Wind Direction (deg)	101	60	68	23	90

Agromet Advisory

General Advisory: Farmers should go for harvesting of early matured paddy at 85% maturity of the panicles in sunny days. They should keep 5 cm water in paddy fields to maintain saturation during PI to grain filling stage. Carry out spraying of insecticides or pesticides in crops during morning and afternoon hours. Complete nursery sowing for winter vegetable crops. They should cover the nursery beds with polythene in night to protect it from cold temperature. Irrigate the vegetable nursery during morning and evening time as below normal rainfall likely to occur in Odisha in upcoming week. Go for sowing black gram and green gram in upland rice fallow condition by utilizing residual soil moisture.

Kharif Rice:

Paddy is at flowering to milking stage. In Early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop. Current weather favors the infestation of **Leaf Folder**. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days. If infestation of **BPH & WBPH** occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 80g Thiamethoxam or 60g Flonicamid per acre in alternate basis. To prevent crop from **Gundhi bug** apply 5 ml neem based insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. Present weather conditions are favourable for the incidence of **Hispa**. To control, go for spraying of chloropyrifos @2.5ml (or) profenophos @ 2ml /l. In **Gall midge** affected areas apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre or Chlorpyrifos 20EC @500ml/acre to manage the pest. For prevention of **false smut** spray the field before panicle initiation or flowering with propiconazole 25%EC @ 2ml/litre of water. Present weather conditions are favourable for incidence of **Bacterial leaf blight**. To control, go for spraying of plantomycin @1g/L or Streptocyclin 0.1g/L + COC 3g/L.

PULSES:

This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

Mustard:

Taking the advantage of available soil moisture due to recent rain, land preparation and sowing of the crop should be started to get optimum yield. Suitable varieties: Toria- Anuradha, Parbati, Sushree, Indian mustard- Mahyco bold, JD 6. Seed rate- 8-10 Kg/ha Spacing- 30 cm X 10 cm, Fertilizer dose: Toria @60:30:30 N-P2O5-K2O Kg/ha Mustard @ 80:40:40 N-P2O5-K2O Kg/ha. Apply 50%N and all P2O5 and K2O fertilizers should be applied as basal.

VEGETABLES:

Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. To control **Fruit and shoot borer** in Brinjal and Okra, Spray Neem based pesticide 1500ppm @600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45% SC @70ml/acre and Thiodicarb @300g/acre at 15 days interval in 200 litre of water. To control **YMV in Okra, thrips in Chilli and aphids in Cowpea** crop, spray Imidacloprid @ 5 ml in 10 litres of water. **Fruits: Banana:** Present weather conditions are favourable for the incidence and spread of **sigatoka leaf spot** disease. To curb the spread of this disease, go for spraying of propiconazole @ 0.1% (or) tebuconazole + trifloxystrobin @ 1ml per litre for 2 -3 times in 25 days interval. **Papaya:** Present weather conditions are favourable for the incidence of **foot rot** in Papaya. To control go for soil drenching around the base of the stem with Metalaxyl @ 2g/L. **Cashewnut:** Present weather conditions are favourable for the incidence of **tea mosquito bug**. To control go for spraying of Imidacloprid @ 5 ml in 10 litres of water at the time of sprouting. At the time of flowering, go for spraying of lambda cyhalothrin @ 0.6ml/l of water.

Animal husbandry:

Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427