




GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com




Bulletin No.-243
Date- 05/11/2021

AGROCLIMATIC ZONES OF CUTTACK DISTRICT:



STANDARD PRECIPITATION INDEX AND NDVI MAP



LEGEND (Medium range block level rainfall forecast)

	No Rainfall	00 mm
	Very light rainfall	01-05 mm
	Light rainfall	06-10 mm
	Light to moderate rainfall	11-30 mm
	Moderate to heavy rainfall	31-65 mm
	Heavy to very heavy rainfall	66-115 mm
	Extremely heavy rainfall	>116 mm

Dr Sujata Sethy
 (Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com




Bulletin No.-243
 Date- 05/11/2021

Past week (29 October-04 November 2021) Cuttack district Cumulative Rainfall departure (%) and average Temperature



Medium range (Today (8.30IST) to Wednesday (8.30IST)) Rainfall (mm) forecast for Cuttack district



Dr Sujata Sethy
 (Nodal officer)



**Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area**



Debasish Jena
 SMS (Agrometeorology)
 Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-243
Date- 05/11/2021


Met Sub division	Extended range outlook for subsequent 1 week (6 th to 12 th days) (10-16 November 2021)		
Odisha	<i>Parameter</i>	<i>Departure (%)</i>	<i>Category</i>
	Rainfall	23.38	Above Normal
	<i>Parameter</i>	<i>Anomaly</i>	<i>Category</i>
	Maximum temperature	0.65	Normal
	Minimum temperature	-0.26	Normal

General Agromet Advisory:- There will be above normal rainfall in subsequent week after 5 days (10-16 November 2021) in Odisha condition, and there will be dry weather is forecasted for upcoming 5 days. Farmers are advised to conserve water level of 5cm during panicle initiation to flowering stage of rice. They may go for harvesting of early sown rice if they are at 85% physiological maturity stage before the subsequent week. Farmers may go for weeding, hoeing followed by earthing up and intercultural operations in the dry days in vegetable fields.

South-west Monsoon has withdrawn from entire Odisha, from 23rd October, 2021

Post Monsoon rainfall forecast for the month of November, 2021

- The spatial distribution of probabilistic forecasts for tercile categories (above normal, normal and below normal) over the country for the November rainfall is shown in Fig. The spatial distribution suggests that **Normal to above normal** is likely over many parts of Odisha.
- Hence Farmers should manage the crop and irrigation activities depending upon the land type, irrigation facility for kharif rice and pre Rabi pulses and vegetables.*



Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427

















GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-243
Date- 05/11/2021

Weather (Past and Forecasted) based Disease and pest in Major Crops and vegetables

Sl No	Crop	Stage and condition of the crop	Anticipated Disease and pest	
1	Rice	Panicle initiation to Flowering stage		Leaf folder
				BPH
				Gundhi bug
				False smut
				Bacterial leaf streak
2	Sugarcane	Grand growth stage to cane development stag		Ring spot
				Top borer
				Woolly Aphid
				Scale insect
				White fly
				Mealy bug
3	Vegetables(Brinjal and okra)	Vegetative to fruiting		Fruit and shoot borer
	Okra			YMV
	Chilli , Cowpea			Thrips in Chilli and Aphids in Cowpea

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-243
Date- 05/11/2021

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31°C and 21°C, respectively. Mean relative humidity of the district was around 81-71%.

Forecasted period: Today(8.30IST) to 10th November (8.30IST)2021

Weather advisory: Dry weather is forecasted for today to Tuesday in the block. The day temperature will be 30-31°C and night temperature will be 19-20°C for upcoming 5 days. Maximum and minimum relative humidity will be 71-87% and 34-43% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Athagad(VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	06/11/2021	07/11/2021	08/11/2021	09/11/2021	10/11/2021
Rainfall (mm)	0	0	0	0	0
T-Max (°C)	31	30	30	31	31
T-Min (°C)	20	19	19	19	20
Cloud Cover (octa)	2	3	2	0	0
RH Max (%)	87	74	73	71	79
RH Min (%)	43	36	36	34	35
Wind Speed (kmph)	6	6	5	6	5
Wind Direction (deg)	11	345	345	335	349

Agromet Advisory

General Advisory: Farmers should go for harvesting of early matured paddy at 85% maturity of the panicles in sunny days. They should keep 5 cm water in paddy fields to maintain saturation during PI to grain filling stage. Carry out spraying of insecticides or pesticides in crops during morning and afternoon hours. Keep watch on the infestation of gundhi bug and BPH in paddy. Complete nursery sowing for winter vegetable crops. Irrigate the vegetable nursery during morning and evening time. Go for sowing black gram and green gram in upland rice fallow condition.

Kharif Rice:

Paddy is at PI to flowering stage. As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop. Current weather favors the infestation of **Leaf Folder**. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days. If infestation of **BPH & WBPH** occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 40g Thiamethoxam or 60g Flonicamid per acre in alternate basis. To prevent crop from **Gundhi bug** apply 5 ml neem based insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. For prevention of **false smut** spray the field before panicle initiation or flowering with propiconazole 25%EC @ 2ml/liter of water. In the event of **bacterial leaf streak** i.e. when the leaves are brown and dry in paddy crop, use plantomycin 1 g + copper oxychloride 1g per liter water only in the clear weather condition.

PULSES:

This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

GROUNDNUT:

Harvesting of Groundnut should be done if 75-80 % of pods are fully matured. The important indication of maturity is yellowing of foliage and necrotic spotting of leaves and dropping of older leaves.

Mustard:

Taking the advantage of available soil moisture due to recent rain, land preparation and sowing of the crop should be started to get optimum yield. Suitable varieties: Toria- Anuradha, Parbati, Sushree, Indian mustard- Mahyco bold, JD 6. Seed rate- 8-10 Kg/ha Spacing- 30 cm X 10 cm, Fertilizer dose: Toria @60:30:30 N-P2O5-K2O Kg/ha Mustard @ 80:40:40 N-P2O5-K2O Kg/ha. Apply 50%N and all P2O5 and K2O fertilizers should be applied as basal.

VEGETABLES:

Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. In case of Cauliflower in this dry days before transplanting during final land preparation apply 8-10 ton FYM per acre and fertilizer like N:P:K each 25-30 Kg/acre should be applied. Short duration Cauliflower varieties (5-6weeks) should be transplanted at distance 60cm×30cm and medium and late duration varieties (3-4 week seedlings) should be transplanted at 60-75cm×40-45cm

- To control Fruit and shoot borer in Brinjal and Okra, Spray Neem based pesticide 1500ppm @600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45% SC @70ml/acre and Thiodicarb @300g/acre at 15 days interval in 200 litre of water. To control YMV in Okra, thrips in Chilli and aphids in Cowpea crop, spray Imidacloprid @ 5 ml in 10 litres of water.

Animal husbandry:

Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-243
Date- 05/11/2021

AGROMET ADVISORY BULLETIN FOR BADAMBA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31°C and 21°C, respectively. Mean relative humidity of the district was around 81-71%.

Forecasted period: Today(8.30IST) to 10th November (8.30IST)2021

Weather advisory: Dry weather is forecasted for today to Tuesday in the block. The day temperature will be 30-31°C and night temperature will be 18-20°C for upcoming 5 days. Maximum and minimum relative humidity will be 66-90% and 33-46% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Badamba (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	06/11/2021	07/11/2021	08/11/2021	09/11/2021	10/11/2021
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	30	30	30	31	31
T-Min(°C)	20	19	18	18	19
Cloud Cover(octa)	2	3	2	0	0
RH Max (%)	90	75	71	66	77
RH Min (%)	46	37	36	33	34
Wind Speed (Kmph)	5	6	5	6	5
Wind Direction (deg)	11	15	360	343	360

Agromet Advisory

General Advisory: Farmers should go for harvesting of early matured paddy at 85% maturity of the panicles in sunny days. They should keep 5 cm water in paddy fields to maintain saturation during PI to grain filling stage. Carry out spraying of insecticides or pesticides in crops during morning and afternoon hours. Keep watch on the infestation of gundhi bug and BPH in paddy. Complete nursery sowing for winter vegetable crops. Irrigate the vegetable nursery during morning and evening time. Go for sowing black gram and green gram in upland rice fallow condition.

Kharif Rice:

Paddy is at PI to flowering stage. As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop. Current weather favors the infestation of Leaf Folder. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days. If infestation of BPH & WBPH occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 40g Thiamethoxam or 60g Flonicamid per acre in alternate basis. To prevent crop from Gundhi bug apply 5 ml neem based insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. For prevention of false smut spray the field before panicle initiation or flowering with propiconazole 25%EC @ 2ml/liter of water. In the event of bacterial leaf streak i.e. when the leaves are brown and dry in paddy crop, use plantomycin 1 g + cupper oxychloride 1g per liter water only in the clear weather condition.

PULSES:

This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

GROUNDNUT:

Harvesting of Groundnut should be done if 75-80 % of pods are fully matured. The important indication of maturity is yellowing of foliage and necrotic spotting of leaves and dropping of older leaves.

Mustard:

Taking the advantage of available soil moisture due to recent rain, land preparation and sowing of the crop should be started to get optimum yield. Suitable varieties: Toria- Anuradha, Parbati, Sushree, Indian mustard- Mahyco bold, JD 6. Seed rate- 8-10 Kg/ha Spacing- 30 cm X 10 cm, Fertilizer dose: Toria @60:30:30 N-P2O5-K2O Kg/ha Mustard @ 80:40:40 N-P2O5-K2O Kg/ha. Apply 50%N and all P2O5 and K2O fertilizers should be applied as basal.

VEGETABLES:

Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. In case of Cauliflower in this dry days before transplanting during final land preparation apply 8-10 ton FYM per acre and fertilizer like N:P:K each 25-30 Kg/acre should be applied. Short duration Cauliflower varieties (5-6weeks) should be transplanted at distance 60cm×30cm and medium and late duration varieties (3-4 week seedlings) should be transplanted at 60-75cm×40-45cm

- To control Fruit and shoot borer in Brinjal and Okra, Spray Neem based pesticide 1500ppm @600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45% SC @70ml/acre and Thiodicarb@300g/acre at 15 days interval in 200 litre of water. To control YMV in Okra, thrips in Chilli and aphids in Cowpea crop, spray Imidacloprid @ 5 ml in 10 litres of water.

Animal husbandry:

Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-243
 Date- 05/11/2021

AGROMET ADVISORY BULLETIN FOR BANKI

District Past week average weather condition

The district received average 0 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31°C and 21°C, respectively. Mean relative humidity of the district was around 81-71%.

Forecasted period: Today(8.30IST) to 10th November (8.30IST)2021

Weather advisory: Dry weather is forecasted for today to Tuesday in the block. The day temperature will be 30-31°C and night temperature will be 18-20°C for upcoming 5 days. Maximum and minimum relative humidity will be 68-89% and 34-44% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Banki (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	06/11/2021	07/11/2021	08/11/2021	09/11/2021	10/11/2021
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	31	30	31	31	31
T-Min(°C)	20	19	19	18	19
Cloud Cover(octa)	2	3	2	0	0
RH Max (%)	89	75	72	68	79
RH Min (%)	44	37	36	34	34
Wind Speed (Kmph)	6	6	6	6	6
Wind Direction (deg)	360	11	360	345	360

Agromet Advisory

General Advisory: Farmers should go for harvesting of early matured paddy at 85% maturity of the panicles in sunny days. They should keep 5 cm water in paddy fields to maintain saturation during PI to grain filling stage. Carry out spraying of insecticides or pesticides in crops during morning and afternoon hours. Keep watch on the infestation of gundhi bug and BPH in paddy. Complete nursery sowing for winter vegetable crops. Irrigate the vegetable nursery during morning and evening time. Go for sowing black gram and green gram in upland rice fallow condition.

Kharif Rice:

Paddy is at PI to flowering stage. As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop. Current weather favors the infestation of Leaf Folder. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days. If infestation of BPH & WBPH occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 40g Thiamethoxam or 60g Flonicamid per acre in alternate basis. To prevent crop from Gundhi bug apply 5 ml neem based insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. For prevention of false smut spray the field before panicle initiation or flowering with propiconazole 25%EC @ 2ml/liter of water. In the event of bacterial leaf streak i.e. when the leaves are brown and dry in paddy crop, use plantomycin 1 g + copper oxychloride 1g per liter water only in the clear weather condition.

PULSES:

This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

GROUNDNUT:

Harvesting of Groundnut should be done if 75-80 % of pods are fully matured. The important indication of maturity is yellowing of foliage and necrotic spotting of leaves and dropping of older leaves.

Mustard:

Taking the advantage of available soil moisture due to recent rain, land preparation and sowing of the crop should be started to get optimum yield. Suitable varieties: Toria- Anuradha, Parbati, Sushree, Indian mustard- Mahyco bold, JD 6. Seed rate- 8-10 Kg/ha Spacing- 30 cm X 10 cm, Fertilizer dose: Toria @60:30:30 N-P2O5-K2O Kg/ha Mustard @ 80:40:40 N-P2O5-K2O Kg/ha. Apply 50%N and all P2O5 and K2O fertilizers should be applied as basal.

VEGETABLES:

Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. In case of Cauliflower in this dry days before transplanting during final land preparation apply 8-10 ton FYM per acre and fertilizer like N:P:K each 25-30 Kg/acre should be applied. Short duration Cauliflower varieties (5-6weeks) should be transplanted at distance 60cm×30cm and medium and late duration varieties (3-4 week seedlings) should be transplanted at 60-75cm×40-45cm

- To control Fruit and shoot borer in Brinjal and Okra, Spray Neem based pesticide 1500ppm @600ml/acre at initial stages. Later ongo for alternate spray of Spinosad 45% SC @70ml/acre and Thiodicarb@300g/acre at 15 days interval in 200 litre of water. To control YMV in Okra, thrips in Chilli and aphids in Cowpea crop, spray Imidacloprid @ 5 ml in 10 litres of water.

Animal husbandry:

Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination.

Dr Sujata Sethy
 (Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena
 SMS (Agrometeorology)
 Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-243
Date- 05/11/2021

AGROMET ADVISORY BULLETIN FOR BANKI-DAMPARA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31°C and 21°C, respectively. Mean relative humidity of the district was around 81-71%.

Forecasted period: Today(8.30IST) to 10th November (8.30IST)2021

Weather Advisory: Dry weather is forecasted for today to Tuesday in the block. The day temperature will be 30-31°C and night temperature will be 18-20°C for upcoming 5 days. Maximum and minimum relative humidity will be 68-88% and 34-44% respectively. **There will be clear to partly cloudy sky for next 5 days.**

Block	Banki-Dampara (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	06/11/2021	07/11/2021	08/11/2021	09/11/2021	10/11/2021
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	31	30	31	31	31
T-Min(°C)	20	19	19	18	19
Cloud Cover(octa)	2	3	2	0	0
RH Max (%)	88	74	72	68	79
RH Min (%)	44	37	36	34	34
Wind Speed (Kmph)	6	6	6	6	6
Wind Direction (deg)	360	11	360	345	360

Agromet Advisory

General Advisory: Farmers should go for harvesting of early matured paddy at 85% maturity of the panicles in sunny days. They should keep 5 cm water in paddy fields to maintain saturation during PI to grain filling stage. Carry out spraying of insecticides or pesticides in crops during morning and afternoon hours. Keep watch on the infestation of gundhi bug and BPH in paddy. Complete nursery sowing for winter vegetable crops. Irrigate the vegetable nursery during morning and evening time. Go for sowing black gram and green gram in upland rice fallow condition.

Kharif Rice:

Paddy is at PI to flowering stage. **As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop.** Current weather favors the infestation of **Leaf Folder**. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days. If infestation of **BPH & WBPH** occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 40g Thiamethoxam or 60g Flonicamid per acre in alternate basis. To prevent crop from **Gundhi bug** apply 5 ml neem based insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. For prevention of **false smut** spray the field before panicle initiation or flowering with propiconazole 25%EC @ 2ml/liter of water. In the event of **bacterial leaf streak** i.e. when the leaves are brown and dry in paddy crop, use plantomycin 1 g + copper oxychloride 1g per liter water only in the clear weather condition.

PULSES:

This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

GROUNDNUT:

Harvesting of Groundnut should be done if 75-80 % of pods are fully matured. The important indication of maturity is yellowing of foliage and necrotic spotting of leaves and dropping of older leaves.

Mustard:

Taking the advantage of available soil moisture due to recent rain, land preparation and sowing of the crop should be started to get optimum yield. Suitable varieties: Toria- Anuradha, Parbati, Sushree, Indian mustard- Mahyco bold, JD 6. Seed rate- 8-10 Kg/ha Spacing- 30 cm X 10 cm, Fertilizer dose: Toria @60:30:30 N-P2O5-K2O Kg/ha Mustard @ 80:40:40 N-P2O5-K2O Kg/ha. Apply 50%N and all P2O5 and K2O fertilizers should be applied as basal.

VEGETABLES:

Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. **In case of Cauliflower in this dry days before transplanting during final land preparation apply 8-10 ton FYM per acre and fertilizer like N:P:K each 25-30 Kg/acre should be applied. Short duration Cauliflower varieties (5-6 weeks) should be transplanted at distance 60cm×30cm and medium and late duration varieties (3-4 week seedlings) should be transplanted at 60-75cm×40-45cm**

- To control Fruit and shoot borer in Brinjal and Okra, Spray Neem based pesticide 1500ppm @600ml/acre at initial stages. Later ongo for alternate spray of Spinosad 45% SC @70ml/acre and Thiodicarb @300g/acre at 15 days interval in 200 litre of water. To control YMV in Okra, thrips in Chilli and aphids in Cowpea crop, spray Imidacloprid @ 5 ml in 10 litres of water.

Animal husbandry:

Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-243
Date- 05/11/2021

AGROMET ADVISORY BULLETIN FOR BARANGA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31°C and 21°C, respectively. Mean relative humidity of the district was around 81-71%.

Forecasted period: Today (8.30IST) to 10th November (8.30IST)2021

Weather advisory: Dry weather is forecasted for today to Tuesday in the block. The day temperature will be 30-31°C and night temperature will be 19-21°C for upcoming 5 days. Maximum and minimum relative humidity will be 71-80% and 34-43% respectively. **There will be clear to partly cloudy sky for next 5 days.**

Block	Baranga (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	06/11/2021	07/11/2021	08/11/2021	09/11/2021	10/11/2021
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	31	30	31	31	31
T-Min(°C)	21	19	20	19	20
Cloud Cover	2	3	2	0	0
RH Max (%)	80	71	72	72	79
RH Min (%)	43	36	37	34	35
Wind Speed (Kmph)	7	6	6	6	7
Wind Direction (deg)	18	360	15	345	18

Agromet Advisory

General Advisory: Farmers should go for harvesting of early matured paddy at 85% maturity of the panicles in sunny days. They should keep 5 cm water in paddy fields to maintain saturation during PI to grain filling stage. Carry out spraying of insecticides or pesticides in crops during morning and afternoon hours. Keep watch on the infestation of gundhi bug and BPH in paddy. Complete nursery sowing for winter vegetable crops. Irrigate the vegetable nursery during morning and evening time. Go for sowing black gram and green gram in upland rice fallow condition.

Kharif Rice:

Paddy is at PI to flowering stage. **As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop.** Current weather favors the infestation of **Leaf Folder**. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days. If infestation of **BPH & WBPH** occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 40g Thiamethoxam or 60g Flonicamid per acre in alternate basis. To prevent crop from **Gundhi bug** apply 5 ml neem based insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. For prevention of **false smut** spray the field before panicle initiation or flowering with propiconazole 25%EC @ 2ml/liter of water. In the event of **bacterial leaf streak** i.e. when the leaves are brown and dry in paddy crop, use plantomycin 1 g + copper oxychloride 1g per liter water only in the clear weather condition.

PULSES:

This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

GROUNDNUT:

Harvesting of Groundnut should be done if 75-80 % of pods are fully matured. The important indication of maturity is yellowing of foliage and necrotic spotting of leaves and dropping of older leaves.

Mustard:

Taking the advantage of available soil moisture due to recent rain, land preparation and sowing of the crop should be started to get optimum yield. Suitable varieties: Toria- Anuradha, Parbati, Sushree, Indian mustard- Mahyco bold, JD 6. Seed rate- 8-10 Kg/ha Spacing- 30 cm X 10 cm, Fertilizer dose: Toria @60:30:30 N-P2O5-K2O Kg/ha Mustard @ 80:40:40 N-P2O5-K2O Kg/ha. Apply 50%N and all P2O5 and K2O fertilizers should be applied as basal.

VEGETABLES:

Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. **In case of Cauliflower in this dry days before transplanting during final land preparation apply 8-10 ton FYM per acre and fertilizer like N:P:K each 25-30 Kg/acre should be applied. Short duration Cauliflower varieties (5-6weeks) should be transplanted at distance 60cm×30cm and medium and late duration varieties (3-4 week seedlings) should be transplanted at 60-75cm×40-45cm.**

- To control Fruit and shoot borer in Brinjal and Okra, Spray Neem based pesticide 1500ppm @600ml/acre at initial stages. Later ongo for alternate spray of Spinosad 45% SC @70ml/acre and Thiodicarb @300g/acre at 15 days interval in 200 litre of water. To control YMV in Okra, thrips in Chilli and aphids in Cowpea crop, spray Imidacloprid @ 5 ml in 10 litres of water.

Animal husbandry:

Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-243
Date- 05/11/2021

AGROMET ADVISORY BULLETIN FOR CUTTACK SADAR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31°C and 21°C, respectively. Mean relative humidity of the district was around 81-71%.

Forecasted period: Today (8.30IST) to 10th November (8.30IST)2021

Weather advisory: Dry weather is forecasted for today to Tuesday in the block. The day temperature will be 30-31°C and night temperature will be 19-21°C for upcoming 5 days. Maximum and minimum relative humidity will be 71-81% and 34-42% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Cuttack sadar (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	06/11/2021	07/11/2021	08/11/2021	09/11/2021	10/11/2021
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	31	30	30	31	31
T-Min(°C)	21	19	20	20	20
Cloud Cover(octa)	2	3	2	0	0
RH Max (%)	81	71	72	73	77
RH Min (%)	42	35	36	34	35
Wind Speed (Kmph)	6	6	6	5	6
Wind Direction (deg)	30	360	15	349	18

Agromet Advisory

General Advisory: Farmers should go for harvesting of early matured paddy at 85% maturity of the panicles in sunny days. They should keep 5 cm water in paddy fields to maintain saturation during PI to grain filling stage. Carry out spraying of insecticides or pesticides in crops during morning and afternoon hours. Complete nursery sowing for winter vegetable crops. They should cover the nursery beds with polythene in night to protect it from cold temperature. Irrigate the vegetable nursery during morning and evening time. Go for sowing black gram and green gram in upland rice fallow condition.

Kharif Rice:

Paddy is at PI to flowering stage. As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop. Current weather favors the infestation of **Leaf Folder**. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days. If infestation of **BPH & WBPH** occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 40g Thiamethoxam or 60g Flonicamid per acre in alternate basis. To prevent crop from **Gundhi bug** apply 5 ml neem based insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. For prevention of **false smut** spray the field before panicle initiation or flowering with propiconazole 25%EC @ 2ml/liter of water. In the event of **bacterial leaf streak** i.e. when the leaves are brown and dry in paddy crop, use plantomycin 1 g + copper oxychloride 1g per liter water only in the clear weather condition.

Sugarcane:

Remove the borer affected tillers and late formed tillers, tie the cane shoot with two or three together with partially dried lower leaves. Present weather conditions are favourable for the incidence of **ring spot disease** in Sugarcane. To control go for spraying of copperoxychloride @ 3g/l (or) Carbendazim @ 1g/l (or) Mancozeb @ 3g/l for three times once in three weeks. To control **top shoot borers** spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water. For control of Sucking pest in sugarcane like **Woolly aphid, Whitefly, Scale insects, mealybug** in present weather conditions go for alternate spraying of thiomethoxam 25WG @ 0.25 gm/lit or imidacloprid 0.45 ml/lit of water or Dinotefuron 20% SG 3gm/lit of Water.

PULSES:

This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

VEGETABLES:

Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. In case of **Cauliflower** in this dry days before transplanting during final land preparation apply 8-10 ton FYM per acre and fertilizer like N:P:K each 25-30 Kg/acre should be applied. Short duration Cauliflower varieties (5-6 weeks) should be transplanted at distance 60cm×30cm and medium and late duration varieties (3-4 week seedlings) should be transplanted at 60-75cm×40-45cm.

To control Fruit and shoot borer in Brinjal and Okra, Spray Neem based pesticide 1500ppm @ 600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45% SC @ 70ml/acre and Thiodicarb @ 300g/acre at 15 days interval in 200 litre of water. To control YMV in Okra, thrips in Chilli and aphids in Cowpea crop, spray Imidacloprid @ 5 ml in 10 litres of water.

Animal husbandry:

Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-243
Date- 05/11/2021

AGROMET ADVISORY BULLETIN FOR KANTAPADA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31°C and 21°C, respectively. Mean relative humidity of the district was around 81-71%.

Forecasted period: Today(8.30IST) to 10th November (8.30IST)2021

Weather advisory: Dry weather is forecasted for today to Tuesday in the block. The day temperature will be 30-31°C and night temperature will be 20-21°C for upcoming 5 days. Maximum and minimum relative humidity will be 71-78% and 35-45% respectively. **There will be clear to partly cloudy sky for next 5days.**

Block	Kantapada (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	06/11/2021	07/11/2021	08/11/2021	09/11/2021	10/11/2021
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	31	30	31	31	31
T-Min(°C)	21	20	20	20	20
Cloud Cover(octa)	2	3	2	0	0
RH Max (%)	77	71	71	74	78
RH Min (%)	45	37	37	35	37
Wind Speed (Kmph)	8	8	7	7	8
Wind Direction (deg)	15	11	360	345	15

Agromet Advisory

General Adviosry: Farmers should go for harvesting of early matured paddy at 85% maturity of the panicles in sunny days. They should keep 5 cm water in paddy fields to maintain saturation during PI to grain filling stage. Carry out spraying of insecticides or pesticides in crops during morning and afternoon hours. Complete nursery sowing for winter vegetable crops. They should cover the nursery beds with polythine in night to protect it from cold temperature. Irrigate the vegetable nursery during morning and evening time. Go for sowing black gram and green gram in upland rice fallow condition.

Kharif Rice:

Paddy is at PI to flowering stage. **As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop.** Current weather favors the infestation of **Leaf Folder**. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days. If infestation of **BPH & WBPH** occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 40g Thiamethoxam or 60g Flonicamid per acre in alternate basis. To prevent crop from **Gundhi bug** apply 5 ml neem baesd insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. For prevention of **false smut** spray the field before panicle initiation or flowering with propiconazole 25%EC@ 2ml/liter of water. In the event of **bacterial leaf streak** i.e. when the leaves are brown and dry in paddy crop, use plantomycin 1 g + cupper oxychloride 1g per liter water only in the clear weather condition.

Sugarcane:

Remove the borer affected tillers and late formed tillers, tie the cane shoot with two or three together with partially dried lower leaves. Present weather conditions are favourable for the incidence of **ring spot disease** in Sugarcane. To control go for spraying of copperoxychloride @ 3g/l (or) Carbendazim @ 1g/l (or) Mancozeb @3g/l for three times once in three weeks. To control **top shoot borers** spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water For control of Sucking pest in sugarcane like **Wooly aphid, Whitefly, Scale insects, mealybug** in present weather conditions go for alternate spraying of thiomethoxam 25WG @ 0.25 gm/lit or imidacloprid 0.45 ml/lit of water or Dinotefuron 20% SG 3gm/lit of Water.

PULSES:

This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

VEGETABLES:

Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. **In case of Cauliflower in this dry days before transplanting during final land preparation apply 8-10 ton FYM per acre and fertilizer like N:P:K each 25-30 Kg/acre should be applied. Short duration Cauliflower varieties (5-6weeks) should be transplanted at distance 60cm×30cm and medium and late duration varieties (3-4 week seedlings) should be transplanted at 60-75cm×40-45cm.**

To control Fruit and shoot borer in Brinjal and Okra, Spray Neem based pesticide 1500ppm @600ml/acre at initial stages. Later ongo for alternate spray of Spinosad 45% SC @70ml/acre and Thiodicarb@300g/acre at 15 days interval in 200 litre of water. To control YMV in Okra, thrips in Chilli and aphids in Cowpea crop, spray Imidacloprid @ 5 ml in 10 litres of water.

Animal husbandry:

Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-243
Date- 05/11/2021

AGROMET ADVISORY BULLETIN FOR MAHANGA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31°C and 21°C, respectively. Mean relative humidity of the district was around 81-71%.

Forecasted period: Today(8.30IST) to 10th November (8.30IST)2021

Weather advisory: Dry weather is forecasted for today to Tuesday in the block. The day temperature will be and it will be 30°C and night temperature will be 19-20°C for upcoming 5 days. Maximum and minimum relative humidity will be 68-75% and 35-44% respectively. There will be clear to partly cloudy sky for next 5days.

Block	Mahanga (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	06/11/2021	07/11/2021	08/11/2021	09/11/2021	10/11/2021
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	30	30	30	30	30
T-Min(°C)	20	19	19	19	20
Cloud Cover(octal)	2	3	2	0	0
RH Max (%)	75	68	69	71	74
RH Min (%)	44	36	37	35	37
Wind Speed (Kmph)	7	7	7	7	7
Wind Direction (deg)	11	18	345	18	349

Agromet Advisory

General Adviosry: Farmers should go for harvesting of early matured paddy at 85% maturity of the panicles in sunny days. They should keep 5 cm water in paddy fields to maintain saturation during PI to grain filling stage. Carry out spraying of insecticides or pesticides in crops during morning and afternoon hours. Complete nursery sowing for winter vegetable crops. They should cover the nursery beds with polythine in night to protect it from cold temperature. Irrigate the vegetable nursery during morning and evening time. Go for sowing black gram and green gram in upland rice fallow condition.

Kharif Rice:

Paddy is at PI to flowering stage. As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop. Current weather favors the infestation of **Leaf Folder**. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days. If infestation of **BPH & WBPH** occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 40g Thiamethoxam or 60g Flonicamid per acre in alternate basis. To prevent crop from **Gundhi bug** apply 5 ml neem baesd insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. For prevention of **false smut** spray the field before panicle initiation or flowering with propiconazole 25%EC@ 2ml/liter of water. In the event of **bacterial leaf streak** i.e. when the leaves are brown and dry in paddy crop, use plantomycin 1 g + cupper oxychloride 1g per liter water only in the clear weather condition.

Sugarcane:

Remove the borer affected tillers and late formed tillers, tie the cane shoot with two or three together with partially dried lower leaves. Present weather conditions are favourable for the incidence of **ring spot disease** in Sugarcane. To control go for spraying of copperoxychloride @ 3g/l (or) Carbendazim @ 1g/l (or) Mancozeb @3g/l for three times once in three weeks. To control **top shoot borers** spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water For control of Sucking pest in sugarcane like **Wooly aphid, Whitefly, Scale insects, mealybug** in present weather conditions go for alternate spraying of thiomethoxam 25WG @ 0.25 gm/lit or imidacloprid 0.45 ml/lit of water or Dinotefuron 20% SG 3gm/lit of Water.

PULSES:

This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

VEGETABLES:

Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. In case of **Cauliflower** in this dry days before transplanting during final land preparation apply 8-10 ton FYM per acre and fertilizer like N:P:K each 25-30 Kg/acre should be applied. Short duration Cauliflower varieties (5-6weeks) should be transplanted at distance 60cm×30cm and medium and late duration varieties (3-4 week seedlings) should be transplanted at 60-75cm×40-45cm.

To control Fruit and shoot borer in Brinjal and Okra, Spray Neem based pesticide 1500ppm @600ml/acre at initial stages. Later ongo for alternate spray of Spinosad 45% SC @70ml/acre and Thiodicarb@300g/acre at 15 days interval in 200 litre of water. To control YMV in Okra, thrips in Chilli and aphids in Cowpea crop, spray Imidacloprid @ 5 ml in 10 litres of water.

Animal husbandry:

Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-243
Date- 05/11/2021

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31°C and 21°C, respectively. Mean relative humidity of the district was around 81-71%.

Forecasted period: Today(8.30IST) to 10th November (8.30IST)2021

Weather advisory: Dry weather is forecasted for today to Tuesday in the block. The day temperature will be 30-31°C and night temperature will be 17-19°C for upcoming 5 days. Maximum and minimum relative humidity will be 64-90% and 31-48% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Narasinghpur (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	06/11/2021	07/11/2021	08/11/2021	09/11/2021	10/11/2021
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	30	30	30	30	31
T-Min(°C)	19	18	18	17	18
Cloud Cover(octa)	2	3	2	0	0
RH Max (%)	90	76	70	64	73
RH Min (%)	48	38	35	31	32
Wind Speed (Kmph)	3	4	5	6	4
Wind Direction (deg)	360	15	349	349	15

Agromet Advisory

General Advisory: Farmers should go for harvesting of early matured paddy at 85% maturity of the panicles in sunny days. They should keep 5 cm water in paddy fields to maintain saturation during PI to grain filling stage. Carry out spraying of insecticides or pesticides in crops during morning and afternoon hours. Keep watch on the infestation of gundhi bug and BPH in paddy. Complete nursery sowing for winter vegetable crops. Irrigate the vegetable nursery during morning and evening time. Go for sowing black gram and green gram in upland rice fallow condition.

Kharif Rice:

Paddy is at PI to flowering stage. As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop. Current weather favors the infestation of **Leaf Folder**. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days. If infestation of **BPH & WBPH** occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 40g Thiamethoxam or 60g Flonicamid per acre in alternate basis. To prevent crop from **Gundhi bug** apply 5 ml neem based insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. For prevention of **false smut** spray the field before panicle initiation or flowering with propiconazole 25%EC @ 2ml/liter of water. In the event of **bacterial leaf streak** i.e. when the leaves are brown and dry in paddy crop, use plantomycin 1 g + cupper oxychloride 1g per liter water only in the clear weather condition.

PULSES:

This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

GROUNDNUT:

Harvesting of Groundnut should be done if 75-80 % of pods are fully matured. The important indication of maturity is yellowing of foliage and necrotic spotting of leaves and dropping of older leaves.

Mustard:

Taking the advantage of available soil moisture due to recent rain, land preparation and sowing of the crop should be started to get optimum yield. Suitable varieties: Toria- Anuradha, Parbati, Sushree, Indian mustard- Mahyco bold, JD 6. Seed rate- 8-10 Kg/ha Spacing- 30 cm X 10 cm, Fertilizer dose: Toria @60:30:30 N-P2O5-K2O Kg/ha Mustard @ 80:40:40 N-P2O5-K2O Kg/ha. Apply 50%N and all P2O5 and K2O fertilizers should be applied as basal.

VEGETABLES:

Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. In case of Cauliflower in this dry days before transplanting during final land preparation apply 8-10 ton FYM per acre and fertilizer like N:P:K each 25-30 Kg/acre should be applied. Short duration Cauliflower varieties (5-6 weeks) should be transplanted at distance 60cm×30cm and medium and late duration varieties (3-4 week seedlings) should be transplanted at 60-75cm×40-45cm

- To control Fruit and shoot borer in Brinjal and Okra, Spray Neem based pesticide 1500ppm @600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45% SC @70ml/acre and Thiodicarb @300g/acre at 15 days interval in 200 litre of water. To control YMV in Okra, thrips in Chilli and aphids in Cowpea crop, spray Imidacloprid @ 5 ml in 10 litres of water.

Animal husbandry:

Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-243
Date- 05/11/2021

AGROMET ADVISORY BULLETIN FOR NIALI

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31°C and 21°C, respectively. Mean relative humidity of the district was around 81-71%.

Forecasted period: Today(8.30IST) to 10th November (8.30IST)2021

Weather advisory: Dry weather is forecasted for today to Tuesday in the block. The day temperature will be 30-31°C and night temperature will be 19-21°C for upcoming 5 days. Maximum and minimum relative humidity will be 71-78% and 36-46% respectively. **There will be clear to partly cloudy sky for next 5 days.**

Block	Niali (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	06/11/2021	07/11/2021	08/11/2021	09/11/2021	10/11/2021
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	31	30	31	31	30
T-Min(°C)	21	19	20	20	20
Cloud Cover(octa)	2	3	2	0	0
RH Max (%)	75	72	71	74	78
RH Min (%)	46	38	38	36	38
Wind Speed(Kmph)	9	9	8	8	8
Wind Direction(deg)	15	15	360	360	15

Agromet Advisory

General Advisory: Farmers should keep the harvested rice in safer dry places and should go for harvesting of early matured paddy at 85% maturity of the panicles in sunny days. They should keep 5 cm water in paddy fields to maintain saturation during PI to grain filling stage. Carry out spraying of insecticides or pesticides in crops during morning and afternoon hours. Complete nursery sowing for winter vegetable crops. They should cover the nursery beds with polythene in night to protect it from cold temperature. Irrigate the vegetable nursery during morning and evening time. Go for sowing black gram and green gram in upland rice fallow condition.

Kharif Rice:

Paddy is at PI to flowering stage. **As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop.** Current weather favors the infestation of **Leaf Folder**. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days. If infestation of **BPH & WBPH** occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 40g Thiamethoxam or 60g Flonicamid per acre in alternate basis. To prevent crop from **Gundhi bug** apply 5 ml neem based insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. For prevention of **false smut** spray the field before panicle initiation or flowering with propiconazole 25%EC @ 2ml/liter of water. In the event of **bacterial leaf streak** i.e. when the leaves are brown and dry in paddy crop, use plantomycin 1 g + copper oxychloride 1g per liter water only in the clear weather condition.

Sugarcane:

Remove the borer affected tillers and late formed tillers, tie the cane shoot with two or three together with partially dried lower leaves. Present weather conditions are favourable for the incidence of **ring spot disease** in Sugarcane. To control go for spraying of copperoxychloride @ 3g/l (or) Carbendazim @ 1g/l (or) Mancozeb @ 3g/l for three times once in three weeks. To control **top shoot borers** spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water. For control of Sucking pest in sugarcane like **Woolly aphid, Whitefly, Scale insects, mealybug** in present weather conditions go for alternate spraying of thiomethoxam 25WG @ 0.25 gm/lit or imidacloprid 0.45 ml/lit of water or Dinotefuron 20% SG 3gm/lit of Water.

PULSES:

This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

VEGETABLES:

Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. **In case of Cauliflower in this dry days before transplanting during final land preparation apply 8-10 ton FYM per acre and fertilizer like N:P:K each 25-30 Kg/acre should be applied. Short duration Cauliflower varieties (5-6weeks) should be transplanted at distance 60cm×30cm and medium and late duration varieties (3-4 week seedlings) should be transplanted at 60-75cm×40-45cm.**

To control Fruit and shoot borer in Brinjal and Okra, Spray Neem based pesticide 1500ppm @ 600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45% SC @ 70ml/acre and Thiodicarb @ 300g/acre at 15 days interval in 200 litre of water. To control YMV in Okra, thrips in Chilli and aphids in Cowpea crop, spray Imidacloprid @ 5 ml in 10 litres of water.

Animal husbandry: Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-243
Date- 05/11/2021

AGROMET ADVISORY BULLETIN FOR NISCHINTAKOILI

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31°C and 21°C, respectively. Mean relative humidity of the district was around 81-71%.

Forecasted period: Today(8.30IST) to 10th November (8.30IST)2021

Weather advisory: Dry weather is forecasted for today to Tuesday in the block. The day temperature will be 30-31°C and night temperature will be 19-20°C for upcoming 5 days. Maximum and minimum relative humidity will be 68-76% and 35-44% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Nischintakoili (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	06/11/2021	07/11/2021	08/11/2021	09/11/2021	10/11/2021
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	30	30	30	31	30
T-Min(°C)	20	19	19	20	20
Cloud Cover(octa)	2	3	2	0	0
RH Max (%)	76	68	69	71	74
RH Min (%)	44	37	37	35	37
Wind Speed (Kmph)	8	7	7	7	7
Wind Direction (deg)	360	18	345	11	360

Agromet Advisory

General Advisory: Farmers should go for harvesting of early matured paddy at 85% maturity of the panicles in sunny days. They should keep 5 cm water in paddy fields to maintain saturation during PI to grain filling stage. Carry out spraying of insecticides or pesticides in crops during morning and afternoon hours. Complete nursery sowing for winter vegetable crops. They should cover the nursery beds with polythene in night to protect it from cold temperature. Irrigate the vegetable nursery during morning and evening time. Go for sowing black gram and green gram in upland rice fallow condition.

Kharif Rice:

Paddy is at PI to flowering stage. As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop. Current weather favors the infestation of **Leaf Folder**. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days. If infestation of **BPH & WBPH** occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 40g Thiamethoxam or 60g Flonicamid per acre in alternate basis. To prevent crop from **Gundhi bug** apply 5 ml neem based insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. For prevention of **false smut** spray the field before panicle initiation or flowering with propiconazole 25%EC @ 2ml/liter of water. In the event of **bacterial leaf streak** i.e. when the leaves are brown and dry in paddy crop, use plantomycin 1 g + copper oxychloride 1g per liter water only in the clear weather condition.

Sugarcane:

Remove the borer affected tillers and late formed tillers, tie the cane shoot with two or three together with partially dried lower leaves. Present weather conditions are favourable for the incidence of **ring spot disease** in Sugarcane. To control go for spraying of copperoxychloride @ 3g/l (or) Carbendazim @ 1g/l (or) Mancozeb @ 3g/l for three times once in three weeks. To control **top shoot borers** spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water. For control of Sucking pest in sugarcane like **Woolly aphid, Whitefly, Scale insects, mealybug** in present weather conditions go for alternate spraying of thiomethoxam 25WG @ 0.25 gm/lit or imidacloprid 0.45 ml/lit of water or Dinotefuron 20% SG 3gm/lit of Water.

PULSES:

This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

VEGETABLES:

Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. In case of **Cauliflower** in this dry days before transplanting during final land preparation apply 8-10 ton FYM per acre and fertilizer like N:P:K each 25-30 Kg/acre should be applied. Short duration Cauliflower varieties (5-6 weeks) should be transplanted at distance 60cm×30cm and medium and late duration varieties (3-4 week seedlings) should be transplanted at 60-75cm×40-45cm.

To control Fruit and shoot borer in Brinjal and Okra, Spray Neem based pesticide 1500ppm @ 600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45% SC @ 70ml/acre and Thiodicarb @ 300g/acre at 15 days interval in 200 litre of water. To control YMV in Okra, thrips in Chilli and aphids in Cowpea crop, spray Imidacloprid @ 5 ml in 10 litres of water.

Animal husbandry:

Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-243
Date- 05/11/2021

AGROMET ADVISORY BULLETIN FOR SALEPUR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31°C and 21°C, respectively. Mean relative humidity of the district was around 81-71%.

Forecasted period: Today(8.30IST) to 10th November (8.30IST)2021

Weather advisory: Dry weather is forecasted for today to Tuesday in the block. The day temperature will be 30-31°C and night temperature will be 19-20°C for upcoming 5 days. Maximum and minimum relative humidity will be 69-76% and 35-44% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Salepur (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	06/11/2021	07/11/2021	08/11/2021	09/11/2021	10/11/2021
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	30	30	30	31	30
T-Min(°C)	20	19	20	20	20
Cloud Cover(octa)	2	3	2	0	0
RH Max (%)	76	69	69	72	74
RH Min (%)	44	36	37	35	37
Wind Speed (Kmph)	7	7	7	7	7
Wind Direction (deg)	360	18	345	11	360

Agromet Advisory

General Advisory: Farmers should go for harvesting of early matured paddy at 85% maturity of the panicles in sunny days. They should keep 5 cm water in paddy fields to maintain saturation during PI to grain filling stage. Carry out spraying of insecticides or pesticides in crops during morning and afternoon hours. Keep watch on the infestation of gundhi bug and BPH in paddy. Complete nursery sowing for winter vegetable crops. Irrigate the vegetable nursery during morning and evening time. Go for sowing black gram and green gram in upland rice fallow condition.

Kharif Rice:

Paddy is at PI to flowering stage. As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop. Current weather favors the infestation of Leaf Folder. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days. If infestation of BPH & WBPH occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 40g Thiamethoxam or 60g Flonicamid per acre in alternate basis. To prevent crop from Gundhi bug apply 5 ml neem based insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. For prevention of false smut spray the field before panicle initiation or flowering with propiconazole 25%EC @ 2ml/liter of water. In the event of bacterial leaf streak i.e. when the leaves are brown and dry in paddy crop, use plantomycin 1 g + copper oxychloride 1g per liter water only in the clear weather condition.

PULSES:

This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

Mustard:

Taking the advantage of available soil moisture due to recent rain, land preparation and sowing of the crop should be started to get optimum yield. Suitable varieties: Toria- Anuradha, Parbati, Sushree, Indian mustard- Mahyco bold, JD 6. Seed rate- 8-10 Kg/ha Spacing- 30 cm X 10 cm, Fertilizer dose: Toria @60:30:30 N-P2O5-K2O Kg/ha Mustard @ 80:40:40 N-P2O5-K2O Kg/ha. Apply 50%N and all P2O5 and K2O fertilizers should be applied as basal.

VEGETABLES:

Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato.

- In case of Cauliflower in this dry days before transplanting during final land preparation apply 8-10 ton FYM per acre and fertilizer like N:P:K each 25-30 Kg/acre should be applied. Short duration Cauliflower varieties (5-6weeks) should be transplanted at distance 60cm×30cm and medium and late duration varieties (3-4 week seedlings) should be transplanted at 60-75cm×40-45cm
- To control Fruit and shoot borer in Brinjal and Okra, Spray Neem based pesticide 1500ppm @600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45% SC @70ml/acre and Thiodicarb@300g/acre at 15 days interval in 200 litre of water. To control YMV in Okra, thrips in Chilli and aphids in Cowpea crop, spray Imidacloprid @ 5 ml in 10 litres of water.

Animal husbandry:

Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-243
Date- 05/11/2021

AGROMET ADVISORY BULLETIN FOR TANGI-CHOUDWAR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31°C and 21°C, respectively. Mean relative humidity of the district was around 81-71%.

Forecasted period: Today(8.30IST) to 10th November (8.30IST)2021

Weather advisory: Dry weather is forecasted for today to Tuesday in the block. The day temperature will be 30°C and night temperature will be 20-21°C for upcoming 5 days. Maximum and minimum relative humidity will be 68-75% and 41-52% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Tangi-Choudwar (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	06/11/2021	07/11/2021	08/11/2021	09/11/2021	10/11/2021
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	30	30	30	30	30
T-Min(°C)	21	20	20	20	20
Cloud Cover(octa)	2	3	2	0	0
RH Max (%)	75	68	72	70	74
RH Min (%)	52	41	43	41	46
Wind Speed (Kmph)	11	11	10	10	10
Wind Direction (deg)	60	60	45	45	60

Agromet Advisory

General Advisory: Farmers should go for harvesting of early matured paddy at 85% maturity of the panicles in sunny days. They should keep 5 cm water in paddy fields to maintain saturation during PI to grain filling stage. Carry out spraying of insecticides or pesticides in crops during morning and afternoon hours. Keep watch on the infestation of gundhi bug and BPH in paddy. Complete nursery sowing for winter vegetable crops. Irrigate the vegetable nursery during morning and evening time. Go for sowing black gram and green gram in upland rice fallow condition.

Kharif Rice:

Paddy is at PI to flowering stage. As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop. Current weather favors the infestation of **Leaf Folder**. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days. If infestation of **BPH & WBPH** occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 40g Thiamethoxam or 60g Flonicamid per acre in alternate basis. To prevent crop from **Gundhi bug** apply 5 ml neem based insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. For prevention of **false smut** spray the field before panicle initiation or flowering with propiconazole 25%EC @ 2ml/liter of water. In the event of **bacterial leaf streak** i.e. when the leaves are brown and dry in paddy crop, use plantomycin 1 g + copper oxychloride 1g per liter water only in the clear weather condition.

PULSES:

This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

Mustard:

Taking the advantage of available soil moisture due to recent rain, land preparation and sowing of the crop should be started to get optimum yield. Suitable varieties: Toria- Anuradha, Parbati, Sushree, Indian mustard- Mahyco bold, JD 6. Seed rate- 8-10 Kg/ha Spacing- 30 cm X 10 cm, Fertilizer dose: Toria @60:30:30 N-P2O5-K2O Kg/ha Mustard @ 80:40:40 N-P2O5-K2O Kg/ha. Apply 50%N and all P2O5 and K2O fertilizers should be applied as basal.

VEGETABLES:

Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. In case of Cauliflower in this dry days before transplanting during final land preparation apply 8-10 ton FYM per acre and fertilizer like N:P:K each 25-30 Kg/acre should be applied. Short duration Cauliflower varieties (5-6weeks) should be transplanted at distance 60cm×30cm and medium and late duration varieties (3-4 week seedlings) should be transplanted at 60-75cm×40-45cm

- To control Fruit and shoot borer in Brinjal and Okra, Spray Neem based pesticide 1500ppm @600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45% SC @70ml/acre and Thiodicarb@300g/acre at 15 days interval in 200 litre of water. To control YMV in Okra, thrips in Chilli and aphids in Cowpea crop, spray Imidacloprid @ 5 ml in 10 litres of water.

Animal husbandry:

Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-243
Date- 05/11/2021

AGROMET ADVISORY BULLETIN FOR TIGIRIA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31°C and 21°C, respectively. Mean relative humidity of the district was around 81-71%.

Forecasted period: Today(8.30IST) to 10th November (8.30IST)2021

Weather advisory: Dry weather is forecasted for today to Tuesday in the block. The day temperature will be 30-31 and night temperature will be 18-20°C for upcoming 5 days. Maximum and minimum relative humidity will be 69-89% and 35-44% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Tigiria (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	06/11/2021	07/11/2021	08/11/2021	09/11/2021	10/11/2021
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	31	30	30	31	31
T-Min(°C)	20	19	19	18	19
Cloud Cover(octa)	2	3	2	0	0
RH Max (%)	89	74	73	69	79
RH Min (%)	44	37	37	35	35
Wind Speed (Kmph)	6	6	5	6	5
Wind Direction (deg)	11	15	345	335	360

Agromet Advisory

General Advisory: Farmers should go for harvesting of early matured paddy at 85% maturity of the panicles in sunny days. They should keep 5 cm water in paddy fields to maintain saturation during PI to grain filling stage. Carry out spraying of insecticides or pesticides in crops during morning and afternoon hours. Keep watch on the infestation of gundhi bug and BPH in paddy. Complete nursery sowing for winter vegetable crops. Irrigate the vegetable nursery during morning and evening time. Go for sowing black gram and green gram in upland rice fallow condition.

Kharif Rice:

Paddy is at PI to flowering stage. As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop. Current weather favors the infestation of Leaf Folder. If observe in field then spray Chlorantraniliprole 18.5% SC @ 3 ml/10 litre of water or Fipronil 5% SC @ 2 ml/litre of water in sunny days. If infestation of BPH & WBPH occurs, it is advised to alter the micro-climate of the rice field by alternate wetting and drying technique. If problem still persists, apply 40g Thiamethoxam or 60g Flonicamid per acre in alternate basis. To prevent crop from Gundhi bug apply 5 ml neem based insecticide (300 ppm) per liter of water or spray 1.5 ml Lambda Cyhalothrin per liter of water. For prevention of false smut spray the field before panicle initiation or flowering with propiconazole 25% EC @ 2ml/liter of water. In the event of bacterial leaf streak i.e. when the leaves are brown and dry in paddy crop, use plantomycin 1 g + copper oxychloride 1g per liter water only in the clear weather condition.

PULSES:

This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture. Apply 4 kg urea, 35 kg DAP and 13 kg MOP per acre during final land preparation. Sow the seeds in line of 30 cm apart with 10-15 cm plant to plant distance.

GROUNDNUT:

Harvesting of Groundnut should be done if 75-80 % of pods are fully matured. The important indication of maturity is yellowing of foliage and necrotic spotting of leaves and dropping of older leaves.

Mustard:

Taking the advantage of available soil moisture due to recent rain, land preparation and sowing of the crop should be started to get optimum yield. Suitable varieties: Toria- Anuradha, Parbati, Sushree, Indian mustard- Mahyco bold, JD 6. Seed rate- 8-10 Kg/ha Spacing- 30 cm X 10 cm, Fertilizer dose: Toria @ 60:30:30 N-P2O5-K2O Kg/ha Mustard @ 80:40:40 N-P2O5-K2O Kg/ha. Apply 50%N and all P2O5 and K2O fertilizers should be applied as basal.

VEGETABLES:

Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. In case of Cauliflower in this dry days before transplanting during final land preparation apply 8-10 ton FYM per acre and fertilizer like N:P:K each 25-30 Kg/acre should be applied. Short duration Cauliflower varieties (5-6 weeks) should be transplanted at distance 60cm×30cm and medium and late duration varieties (3-4 week seedlings) should be transplanted at 60-75cm×40-45cm

- To control Fruit and shoot borer in Brinjal and Okra, Spray Neem based pesticide 1500ppm @ 600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45% SC @ 70ml/acre and Thiodicarb @ 300g/acre at 15 days interval in 200 litre of water. To control YMV in Okra, thrips in Chilli and aphids in Cowpea crop, spray Imidacloprid @ 5 ml in 10 litres of water.

Animal husbandry:

Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-243
 Date- 05/11/2021

				
MEGHDOOT APP	DAMINI APP	FACEBOOK (Damu Cuttack)	BLOCK SPECIFIC WHAT'S APP GROUPS	@DAMUCUTTACK



- Kindly install and use **“Meghdoot mobile App”** for Cuttack district weather forecast, warnings and weather based agromet advisory services.

<https://play.google.com/store/apps/details?id=com.aas.meghdoot>

- Kindly install and use **“Damini mobile App”** for location specific for lightening forecast and warnings.

<https://play.google.com/store/apps/details?id=com.lightening.live.damini>

- Follow the **block specific whats app groups** and join in your block wise groups for getting weather based agromet advisory services twice in a week (Tuesday and Friday) and other weather warning based farm management advisories.

1	ICAR-NRRI, Cuttack, Odisha	https://icar-nrri.in/agromate/	
2	Agrimet division, IMD, Pune	https://imdagrmet.gov.in/imdproject/AGIndex.php	

- Follow the **Official Website links** to get the weather based agromet advisory service bulletins (**English & Odia**) twice in a week (Tuesday and Friday).

Dr Sujata Sethy
 (Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
 SMS (Agrometeorology)
 Contact No-9861445080/7008453427