




GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com




Bulletin No.-272
Date- 18/02/2022

AGROCLIMATIC ZONES OF CUTTACK DISTRICT:




STANDARD PRECIPITATION INDEX AND NDVI MAP



NOAA/VIIRS/BLENDED NDVI Composite ending on Week no 06
 (05.02.2022 to 11.02.2022) over Agricultural region

Odisha



Agriculture vigour is moderate over the state.

As per SPI value (cumulative 4weeks -20 January 2022 – 16 February 2022), mildly wet condition was there in Cuttack district.

According to NDVI map (05.02.2022 to 11.02.2022) the agriculture vigor was light over most eastern blocks and light over some western blocks of Cuttack district.

LEGEND (Medium range block level rainfall forecast)

	No Rainfall	00 mm
	Very light rainfall	01-05 mm
	Light rainfall	06-10 mm
	Light to moderate rainfall	11-30 mm
	Moderate to heavy rainfall	31-65 mm
	Heavy to very heavy rainfall	66-115 mm
	Extremely heavy rainfall	>116 mm

Dr Sujata Sethy
 (Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com





Bulletin No.-272
Date- 18/02/2022

Past week (11-17 Feb, 2022) cumulative rainfall and (12-18 Feb) average minimum temperature and its departure from normal for Cuttack district



Medium range (Today (8.30IST) to Wednesday (8.30IST)) Rainfall (mm) forecast for Cuttack district



Dr Sujata Sethy
 (Nodal officer)



**Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area**



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-272
 Date- 18/02/2022

Met Sub division	Extended range outlook for subsequent 1 week (6 th to 12 th days) (23February-01 March 2022)		
Odisha	<i>Parameter</i>	<i>Departure (%)</i>	<i>Category</i>
	Rainfall	573.84	Above Normal
	<i>Parameter</i>	<i>Anomaly</i>	<i>Category</i>
	Maximum temperature	-1.45	Normal
	Minimum temperature	-1.21	Normal

General Agromet Advisory:-As There will be above normal rainfall is forecasted in subsequent week, after upcoming 5 days, so farmers are advised to complete their pre summer sowing activity in upcoming 5days as soon as possible and should pause irrigation activities in the subsequent week(the subsequent week rainfall activity will be helpful for crops mainly pulses, mustard and groundnut as maximum crops at flowering stage) .

Dr Sujata Sethy
 (Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena
 SMS (Agrometeorology)
 Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)















Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-272

Date- 18/02/2022

Weather (Past and Forecasted) based Disease and pest in Major Crops and vegetables

Sl No	Crop	Stage and condition of the crop	Anticipated Disease and pest	
1	Pulses (Green gram and black gram)	Vegetative stage to flowering		Powdery Mildew
				Flea beetles
				Leaf crinkle virus
2	Mustard	Vegetative stage to flowering		Powdery Mildew
				White Rust
				Painted Bug
				Aphid
3	Rabi Groundnut	Vegetative stage to flowering		Thrips
				Spodoptera/Tobacco caterpillar
4	Vegetables (Tomato, Chilli, Brinjal)	Vegetative to fruiting		Mites
				Chili Aphid
5	Tomato	Flowering to Fruiting		Late blight
				White fly
				Flower drop

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area












Debasish Jena
 SMS (Agrometeorology)
 Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-272
Date- 18/02/2022

6	Potato	Sprouting/Vegetative stage		Potato tuber moth
				Late blight
7	Onion	Flowering		Purple blight
				Stemphylium Blight
7	Cabbage and Cauliflower	Vegetative to Flowering		Diamond back moth
8	Mango	Vegetative to Flowering		Mango hooper
				Mealy Bug
				Powdery Mildew
9	Papaya	Fruting stage		Mealy bug

Dr Sujata Sethy
 (Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA INDIA METEOROLOGICAL DEPARTMENT DISTRICT AGROMET UNIT KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-272

Date- 18/02/2022

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0.1 mm rainfall during the last week. The mean daily maximum and minimum temperature was 29.14°C (-2.66°C from normal) and 15.57°C (-2.3°C from normal), respectively. Mean relative humidity of the district was around 58-79%.

Forecasted period: Today(8.30IST) to 23th February (8.30IST)2022

Weather advisory: Very light rainfall is forecasted for Sunday and Dry weather is forecasted for other 4days in the block. The day temperature will be 31-33°C and night temperature will be 17-19°C for upcoming 5 days. Maximum and minimum relative humidity will be 77-93% and 11-40% respectively. There will be partly cloudy to overcast sky for next 5days.

Block	Athagad (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	18/02/2022	19/02/2022	20/02/2022	21/02/2022	22/02/2022
Rainfall (mm)	0	0	1	0	0
T-Max (°C)	31	31	33	33	31
T-Min (°C)	19	19	19	17	17
Cloud Cover (octa)	3	2	4	5	0
RH Max (%)	78	92	79	77	93
RH Min (%)	40	28	16	12	11
Wind Speed (kmph)	7	8	12	9	7
Wind Direction (deg)	155	169	207	203	23

Agromet Advisory

General advisory: Farmers should undertake field preparation for cultivation of different summer vegetables like sponge gourd, bottle gourd, pumpkin, water melon, musk melon etc. If seedlings are already raised in polythene bags; transplant in main field @ 2-3 seedlings per pit. Irrigation should be paused on Sunday as there will be rainfall is forecasted.

Pulses:

Forecasted weather may aggravate the infestation of **Flea beetle and powdery mildew** in pulses. Spray neem oil 5 ml/L of water to control Flea beetle and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew. There may be chances of **leaf crinkle disease** infestation due to past foggy and cold conditions. To manage the insect vector of the disease like whitefly and aphid spray Thiamethoxam 25% W. G @ 40 gram/acre or Acetamiprid 20% S.P. @ 50 gram/acre or Imidacloprid 17.8% SL @ 60-ml acre by mixing in 200 litre of water in 15 days interval.

Rabi-Groundnut:

Farmers are advised to do intercultivation and apply Gypsum @200 kg/ acre in Groundnut which is at flowering stage. forecasted warm and dry days with low RH favours infestation of **thrips**, to control thrips spray Thiamethoxam 25% W.G @ 40 gram/acre or Imidacloprid 17.8% SL @ 50-ml/acre by mixing it in 200 litre of water. For control of **Tobacco caterpillar and spodotera** for spraying of fipronil 5% SL 400ml (or) indoxacarb 14.5% SC 80gm or neem oil 1000ml @ 200 liter of water per acre.

Mustard: To meet the deficiency of Sulfur applies 200 gram sulfur per acre. **Powdery Mildew:** Spray Sulfex 2 gram/lit. To control **white rust disease** in mustard, spray Copper oxychloride @ 3g per litre of water. **Painted bug and Aphid attack** may take place in mustard crop, apply Thiamethoxam 25% WG @ 3g per 10 litres of water.

Vegetables: **Thrips and mites** are prominent pests in vegetables at present. To control both, spray Bifenthrin @ 1g/litre or Profenophos @ 2ml/litre of water. To control the **sucking pests like aphid & white fly**, spray Imidacloprid @ 5ml/10 lit of water.

Tomato:

Transplanted seedlings are at 6-8 week stage in main field. **Due to water stress in past, deficiency of boron or due to hormonal disbalance flower drop occurs.** Apply timely and required amount of irrigation. Spray Boron 20% 2 gram/lit during early flowering and 15 days later. To manage **late blight** in tomato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days interval. To control **white fly** Spray Imidacloprid (18.7% S.L) 0.8 ml/lit of water or Acetamiprid 1 ml/lit of water or Diafenthiuron 12.5 gram/ 15 lit of water.

Potato: Farmers are advised to start earthing-up in potato if crop has attained a height of 15-22 cm. If so required, the practice may be repeated after 15 days. There are chances of **Potato Tuber Moth** infestation in potato. To manage this pest chemically spray Spinosad 45 % SC @ 75-ml/acre or Emamectin Benzoate 5% SG @ 80-gram/ acre or Chlorantraniliprole 18.5 % SC @ 80- ml/acre. To manage **late blight** in potato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days interval.

Onion: Do inter culture and hand weeding may be carried out at 45-50 DAT and application of 30-kg of urea and 15-kg of MOP fertilizer should be done at this stage. Onion being a shallow rooted crop, **needs frequent light irrigation to maintain optimum soil moisture for proper growth and bulb development.** Rabi crop needs 10-15 irrigations at 7-10 days interval depending upon the soil moisture condition. There are chances of **purple blotch and Stemphylium blight** disease incidence in onion crop. To manage purple blotch and Stemphylium leaf blight disease in onion spray Mancozeb 75% WP @ 600gram/acre or Difenoconazole 25% E.C @ 200ml/acre or Tebuconazole 25.9% E.C @ 100ml/acre by mixing in 200 litre of water at 15 days interval

Fruits: To control **mealy bug infestation in papaya, guava, mango** wet the soil near the root zone of the plant with Chlorpyrifos @ 4 ml per litre of water and spray the same insecticide on plant @ 2 ml per litre of water; do not pluck fruits within 15 days of spraying

Mango: Present weather conditions are favourable for the incidence of **Mango hoppers**. To control go for spraying of phosphamidon @ 0.5ml/l (or) dimethoate 2ml/l (or) chlorpyrifos @ 2.5ml/l. during initial stages of flowering. Cloudy and foggy weather may aggravate the infestation of **powdery mildew** in Mango. Spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew before floral initiation.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-272
Date- 18/02/2022

AGROMET ADVISORY BULLETIN FOR BADAMBA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0.1 mm rainfall during the last week. The mean daily maximum and minimum temperature was 29.14°C (-2.6°C from normal) and 15.57°C (-2.3°C from normal), respectively. Mean relative humidity of the district was around 58-79%.

Forecasted period: Today(8.30IST) to 23th February (8.30IST)2022

Weather advisory: Very light rainfall is forecasted for Sunday and Dry weather is forecasted for other 4 days in the block. The day temperature will be 31-33°C and night temperature will be 16-18°C for upcoming 5 days. Maximum and minimum relative humidity will be 68-95% and 11-36% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Badamba (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	18/02/2022	19/02/2022	20/02/2022	21/02/2022	22/02/2022
Rainfall(mm)	0	0	2	0	0
T-Max(°C)	31	31	33	33	31
T-Min(°C)	18	18	18	16	16
Cloud Cover(octa)	3	2	3	5	0
RH Max (%)	87	90	83	68	95
RH Min (%)	36	27	15	10	11
Wind Speed (Kmph)	7	7	12	9	6
Wind Direction (deg)	158	162	207	326	45

Agromet Advisory

General advisory: Farmers should undertake field preparation for cultivation of different summer vegetables like sponge gourd, bottle gourd, pumpkin, water melon, musk melon etc. If seedlings are already raised in polythene bags; transplant in main field @ 2-3 seedlings per pit. Irrigation should be paused on Sunday as there will be rainfall is forecasted.

Pulses:

Forecasted weather may aggravate the infestation of **Flea beetle and powdery mildew** in pulses. Spray neem oil 5 ml/L of water to control Flea beetle and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew. There may be chances of **leaf crinkle disease** infestation due to past foggy and cold conditions. To manage the insect vector of the disease like whitefly and aphid spray Thiamethoxam 25% W. G @ 40 gram/acre or Acetamiprid 20% S.P. @ 50 gram/acre or Imidacloprid 17.8% SL @ 60-ml acre by mixing in 200 litre of water in 15 days interval.

Rabi-Groundnut:

Farmers are advised to do intercultivation and apply Gypsum @200 kg/ acre in Groundnut which is at flowering stage. forecasted warm and dry days with low RH favours infestation of **thrips**, to control thrips spray Thiamethoxam 25% W.G @ 40 gram/acre or Imidacloprid 17.8% SL @ 50-ml/acre by mixing it in 200 litre of water. For control of **Tobacco caterpillar and spodoterago** for spraying of fipronil 5% SL 400ml (or) indoxacarb 14.5% SC 80gm or neem oil 1000ml @ 200 liter of water per acre.

Mustard: To meet the deficiency of Sulfur applies 200 gram sulfur per acre. **Powdery Mildew:** Spray Sulfex 2 gram/lit. To control **white rust disease** in mustard, spray Copper oxychloride @ 3g per litre of water. **Painted bug and Aphid attack** may take place in mustard crop, apply Thiamethoxam 25% WG @ 3g per 10 litres of water.

Vegetables:

Thrips and mites are prominent pests in vegetables at present. To control both, spray Bifenthrin @1g/litre or Profenophos @2ml/litre of water. To control the **sucking pests like aphid & white fly**, spray Imidacloprid @ 5ml/10 lit of water.

Tomato:

- Transplanted seedlings are at 6-8 week stage in main field. Due to water stress in past, deficiency of boron or due to hormonal disbalance **flower drop occurs**. Apply timely and required amount of irrigation. Spray Boron 20% 2 gram/lit during early flowering and 15 days later. To manage **late blight** in tomato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days interval. To control **white fly** Spray Imidacloprid (18.7% S.L) 0.8 ml/lit of water or Acidamiprid 1 ml/lit of water or Diafenthiuron 12.5 gram/ 15 lit of water.

- Potato:** Farmers are advised to start earthing-up in potato if crop has attained a height of 15-22 cm. If so required, the practice may be repeated after 15 days. There are chances of **Potato Tuber Moth** infestation in potato. To manage this pest chemically spray Spinosad 45 % SC @ 75-ml/acre or Emamectin Benzoate 5% SG @ 80-gram/ acre or Chlorantraniliprole 18.5 % SC @80- ml/acre. To manage **late blight** in potato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days interval.

- Onion:** Do inter culture and hand weeding may be carried out at 45-50 DAT and application of 30-kg of urea and 15-kg of MOP fertilizer should be done at this stage. Onion being a shallow rooted crop, **needs frequent light irrigation to maintain optimum soil moisture for proper growth and bulb development**. Rabi crop needs 10-15 irrigations at 7-10 days interval depending upon the soil moisture condition. There are chances of **purple blotch and Stemphyllium blight** disease incidence in onion crop. To manage purple blotch and Stemphyllium leaf blight disease in onion spray Mancozeb 75% WP @ 600gram/acre or Difenoconazole 25% E.C @ 200ml/acre or Tebuconazole 25.9% E.C @ 100ml/acre by mixing in 200 litre of water at 15 days interval

- Fruits:** To control **mealy bug infestation** in papaya, guava, mango wet the soil near the root zone of the plant with Chlorpyrifos @ 4 ml per litre of water and spray the same insecticide on plant @ 2 ml per litre of water; do not pluck fruits within 15 days of spraying

- Mango:** Present weather conditions are favourable for the incidence of **Mango hoppers**. To control go for spraying of phosphamidon @ 0.5ml/l (or) dimethoate 2ml/l (or) chlorpyrifos @2.5ml/l. during initial stages of flowering. Cloudy and foggy weather may aggravate the infestation of **powdery mildew** in Mango. Spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew before floral initiation.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-272
Date- 18/02/2022

AGROMET ADVISORY BULLETIN FOR BANKI

District Past week average weather condition

The district received average 0.1 mm rainfall during the last week. The mean daily maximum and minimum temperature was 29.14°C (-2.6°C from normal) and 15.57°C (-2.3°C from normal), respectively. Mean relative humidity of the district was around 58-79%.

Forecasted period: Today(8.30IST) to 23th February (8.30IST)2022

Weather advisory: *Very light rainfall is forecasted for Sunday and Dry weather is forecasted for other 4 days in the block.* The day temperature will be 31-34°C and night temperature will be 17-19°C for upcoming 5 days. Maximum and minimum relative humidity will be 81-94% and 11-39% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Banki (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	18/02/2022	19/02/2022	20/02/2022	21/02/2022	22/02/2022
Rainfall(mm)	0	0	1	0	0
T-Max(°C)	31	31	34	33	31
T-Min(°C)	18	18	19	17	17
Cloud Cover(octa)	3	2	3	5	0
RH Max (%)	83	92	81	82	94
RH Min (%)	39	30	16	12	11
Wind Speed (Kmph)	7	9	14	9	6
Wind Direction (deg)	152	244	207	248	68

Agromet Advisory

General advisory: Farmers should undertake field preparation for cultivation of different summer vegetables like sponge gourd, bottle gourd, pumpkin, water melon, musk melon etc. If seedlings are already raised in polythene bags; transplant in main field @ 2-3 seedlings per pit. Irrigation should be paused on Sunday as there will be rainfall is forecasted.

Pulses:

Forecasted weather may aggravate the infestation of **Flea beetle and powdery mildew** in pulses. Spray neem oil 5 ml/L of water to control Flea beetle and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9% EC per litre water for controlling Powdery mildew. There may be chances of **leaf crinkle disease** infestation due to past foggy and cold conditions. To manage the insect vector of the disease like whitefly and aphid spray Thiamethoxam 25% W. G @ 40 gram/acre or Acetamiprid 20% S.P. @ 50 gram/acre or Imidacloprid 17.8% SL @ 60-ml acre by mixing in 200 litre of water in 15 days interval.

Rabi-Groundnut:

Farmers are advised to do intercultivation and apply Gypsum @200 kg/ acre in Groundnut which is at flowering stage. forecasted warm and dry days with low RH favours infestation of **thrips**, to control thrips spray Thiamethoxam 25% W.G @ 40 gram/acre or Imidacloprid 17.8% SL @ 50-ml/acre by mixing it in 200 litre of water. For control of **Tobacco caterpillar and spodotera** for spraying of fipronil 5% SL 400ml (or) indoxacarb 14.5% SC 80gm or neem oil 1000ml @ 200 liter of water per acre.

Mustard: To meet the deficiency of Sulfur applies 200 gram sulfur per acre. **Powdery Mildew:** Spray Sulfex 2 gram/lit. To control **white rust disease** in mustard, spray Copper oxychloride @ 3g per litre of water. **Painted bug and Aphid attack** may take place in mustard crop, apply Thiamethoxam 25% WG @ 3g per 10 litres of water.

Vegetables:

Thrips and mites are prominent pests in vegetables at present. To control both, spray Bifenthrin @1g/litre or Profenophos @2ml/litre of water. To control the sucking pests like aphid & white fly, spray Imidacloprid @ 5ml/10 lit of water.

Tomato:

- Transplanted seedlings are at 6-8 week stage in main field. Due to water stress in past, deficiency of boron or due to hormonal disbalance **flower drop occurs**. Apply timely and required amount of irrigation. Spray Boron 20% 2 gram/lit during early flowering and 15 days later. To manage **late blight** in tomato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days interval. To control **white fly** Spray Imidacloprid (18.7% S.L) 0.8 ml/lit of water or Acetamiprid 1 ml/lit of water or Diafenthiuron 12.5 gram/ 15 lit of water.

- Potato:** Farmers are advised to start earthing-up in potato if crop has attained a height of 15-22 cm. If so required, the practice may be repeated after 15 days. There are chances of **Potato Tuber Moth** infestation in potato. To manage this pest chemically spray Spinosad 45 % SC @ 75-ml/acre or Emamectin Benzoate 5% SG @ 80-gram/ acre or Chlorantraniliprole 18.5 % SC @80- ml/acre. To manage **late blight** in potato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days interval.

- Onion:** Do inter culture and hand weeding may be carried out at 45-50 DAT and application of 30-kg of urea and 15-kg of MOP fertilizer should be done at this stage. Onion being a shallow rooted crop, **needs frequent light irrigation to maintain optimum soil moisture for proper growth and bulb development**. Rabi crop needs 10-15 irrigations at 7-10 days interval depending upon the soil moisture condition. There are chances of **purple blotch and Stemphylium blight** disease incidence in onion crop. To manage purple blotch and Stemphylium leaf blight disease in onion spray Mancozeb 75% WP @ 600gram/acre or Difenoconazole 25% E.C @ 200ml/acre or Tebuconazole 25.9% E.C @ 100ml/acre by mixing in 200 litre of water at 15 days interval

- Fruits:** To control **mealy bug infestation in papaya, guava, mango** wet the soil near the root zone of the plant with Chlorpyrifos @ 4 ml per litre of water and spray the same insecticide on plant @ 2 ml per litre of water; do not pluck fruits within 15 days of spraying

- Mango:** Present weather conditions are favourable for the incidence of **Mango hoppers**. To control go for spraying of phosphamidon @ 0.5ml/l (or) dimethoate 2ml/l (or) chlorpyrifos @2.5ml/l. during initial stages of flowering. Cloudy and foggy weather may aggravate the infestation of **powdery mildew** in Mango. Spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9% EC per litre water for controlling Powdery mildew before floral initiation.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA INDIA METEOROLOGICAL DEPARTMENT DISTRICT AGROMET UNIT KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-272

Date- 18/02/2022

AGROMET ADVISORY BULLETIN FOR BANKI-DAMPARA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0.1 mm rainfall during the last week. The mean daily maximum and minimum temperature was 29.14°C (-2.6°C from normal) and 15.57°C (-2.3°C from normal), respectively. Mean relative humidity of the district was around 58-79%.

Forecasted period: Today(8.30IST) to 23th February (8.30IST)2022

Weather Advisory: Very light rainfall is forecasted for Sunday and Dry weather is forecasted for other 4 days in the block. The day temperature will be 31-33°C and night temperature will be 17-19°C for upcoming 5 days. Maximum and minimum relative humidity will be 80-94% and 11-39% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Banki-Dampara (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	18/02/2022	19/02/2022	20/02/2022	21/02/2022	22/02/2022
Rainfall(mm)	0	0	2	0	0
T-Max(°C)	31	31	33	33	31
T-Min(°C)	18	19	19	17	17
Cloud Cover(octa)	3	2	3	5	0
RH Max (%)	82	92	80	83	94
RH Min (%)	39	30	16	12	11
Wind Speed (Kmph)	7	9	14	9	6
Wind Direction (deg)	152	169	207	248	23

Agromet Advisory

General advisory: Farmers should undertake field preparation for cultivation of different summer vegetables like sponge gourd, bottle gourd, pumpkin, water melon, musk melon etc. If seedlings are already raised in polythene bags; transplant in main field @ 2-3 seedlings per pit. Irrigation should be paused on Sunday as there will be rainfall is forecasted.

Pulses:

Forecasted weather may aggravate the infestation of **Flea beetle and powdery mildew** in pulses. Spray neem oil 5 ml/L of water to control Flea beetle and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew. There may be chances of **leaf crinkle disease** infestation due to past foggy and cold conditions. To manage the insect vector of the disease like whitefly and aphid spray Thiamethoxam 25% W. G @ 40 gram/acre or Acetamiprid 20% S.P. @ 50 gram/acre or Imidacloprid 17.8% SL @ 60-ml acre by mixing in 200 litre of water in 15 days interval.

Rabi-Groundnut:

Farmers are advised to do intercultivation and apply Gypsum @200 kg/ acre in Groundnut which is at flowering stage. forecasted warm and dry days with low RH favours infestation of **thrips**, to control thrips spray Thiamethoxam 25% W.G @ 40 gram/acre or Imidacloprid 17.8% SL @ 50-ml/acre by mixing it in 200 litre of water. For control of **Tobacco caterpillar and spodoterago** for spraying of fipronil 5% SL 400ml (or) indoxacarb 14.5% SC 80gm or neem oil 1000ml @ 200 liter of water per acre.

Mustard: To meet the deficiency of Sulfur applies 200 gram sulfur per acre. **Powdery Mildew:** Spray Sulfex 2 gram/lit. To control **white rust disease** in mustard, spray Copper oxychloride @ 3g per litre of water. **Painted bug and Aphid attack** may take place in mustard crop, apply Thiamethoxam 25% WG @ 3g per 10 litres of water.

Vegetables:

Thrips and mites are prominent pests in vegetables at present. To control both, spray Bifenthrin @1g/litre or Profenophos @2ml/litre of water. To control the sucking pests like aphid & white fly, spray Imidacloprid @ 5ml/10 lit of water.

Tomato:

Transplanted seedlings are at 6-8 week stage in main field. Due to water stress in past, deficiency of boron or due to hormonal disbalance **flower drop occurs**. Apply timely and required amount of irrigation. Spray Boron 20% 2 gram/lit during early flowering and 15 days later. To manage **late blight** in tomato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days interval. To control **white fly** Spray Imidacloprid (18.7% S.L) 0.8 ml/lit of water or Acetamiprid 1 ml/lit of water or Diafenthiuron 12.5 gram/ 15 lit of water.

Potato: Farmers are advised to start earthing-up in potato if crop has attained a height of 15-22 cm. If so required, the practice may be repeated after 15 days. There are chances of **Potato Tuber Moth** infestation in potato. To manage this pest chemically spray Spinosad 45 % SC @ 75-ml/acre or Emamectin Benzoate 5% SG @ 80-gram/ acre or Chlorantraniliprole 18.5 % SC @80- ml/acre. To manage **late blight** in potato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days interval.

Onion: Do inter culture and hand weeding may be carried out at 45-50 DAT and application of 30-kg of urea and 15-kg of MOP fertilizer should be done at this stage. Onion being a shallow rooted crop, **needs frequent light irrigation to maintain optimum soil moisture for proper growth and bulb development**. Rabi crop needs 10-15 irrigations at 7-10 days interval depending upon the soil moisture condition. There are chances of **purple blotch and Stemphylium blight** disease incidence in onion crop. To manage purple blotch and Stemphylium leaf blight disease in onion spray Mancozeb 75% WP @ 600gram/acre or Difenoconazole 25% E.C @ 200ml/acre or Tebuconazole 25.9% E.C @ 100ml/acre by mixing in 200 litre of water at 15 days interval

Fruits: To control **mealy bug infestation** in papaya, guava, mango wet the soil near the root zone of the plant with Chlorpyrifos @ 4 ml per litre of water and spray the same insecticide on plant @ 2 ml per litre of water; do not pluck fruits within 15 days of spraying

Mango: Present weather conditions are favourable for the incidence of **Mango hoppers**. To control go for spraying of phosphamidon @ 0.5ml/l (or) dimethoate 2ml/l (or) chlorpyrifos @2.5ml/l. during initial stages of flowering. Cloudy and foggy weather may aggravate the infestation of **powdery mildew** in Mango. Spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew before floral initiation.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-272
Date- 18/02/2022

AGROMET ADVISORY BULLETIN FOR BARANGA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0.1 mm rainfall during the last week. The mean daily maximum and minimum temperature was 29.14°C (-2.6°C from normal) and 15.57°C (-2.3°C from normal), respectively. Mean relative humidity of the district was around 58-79%.

Forecasted period: Today(8.30IST) to 23th February (8.30IST)2022

Weather advisory: *Very light rainfall is forecasted for Sunday and Dry weather is forecasted for other 4 days in the block.* The day temperature will be 30-33°C and night temperature will be 17-20°C for upcoming 5 days. Maximum and minimum relative humidity will be 79-90% and 14-41% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Baranga (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	18/02/2022	19/02/2022	20/02/2022	21/02/2022	22/02/2022
Rainfall(mm)	0	0	3	0	0
T-Max(°C)	30	31	33	33	31
T-Min(°C)	19	19	20	17	17
Cloud Cover	2	2	4	5	0
RH Max (%)	75	90	79	89	89
RH Min (%)	41	35	19	15	14
Wind Speed (Kmph)	7	9	15	9	8
Wind Direction (deg)	150	203	214	360	27

Agromet Advisory

General advisory: Farmers should undertake field preparation for cultivation of different summer vegetables like sponge gourd, bottle gourd, pumpkin, water melon, musk melon etc. If seedlings are already raised in polythene bags; transplant in main field @ 2-3 seedlings per pit. Irrigation should be paused on Sunday as there will be rainfall is forecasted.

Pulses:

Forecasted weather may aggravate the infestation of **Flea beetle and powdery mildew** in pulses. Spray neem oil 5 ml/L of water to control Flea beetle and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9% EC per litre water for controlling Powdery mildew. There may be chances of **leaf crinkle disease** infestation due to past foggy and cold conditions. To manage the insect vector of the disease like whitefly and aphid spray Thiamethoxam 25% W. G @ 40 gram/acre or Acetamiprid 20% S.P. @ 50 gram/acre or Imidacloprid 17.8% SL @ 60-ml acre by mixing in 200 litre of water in 15 days interval.

Rabi-Groundnut:

Farmers are advised to do intercultivation and apply Gypsum @200 kg/ acre in Groundnut which is at flowering stage. forecasted warm and dry days with low RH favours infestation of **thrips**, to control thrips spray Thiamethoxam 25% W.G @ 40 gram/acre or Imidacloprid 17.8% SL @ 50-ml/acre by mixing it in 200 litre of water. For control of **Tobacco caterpillar and spodotera** go for spraying of fipronil 5% SL 400ml (or) indoxacarb 14.5% SC 80gm or neem oil 1000ml @ 200 liter of water per acre.

Mustard: To meet the deficiency of Sulfur applies 200 gram sulfur per acre. **Powdery Mildew:** Spray Sulfex 2 gram/lit. To control **white rust disease** in mustard, spray Copper oxychloride @ 3g per litre of water. **Painted bug and Aphid attack** may take place in mustard crop, apply Thiamethoxam 25% WG @ 3g per 10 litres of water.

Vegetables:

Thrips and mites are prominent pests in vegetables at present. To control both, spray Bifenthrin @ 1g/litre or Profenophos @ 2ml/litre of water. To control the **sucking pests like aphid & white fly**, spray Imidacloprid @ 5ml/10 lit of water.

- **Tomato:**
- Transplanted seedlings are at 6-8 week stage in main field. **Due to water stress in past, deficiency of boron or due to hormonal disbalance flower drop occurs.** Apply timely and required amount of irrigation. Spray Boron 20% 2 gram/lit during early flowering and 15 days later. To manage **late blight** in tomato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days interval. To control **white fly** Spray Imidacloprid (18.7% S.L) 0.8 ml/lit of water or Acetamiprid 1 ml/lit of water or Diafenthiuron 12.5 gram/ 15 lit of water.
- **Potato:** Farmers are advised to start earthing-up in potato if crop has attained a height of 15-22 cm. If so required, the practice may be repeated after 15 days. There are chances of **Potato Tuber Moth** infestation in potato. To manage this pest chemically spray Spinosad 45 % SC @ 75-ml/acre or Emamectin Benzoate 5% SG @ 80-gram/ acre or Chlorantraniliprole 18.5 % SC @ 80- ml/acre. To manage **late blight** in potato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days interval.
- **Onion:** Do inter culture and hand weeding may be carried out at 45-50 DAT and application of 30-kg of urea and 15-kg of MOP fertilizer should be done at this stage. Onion being a shallow rooted crop, **needs frequent light irrigation to maintain optimum soil moisture for proper growth and bulb development.** Rabi crop needs 10-15 irrigations at 7-10 days interval depending upon the soil moisture condition. There are chances of **purple blotch and Stemphylium blight** disease incidence in onion crop. To manage purple blotch and Stemphylium leaf blight disease in onion spray Mancozeb 75% WP @ 600gram/acre or Difenconazole 25% E.C @ 200ml/acre or Tebuconazole 25.9% E.C @ 100ml/acre by mixing in 200 litre of water at 15 days interval
- **Fruits:** To control **mealy bug infestation in papaya, guava, mango** wet the soil near the root zone of the plant with Chlorpyrifos @ 4 ml per litre of water and spray the same insecticide on plant @ 2 ml per litre of water; do not pluck fruits within 15 days of spraying
- **Mango:** Present weather conditions are favourable for the incidence of **Mango hoppers**. To control go for spraying of phosphamidon @ 0.5ml/l (or) dimethoate 2ml/l (or) chlorpyrifos @ 2.5ml/l. during initial stages of flowering. Cloudy and foggy weather may aggravate the infestation of **powdery mildew** in Mango. Spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9% EC per litre water for controlling Powdery mildew before floral initiation.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)



Bulletin No.-272
Date- 18/02/2022

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

AGROMET ADVISORY BULLETIN FOR CUTTACK SADAR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0.1 mm rainfall during the last week. The mean daily maximum and minimum temperature was 29.14°C (-2.6°C from normal) and 15.57°C (-2.3°C from normal), respectively. Mean relative humidity of the district was around 58-79%.

Forecasted period: Today(8.30IST) to 23th February (8.30IST)2022

Weather advisory: *Very light rainfall is forecasted for Sunday and Dry weather is forecasted for other 4 days in the block.* The day temperature will be 30-33°C and night temperature will be 17-20°C for upcoming 5 days. Maximum and minimum relative humidity will be 73-91% and 14-40% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Cuttack sadar (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	18/02/2022	19/02/2022	20/02/2022	21/02/2022	22/02/2022
Rainfall(mm)	0	0	4	0	0
T-Max(°C)	30	31	33	33	31
T-Min(°C)	19	19	20	18	17
Cloud Cover(octa)	2	2	4	5	0
RH Max (%)	73	91	79	84	88
RH Min (%)	40	32	17	14	14
Wind Speed (Kmph)	6	9	14	9	8
Wind Direction (deg)	158	195	207	345	34

Agromet Advisory

General advisory: Farmers should undertake field preparation for cultivation of different summer vegetables like sponge gourd, bottle gourd, pumpkin, water melon, musk melon etc. If seedlings are already raised in polythene bags; transplant in main field @ 2-3 seedlings per pit. Irrigation should be paused on Sunday as there will be rainfall is forecasted.

Pulses:

Forecasted weather may aggravate the infestation of **Flea beetle and powdery mildew** in pulses. Spray neem oil 5 ml/L of water to control Flea beetle and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9% EC per litre water for controlling Powdery mildew. There may be chances of **leaf crinkle disease** infestation due to past foggy and cold conditions. To manage the insect vector of the disease like whitefly and aphid spray Thiamethoxam 25% W. G @ 40 gram/acre or Acetamiprid 20% S.P. @ 50 gram/acre or Imidacloprid 17.8% SL @ 60-ml acre by mixing in 200 litre of water in 15 days interval.

Rabi-Groundnut:

Farmers are advised to do intercultivation and apply Gypsum @200 kg/ acre in Groundnut which is at flowering stage. forecasted warm and dry days with low RH favours infestation of **thrips**, to control thrips spray Thiamethoxam 25% W.G @ 40 gram/acre or Imidacloprid 17.8% SL @ 50-ml/acre by mixing it in 200 litre of water. For control of **Tobacco caterpillar and spodotera** go for spraying of fipronil 5% SL 400ml (or) indoxacarb 14.5% SC 80gm or neem oil 1000ml @ 200 liter of water per acre.

Mustard: To meet the deficiency of Sulfur applies 200 gram sulfur per acre. **Powdery Mildew:** Spray Sulfex 2 gram/lit. To control **white rust disease** in mustard, spray Copper oxychloride @ 3g per litre of water. **Painted bug and Aphid attack** may take place in mustard crop, apply Thiamethoxam 25% WG @ 3g per 10 litres of water.

Vegetables:

Thrips and mites are prominent pests in vegetables at present. To control both, spray Bifenthrin @ 1g/litre or Profenophos @ 2ml/litre of water. To control the sucking pests like aphid & white fly, spray Imidacloprid @ 5ml/10 lit of water.

- Tomato:**
- Transplanted seedlings are at 6-8 week stage in main field. Due to water stress in past, deficiency of boron or due to hormonal disbalance **flower drop occurs**. Apply timely and required amount of irrigation. Spray Boron 20% 2 gram/lit during early flowering and 15 days later. To manage **late blight** in tomato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days interval. To control **white fly** Spray Imidacloprid (18.7% S.L) 0.8 ml/lit of water or Acetamiprid 1 ml/lit of water or Diafenthiuron 12.5 gram/ 15 lit of water.
- Potato:** Farmers are advised to start earthing-up in potato if crop has attained a height of 15-22 cm. If so required, the practice may be repeated after 15 days. There are chances of **Potato Tuber Moth** infestation in potato. To manage this pest chemically spray Spinosad 45 % SC @ 75-ml/acre or Emamectin Benzoate 5% SG @ 80-gram/ acre or Chlorantraniliprole 18.5 % SC @ 80- ml/acre. To manage **late blight** in potato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days interval.
- Onion:** Do inter culture and hand weeding may be carried out at 45-50 DAT and application of 30-kg of urea and 15-kg of MOP fertilizer should be done at this stage. Onion being a shallow rooted crop, **needs frequent light irrigation to maintain optimum soil moisture for proper growth and bulb development**. Rabi crop needs 10-15 irrigations at 7-10 days interval depending upon the soil moisture condition. There are chances of **purple blotch and Stemphyllium blight** disease incidence in onion crop. To manage purple blotch and Stemphyllium leaf blight disease in onion spray Mancozeb 75% WP @ 600gram/acre or Difenoconazole 25% E.C @ 200ml/acre or Tebuconazole 25.9% E.C @ 100ml/acre by mixing in 200 litre of water at 15 days interval
- Fruits:** To control **mealy bug infestation in papaya, guava, mango** wet the soil near the root zone of the plant with Chlorpyrifos @ 4 ml per litre of water and spray the same insecticide on plant @ 2 ml per litre of water; do not pluck fruits within 15 days of spraying
- Mango:** Present weather conditions are favourable for the incidence of **Mango hoppers**. To control go for spraying of phosphamidon @ 0.5ml/l (or) dimethoate 2ml/l (or) chlorpyrifos @ 2.5ml/l. during initial stages of flowering. Cloudy and foggy weather may aggravate the infestation of **powdery mildew** in Mango. Spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9% EC per litre water for controlling Powdery mildew before floral initiation.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-272
Date- 18/02/2022

AGROMET ADVISORY BULLETIN FOR KANTAPADA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 2mm and district received average 0.1 mm rainfall during the last week. The mean daily maximum and minimum temperature was 29.14°C (-2.6°C from normal) and 15.57°C (-2.3°C from normal), respectively. Mean relative humidity of the district was around 58-79%.

Forecasted period: Today(8.30IST) to 23th February (8.30IST)2022

Weather advisory: Light rainfall is forecasted for Sunday and dry weather is forecasted for other 4 days in the block. The day temperature will be 30-32°C and night temperature will be 17-20°C for upcoming 5 days. Maximum and minimum relative humidity will be 76-95% and 18-42% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Kantapada (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	18/02/2022	19/02/2022	20/02/2022	21/02/2022	22/02/2022
Rainfall(mm)	0	0	6	0	0
T-Max(°C)	30	31	32	32	30
T-Min(°C)	19	20	20	18	17
Cloud Cover(octa)	2	2	4	5	0
RH Max (%)	76	88	80	95	87
RH Min (%)	42	39	25	18	18
Wind Speed (Kmph)	7	10	17	9	9
Wind Direction (deg)	30	198	210	15	23

Agromet Advisory

General advisory: Farmers should undertake field preparation for cultivation of different summer vegetables like sponge gourd, bottle gourd, pumpkin, water melon, musk melon etc. If seedlings are already raised in polythene bags; transplant in main field @ 2-3 seedlings per pit. Irrigation should be paused on Sunday as there will be rainfall is forecasted.

Pulses:

Forecasted weather may aggravate the infestation of **Flea beetle and powdery mildew** in pulses. Spray neem oil 5 ml/L of water to control Flea beetle and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew. There may be chances of **leaf crinkle disease** infestation due to past foggy and cold conditions. To manage the insect vector of the disease like whitefly and aphid spray Thiamethoxam 25% W. G @ 40 gram/acre or Acetamiprid 20% S.P. @ 50 gram/acre or Imidacloprid 17.8% SL @ 60-ml acre by mixing in 200 litre of water in 15 days interval.

Rabi-Groundnut:

Farmers are advised to do intercultivation and apply Gypsum @200 kg/ acre in Groundnut which is at flowering stage. forecasted warm and dry days with low RH favours infestation of **thrips**, to control thrips spray Thiamethoxam 25% W.G @ 40 gram/acre or Imidacloprid 17.8% SL @ 50-ml/acre by mixing it in 200 litre of water. For control of **Tobacco caterpillar and spodoterago** for spraying of fipronil 5% SL 400ml (or) indoxacarb 14.5% SC 80gm or neem oil 1000ml @ 200 liter of water per acre.

Mustard: To meet the deficiency of Sulfur applies 200 gram sulfur per acre. **Powdery Mildew:** Spray Sulfex 2 gram/lit. To control **white rust disease** in mustard, spray Copper oxychloride @ 3g per litre of water. **Painted bug and Aphid attack** may take place in mustard crop, apply Thiamethoxam 25% WG @ 3g per 10 litres of water.

Vegetables:

Thrips and mites are prominent pests in vegetables at present. To control both, spray Bifenthrin @1g/litre or Profenophos @2ml/litre of water. To control the **sucking pests like aphid & white fly**, spray Imidacloprid @ 5ml/10 lit of water.

Tomato:

- Transplanted seedlings are at 6-8 week stage in main field. Due to water stress in past, deficiency of boron or due to hormonal disbalance **flower drop occurs**. Apply timely and required amount of irrigation. Spray Boron 20% 2 gram/lit during early flowering and 15 days later. To manage **late blight** in tomato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days interval. To control **white fly** Spray Imidacloprid (18.7% S.L) 0.8 ml/lit of water or Acetamiprid 1 ml/lit of water or Diafenthiuron 12.5 gram/ 15 lit of water.

- Potato:** Farmers are advised to start earthing-up in potato if crop has attained a height of 15-22 cm. If so required, the practice may be repeated after 15 days. There are chances of **Potato Tuber Moth** infestation in potato. To manage this pest chemically spray Spinosad 45 % SC @ 75-ml/acre or Emamectin Benzoate 5% SG @ 80-gram/ acre or Chlorantraniliprole 18.5 % SC @80- ml/acre. To manage **late blight** in potato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days interval.

- Onion:** Do inter culture and hand weeding may be carried out at 45-50 DAT and application of 30-kg of urea and 15-kg of MOP fertilizer should be done at this stage. Onion being a shallow rooted crop, **needs frequent light irrigation to maintain optimum soil moisture for proper growth and bulb development**. Rabi crop needs 10-15 irrigations at 7-10 days interval depending upon the soil moisture condition. There are chances of **purple blotch and Stemphylium blight** disease incidence in onion crop. To manage purple blotch and Stemphylium leaf blight disease in onion spray Mancozeb 75% WP @ 600gram/acre or Difenconazole 25% E.C @ 200ml/acre or Tebuconazole 25.9% E.C @ 100ml/acre by mixing in 200 litre of water at 15 days interval

- Fruits:** To control **mealy bug infestation** in papaya, guava, mango wet the soil near the root zone of the plant with Chlorpyrifos @ 4 ml per litre of water and spray the same insecticide on plant @ 2 ml per litre of water; do not pluck fruits within 15 days of spraying

- Mango:** Present weather conditions are favourable for the incidence of **Mango hoppers**. To control go for spraying of phosphamidon @ 0.5ml/l (or) dimethoate 2ml/l (or) chlorpyrifos @2.5ml/l. during initial stages of flowering. Cloudy and foggy weather may aggravate the infestation of **powdery mildew** in Mango. Spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew before floral initiation.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)



Bulletin No.-272
Date- 18/02/2022

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

AGROMET ADVISORY BULLETIN FOR MAHANGA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0.1 mm rainfall during the last week. The mean daily maximum and minimum temperature was 29.14°C (-2.6°C from normal) and 15.57°C (-2.3°C from normal), respectively. Mean relative humidity of the district was around 58-79%.

Forecasted period: Today(8.30IST) to 23th February (8.30IST)2022

Weather advisory: Light rainfall is forecasted for Sunday and dry weather is forecasted for other 4 days in the block. The day temperature will be and it will be 30-32°C and night temperature will be 17-20°C for upcoming 5 days. Maximum and minimum relative humidity will be 70-89% and 18-35% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Mahanga (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	18/02/2022	19/02/2022	20/02/2022	21/02/2022	22/02/2022
Rainfall(mm)	0	0	6	0	0
T-Max(°C)	31	31	32	32	30
T-Min(°C)	19	19	20	17	17
Cloud Cover(octal)	2	2	4	5	0
RH Max (%)	70	89	77	87	76
RH Min (%)	35	35	20	18	21
Wind Speed (Kmph)	8	10	15	11	10
Wind Direction (deg)	338	195	214	18	26

Agromet Advisory

General advisory: Farmers should undertake field preparation for cultivation of different summer vegetables like sponge gourd, bottle gourd, pumpkin, water melon, musk melon etc. If seedlings are already raised in polythene bags; transplant in main field @ 2-3 seedlings per pit. Irrigation should be paused on Sunday as there will be rainfall is forecasted.

Pulses:

Forecasted weather may aggravate the infestation of **Flea beetle and powdery mildew** in pulses. Spray neem oil 5 ml/L of water to control Flea beetle and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew. There may be chances of **leaf crinkle disease** infestation due to past foggy and cold conditions. To manage the insect vector of the disease like whitefly and aphid spray Thiamethoxam 25% W. G @ 40 gram/acre or Acetamiprid 20% S.P. @ 50 gram/acre or Imidacloprid 17.8% SL @ 60-ml acre by mixing in 200 litre of water in 15 days interval.

Rabi-Groundnut:

Farmers are advised to do intercultivation and apply Gypsum @200 kg/ acre in Groundnut which is at flowering stage. forecasted warm and dry days with low RH favours infestation of **thrips**, to control thrips spray Thiamethoxam 25% W.G @ 40 gram/acre or Imidacloprid 17.8% SL @ 50-ml/acre by mixing it in 200 litre of water. For control of **Tobacco caterpillar and spodoterago** for spraying of fipronil 5% SL 400ml (or) indoxacarb 14.5% SC 80gm or neem oil 1000ml @ 200 liter of water per acre.

Mustard: To meet the deficiency of Sulfur applies 200 gram sulfur per acre. **Powdery Mildew:** Spray Sulfex 2 gram/lit. To control **white rust disease** in mustard, spray Copper oxychloride @ 3g per litre of water. **Painted bug and Aphid attack** may take place in mustard crop, apply Thiamethoxam 25% WG @ 3g per 10 litres of water.

Vegetables:

Thrips and mites are prominent pests in vegetables at present. To control both, spray Bifenthrin @1g/litre or Profenophos @2ml/litre of water. To control the **sucking pests like aphid & white fly**, spray Imidacloprid @ 5ml/10 lit of water.

Tomato:

- Transplanted seedlings are at 6-8 week stage in main field. Due to water stress in past, deficiency of boron due to hormonal disbalance **flower drop occurs**. Apply timely and required amount of irrigation. Spray Boron 20% 2 gram/lit during early flowering and 15 days later. To manage **late blight** in tomato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days interval. To control **white fly** Spray Imidacloprid (18.7% S.L) 0.8 ml/lit of water or Acetamiprid 1 ml/lit of water or Diafenthiuron 12.5 gram/ 15 lit of water.

- Potato:** Farmers are advised to start earthing-up in potato if crop has attained a height of 15-22 cm. If so required, the practice may be repeated after 15 days. There are chances of **Potato Tuber Moth** infestation in potato. To manage this pest chemically spray Spinosad 45 % SC @ 75-ml/acre or Emamectin Benzoate 5% SG @ 80-gram/ acre or Chlorantraniliprole 18.5 % SC @80- ml/acre. To manage **late blight** in potato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days interval.

- Onion:** Do inter culture and hand weeding may be carried out at 45-50 DAT and application of 30-kg of urea and 15-kg of MOP fertilizer should be done at this stage. Onion being a shallow rooted crop, **needs frequent light irrigation to maintain optimum soil moisture for proper growth and bulb development**. Rabi crop needs 10-15 irrigations at 7-10 days interval depending upon the soil moisture condition. There are chances of **purple blotch and Stemphylium blight** disease incidence in onion crop. To manage purple blotch and Stemphylium leaf blight disease in onion spray Mancozeb 75% WP @ 600gram/acre or Difenconazole 25% E.C @ 200ml/acre or Tebuconazole 25.9% E.C @ 100ml/acre by mixing in 200 litre of water at 15 days interval

- Fruits:** To control **mealy bug infestation** in papaya, guava, mango wet the soil near the root zone of the plant with Chlorpyrifos @ 4 ml per litre of water and spray the same insecticide on plant @ 2 ml per litre of water; do not pluck fruits within 15 days of spraying

- Mango:** Present weather conditions are favourable for the incidence of **Mango hoppers**. To control go for spraying of phosphamidon @ 0.5ml/l (or) dimethoate 2ml/l (or) chlorpyrifos @2.5ml/l. during initial stages of flowering. Cloudy and foggy weather may aggravate the infestation of **powdery mildew** in Mango. Spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew before floral initiation.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)



Bulletin No.-272
Date- 18/02/2022

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

AGROMET ADVISORY BULLETIN FOR NARASINGHPUR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0.1 mm rainfall during the last week. The mean daily maximum and minimum temperature was 29.14°C (-2.6°C from normal) and 15.57°C (-2.3°C from normal), respectively. Mean relative humidity of the district was around 58-79%.

Forecasted period: Today(8.30IST) to 23th February (8.30IST)2022

Weather advisory: Very light rainfall is forecasted for Sunday and Dry weather is forecasted for other 4 days in the block. The day temperature will be 30-33°C and night temperature will be 15-17°C for upcoming 5 days. Maximum and minimum relative humidity will be 45-93% and 10-28% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Narasinghpur (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	18/02/2022	19/02/2022	20/02/2022	21/02/2022	22/02/2022
Rainfall(mm)	0	0	2	0	0
T-Max(°C)	33	31	32	32	30
T-Min(°C)	17	17	17	15	15
Cloud Cover(octa)	3	2	2	5	0
RH Max (%)	93	85	76	45	93
RH Min (%)	28	23	17	10	10
Wind Speed (Kmph)	6	5	9	9	6
Wind Direction (deg)	158	165	333	315	23

Agromet Advisory

General advisory: Farmers should undertake field preparation for cultivation of different summer vegetables like sponge gourd, bottle gourd, pumpkin, water melon, musk melon etc. If seedlings are already raised in polythene bags; transplant in main field @ 2-3 seedlings per pit. Irrigation should be paused on Sunday as there will be rainfall is forecasted.

Pulses:

Forecasted weather may aggravate the infestation of **Flea beetle and powdery mildew** in pulses. Spray neem oil 5 ml/L of water to control Flea beetle and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew. There may be chances of **leaf crinkle disease** infestation due to past foggy and cold conditions. To manage the insect vector of the disease like whitefly and aphid spray Thiamethoxam 25% W. G @ 40 gram/acre or Acetamiprid 20% S.P. @ 50 gram/acre or Imidacloprid 17.8% SL @ 60-ml acre by mixing in 200 litre of water in 15 days interval.

Rabi-Groundnut:

Farmers are advised to do intercultivation and apply Gypsum @200 kg/ acre in Groundnut which is at flowering stage. forecasted warm and dry days with low RH favours infestation of **thrips**, to control thrips spray Thiamethoxam 25% W.G @ 40 gram/acre or Imidacloprid 17.8% SL @ 50-ml/acre by mixing it in 200 litre of water. For control of **Tobacco caterpillar and spodoterago** for spraying of fipronil 5% SL 400ml (or) indoxacarb 14.5% SC 80gm or neem oil 1000ml @ 200 liter of water per acre.

Mustard: To meet the deficiency of Sulfur applies 200 gram sulfur per acre. **Powdery Mildew:** Spray Sulfex 2 gram/lit. To control **white rust disease** in mustard, spray Copper oxychloride @ 3g per litre of water. **Painted bug and Aphid attack** may take place in mustard crop, apply Thiamethoxam 25% WG @ 3g per 10 litres of water.

Vegetables:

Thrips and mites are prominent pests in vegetables at present. To control both, spray Bifenthrin @1g/litre or Profenophos @2ml/litre of water. To control the **sucking pests like aphid & white fly**, spray Imidacloprid @ 5ml/10 lit of water.

Tomato:

- Transplanted seedlings are at 6-8 week stage in main field. Due to water stress in past, deficiency of boron or due to hormonal disbalance **flower drop occurs**. Apply timely and required amount of irrigation. Spray Boron 20% 2 gram/lit during early flowering and 15 days later. To manage **late blight** in tomato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days interval. To control **white fly** Spray Imidacloprid (18.7% S.L) 0.8 ml/lit of water or Acetamiprid 1 ml/lit of water or Diafenthiuron 12.5 gram/ 15 lit of water.

- Potato:** Farmers are advised to start earthing-up in potato if crop has attained a height of 15-22 cm. If so required, the practice may be repeated after 15 days. There are chances of **Potato Tuber Moth** infestation in potato. To manage this pest chemically spray Spinosad 45 % SC @ 75-ml/acre or Emamectin Benzoate 5% SG @ 80-gram/ acre or Chlorantraniliprole 18.5 % SC @80- ml/acre. To manage **late blight** in potato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days interval.

- Onion:** Do inter culture and hand weeding may be carried out at 45-50 DAT and application of 30-kg of urea and 15-kg of MOP fertilizer should be done at this stage. Onion being a shallow rooted crop, **needs frequent light irrigation to maintain optimum soil moisture for proper growth and bulb development**. Rabi crop needs 10-15 irrigations at 7-10 days interval depending upon the soil moisture condition. There are chances of **purple blotch and Stemphyllium blight** disease incidence in onion crop. To manage purple blotch and Stemphyllium leaf blight disease in onion spray Mancozeb 75% WP @ 600gram/acre or Difenoconazole 25% E.C @ 200ml/acre or Tebuconazole 25.9% E.C @ 100ml/acre by mixing in 200 litre of water at 15 days interval

- Fruits:** To control **mealy bug infestation** in papaya, guava, mango wet the soil near the root zone of the plant with Chlorpyrifos @ 4 ml per litre of water and spray the same insecticide on plant @ 2 ml per litre of water; do not pluck fruits within 15 days of spraying

- Mango:** Present weather conditions are favourable for the incidence of **Mango hoppers**. To control go for spraying of phosphamidon @ 0.5ml/l (or) dimethoate 2ml/l (or) chlorpyrifos @2.5ml/l. during initial stages of flowering. Cloudy and foggy weather may aggravate the infestation of **powdery mildew** in Mango. Spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew before floral initiation.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-272
Date- 18/02/2022

AGROMET ADVISORY BULLETIN FOR NIALI

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0.1 mm rainfall during the last week. The mean daily maximum and minimum temperature was 29.14°C (-2.6°C from normal) and 15.57°C (-2.3°C from normal), respectively. Mean relative humidity of the district was around 58-79%.

Forecasted period: Today(8.30IST) to 23th February (8.30IST)2022

Weather advisory: *Very light rainfall is forecasted for Sunday and Dry weather is forecasted for other 4 days in the block.* The day temperature will be 30-32°C and night temperature will be 17-20°C for upcoming 5 days. Maximum and minimum relative humidity will be 77-93% and 20-44% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Niali (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	18/02/2022	19/02/2022	20/02/2022	21/02/2022	22/02/2022
Rainfall(mm)	0	0	1	0	0
T-Max(°C)	30	31	32	32	31
T-Min(°C)	19	20	20	18	17
Cloud Cover(octa)	2	2	4	5	0
RH Max (%)	77	86	82	93	87
RH Min (%)	44	40	32	20	22
Wind Speed(Kmph)	8	11	18	10	10
Wind Direction(deg)	116	189	205	15	45

Agromet Advisory

General advisory: Farmers should undertake field preparation for cultivation of different summer vegetables like sponge gourd, bottle gourd, pumpkin, water melon, musk melon etc. If seedlings are already raised in polythene bags; transplant in main field @ 2-3 seedlings per pit. Irrigation should be paused on Sunday as there will be rainfall is forecasted.

Pulses:

Forecasted weather may aggravate the infestation of **Flea beetle and powdery mildew** in pulses. Spray neem oil 5 ml/L of water to control Flea beetle and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew. There may be chances of **leaf crinkle disease** infestation due to past foggy and cold conditions. To manage the insect vector of the disease like whitefly and aphid spray Thiamethoxam 25% W. G @ 40 gram/acre or Acetamiprid 20% S.P. @ 50 gram/acre or Imidacloprid 17.8% SL @ 60-ml acre by mixing in 200 litre of water in 15 days interval.

Rabi-Groundnut:

Farmers are advised to do intercultivation and apply Gypsum @200 kg/ acre in Groundnut which is at flowering stage. forecasted warm and dry days with low RH favours infestation of **thrips**, to control thrips spray Thiamethoxam 25% W.G @ 40 gram/acre or Imidacloprid 17.8% SL @ 50-ml/acre by mixing it in 200 litre of water. For control of **Tobacco caterpillar and spodoterago** for spraying of fipronil 5% SL 400ml (or) indoxacarb 14.5% SC 80gm or neem oil 1000ml @ 200 liter of water per acre.

Mustard: To meet the deficiency of Sulfur applies 200 gram sulfur per acre. **Powdery Mildew:** Spray Sulfex 2 gram/lit. To control **white rust disease** in mustard, spray Copper oxychloride @ 3g per litre of water. **Painted bug and Aphid attack** may take place in mustard crop, apply Thiamethoxam 25% WG @ 3g per 10 litres of water.

Vegetables:

Thrips and mites are prominent pests in vegetables at present. To control both, spray Bifenthrin @1g/litre or Profenophos @2ml/litre of water. To control the **sucking pests like aphid & white fly**, spray Imidacloprid @ 5ml/10 lit of water.

Tomato:

Transplanted seedlings are at 6-8 week stage in main field. **Due to water stress in past, deficiency of boron due to hormonal disbalance flower drop occurs.** Apply timely and required amount of irrigation. Spray Boron 20% 2 gram/lit during early flowering and 15 days later. To manage **late blight** in tomato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days interval. To control **white fly** Spray Imidacloprid (18.7% S.L) 0.8 ml/lit of water or Acetamiprid 1 ml/lit of water or Diafenthiuron 12.5 gram/ 15 lit of water.

Potato: Farmers are advised to start earthing-up in potato if crop has attained a height of 15-22 cm. If so required, the practice may be repeated after 15 days. There are chances of **Potato Tuber Moth** infestation in potato. To manage this pest chemically spray Spinosad 45 % SC @ 75-ml/acre or Emamectin Benzoate 5% SG @ 80-gram/ acre or Chlorantraniliprole 18.5 % SC @80- ml/acre. To manage **late blight** in potato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days interval.

Onion: Do inter culture and hand weeding may be carried out at 45-50 DAT and application of 30-kg of urea and 15-kg of MOP fertilizer should be done at this stage. Onion being a shallow rooted crop, **needs frequent light irrigation to maintain optimum soil moisture for proper growth and bulb development.** Rabi crop needs 10-15 irrigations at 7-10 days interval depending upon the soil moisture condition. There are chances of **purple blotch and Stemphylium blight** disease incidence in onion crop. To manage purple blotch and Stemphylium leaf blight disease in onion spray Mancozeb 75% WP @ 600gram/acre or Difenconazole 25% E.C @ 200ml/acre or Tebuconazole 25.9% E.C @ 100ml/acre by mixing in 200 litre of water at 15 days interval

Fruits: To control **mealy bug infestation** in papaya, guava, mango wet the soil near the root zone of the plant with Chlorpyrifos @ 4 ml per litre of water and spray the same insecticide on plant @ 2 ml per litre of water; do not pluck fruits within 15 days of spraying

Mango: Present weather conditions are favourable for the incidence of **Mango hoppers**. To control go for spraying of phosphamidon @ 0.5ml/l (or) dimethoate 2ml/l (or) chlorpyrifos @2.5ml/l. during initial stages of flowering. Cloudy and foggy weather may aggravate the infestation of **powdery mildew** in Mango. Spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew before floral initiation.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-272
Date- 18/02/2022

AGROMET ADVISORY BULLETIN FOR NISCHINTAKOILI

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0.1 mm rainfall during the last week. The mean daily maximum and minimum temperature was 29.14°C (-2.6°C from normal) and 15.57°C (-2.3°C from normal), respectively. Mean relative humidity of the district was around 58-79%.

Forecasted period: Today(8.30IST) to 23th February (8.30IST)2022

Weather advisory: Light rainfall is forecasted for Sunday and dry weather is forecasted for other 4 days in the block. The day temperature will be 30-32°C and night temperature will be 17-20°C for upcoming 5 days. Maximum and minimum relative humidity will be 71-92% and 18-37% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Nischintakoili (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	18/02/2022	19/02/2022	20/02/2022	21/02/2022	22/02/2022
Rainfall(mm)	0	0	6	0	0
T-Max(°C)	31	31	32	32	30
T-Min(°C)	19	19	20	18	17
Cloud Cover(octa)	2	2	4	5	0
RH Max (%)	71	89	76	92	78
RH Min (%)	36	37	21	18	21
Wind Speed (Kmph)	8	10	16	10	10
Wind Direction (deg)	135	203	225	15	26

Agromet Advisory

General advisory: Farmers should undertake field preparation for cultivation of different summer vegetables like sponge gourd, bottle gourd, pumpkin, water melon, musk melon etc. If seedlings are already raised in polythene bags; transplant in main field @ 2-3 seedlings per pit. Irrigation should be paused on Sunday as there will be rainfall is forecasted.

Pulses:

Forecasted weather may aggravate the infestation of Flea beetle and powdery mildew in pulses. Spray neem oil 5 ml/L of water to control Flea beetle and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew. There may be chances of leaf crinkle disease infestation due to past foggy and cold conditions. To manage the insect vector of the disease like whitefly and aphid spray Thiamethoxam 25% W. G @ 40 gram/acre or Acetamiprid 20% S.P. @ 50 gram/acre or Imidacloprid 17.8% SL @ 60-ml acre by mixing in 200 litre of water in 15 days interval.

Rabi-Groundnut:

Farmers are advised to do intercultivation and apply Gypsum @200 kg/ acre in Groundnut which is at flowering stage. forecasted warm and dry days with low RH favours infestation of thrips, to control thrips spray Thiamethoxam 25% W.G @ 40 gram/acre or Imidacloprid 17.8% SL @ 50-ml/acre by mixing it in 200 litre of water. For control of Tobacco caterpillar and spodotera go for spraying of fipronil 5% SL 400ml (or) indoxacarb 14.5% SC 80gm or neem oil 1000ml @ 200 liter of water per acre.

Mustard: To meet the deficiency of Sulfur applies 200 gram sulfur per acre. Powdery Mildew: Spray Sulfex 2 gram/lit. To control white rust disease in mustard, spray Copper oxychloride @ 3g per litre of water. Painted bug and Aphid attack may take place in mustard crop, apply Thiamethoxam 25% WG @ 3g per 10 litres of water.

Vegetables:

Thrips and mites are prominent pests in vegetables at present. To control both, spray Bifenthrin @1g/litre or Profenophos @2ml/litre of water. To control the sucking pests like aphid & white fly, spray Imidacloprid @ 5ml/10 lit of water.

Tomato:

Transplanted seedlings are at 6-8 week stage in main field. Due to water stress in past, deficiency of boron or due to hormonal disbalance flower drop occurs. Apply timely and required amount of irrigation. Spray Boron 20% 2 gram/lit during early flowering and 15 days later. To manage late blight in tomato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days interval. To control white fly Spray Imidacloprid (18.7% S.L) 0.8 ml/lit of water or Acetamiprid 1 ml/lit of water or Diafenthiuron 12.5 gram/ 15 lit of water.

Potato: Farmers are advised to start earthing-up in potato if crop has attained a height of 15-22 cm. If so required, the practice may be repeated after 15 days. There are chances of Potato Tuber Moth infestation in potato. To manage this pest chemically spray Spinosad 45 % SC @ 75-ml/acre or Emamectin Benzoate 5% SG @ 80-gram/ acre or Chlorantraniliprole 18.5 % SC @80- ml/acre. To manage late blight in potato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days interval.

Onion: Do inter culture and hand weeding may be carried out at 45-50 DAT and application of 30-kg of urea and 15-kg of MOP fertilizer should be done at this stage. Onion being a shallow rooted crop, needs frequent light irrigation to maintain optimum soil moisture for proper growth and bulb development. Rabi crop needs 10-15 irrigations at 7-10 days interval depending upon the soil moisture condition. There are chances of purple blotch and Stemphylium blight disease incidence in onion crop. To manage purple blotch and Stemphylium leaf blight disease in onion spray Mancozeb 75% WP @ 600gram/acre or Difenoconazole 25% E.C @ 200ml/acre or Tebuconazole 25.9% E.C @ 100ml/acre by mixing in 200 litre of water at 15 days interval

Fruits: To control mealy bug infestation in papaya, guava, mango wet the soil near the root zone of the plant with Chlorpyrifos @ 4 ml per litre of water and spray the same insecticide on plant @ 2 ml per litre of water; do not pluck fruits within 15 days of spraying

Mango: Present weather conditions are favourable for the incidence of Mango hoppers. To control go for spraying of phosphamidon @ 0.5ml/l (or) dimethoate 2ml/l (or) chlorpyrifos @2.5ml/l. during initial stages of flowering. Cloudy and foggy weather may aggravate the infestation of powdery mildew in Mango. Spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew before floral initiation.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-272
Date- 18/02/2022

AGROMET ADVISORY BULLETIN FOR SALEPUR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0.1 mm rainfall during the last week. The mean daily maximum and minimum temperature was 29.14°C (-2.6°C from normal) and 15.57°C (-2.3°C from normal), respectively. Mean relative humidity of the district was around 58-79%.

Forecasted period: Today(8.30IST) to 23rd February (8.30IST)2022

Weather advisory: *Very light rainfall is forecasted for Sunday and Dry weather is forecasted for other 4 days in the block.* The day temperature will be 30-33°C and night temperature will be 18-20°C for upcoming 5 days. Maximum and minimum relative humidity will be 71-91% and 18-37% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Salepur (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	18/02/2022	19/02/2022	20/02/2022	21/02/2022	22/02/2022
Rainfall(mm)	0	0	5	0	0
T-Max(°C)	31	30	33	32	30
T-Min(°C)	19	19	20	18	18
Cloud Cover(octa)	2	2	4	5	0
RH Max (%)	71	89	77	91	79
RH Min (%)	37	36	20	18	20
Wind Speed (Kmph)	7	10	15	10	9
Wind Direction (deg)	158	203	225	15	26

Agromet Advisory

General advisory: Farmers should undertake field preparation for cultivation of different summer vegetables like sponge gourd, bottle gourd, pumpkin, water melon, musk melon etc. If seedlings are already raised in polythene bags; transplant in main field @ 2-3 seedlings per pit. Irrigation should be paused on Sunday as there will be rainfall is forecasted.

Pulses:

Forecasted weather may aggravate the infestation of **Flea beetle and powdery mildew** in pulses. Spray neem oil 5 ml/L of water to control Flea beetle and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew. There may be chances of **leaf crinkle disease** infestation due to past foggy and cold conditions. To manage the insect vector of the disease like whitefly and aphid spray Thiamethoxam 25% W. G @ 40 gram/acre or Acetamiprid 20% S.P. @ 50 gram/acre or Imidacloprid 17.8% SL @ 60-ml acre by mixing in 200 litre of water in 15 days interval.

Rabi-Groundnut:

Farmers are advised to do intercultivation and apply Gypsum @200 kg/ acre in Groundnut which is at flowering stage. forecasted warm and dry days with low RH favours infestation of **thrips**, to control thrips spray Thiamethoxam 25% W.G @ 40 gram/acre or Imidacloprid 17.8% SL @ 50-ml/acre by mixing it in 200 litre of water. For control of **Tobacco caterpillar and spodoterago** for spraying of fipronil 5% SL 400ml (or) indoxacarb 14.5% SC 80gm or neem oil 1000ml @ 200 liter of water per acre.

Mustard: To meet the deficiency of Sulfur applies 200 gram sulfur per acre. **Powdery Mildew:** Spray Sulfex 2 gram/lit. To control **white rust disease** in mustard, spray Copper oxychloride @ 3g per litre of water. **Painted bug and Aphid attack** may take place in mustard crop, apply Thiamethoxam 25% WG @ 3g per 10 litres of water.

Vegetables:

Thrips and mites are prominent pests in vegetables at present. To control both, spray Bifenthrin @1g/litre or Profenophos @2ml/litre of water. To control the **sucking pests like aphid & white fly**, spray Imidacloprid @ 5ml/10 lit of water.

Tomato:

Transplanted seedlings are at 6-8 week stage in main field. Due to water stress in past, deficiency of boron due to hormonal disbalance **flower drop occurs**. Apply timely and required amount of irrigation. Spray Boron 20% 2 gram/lit during early flowering and 15 days later. To manage **late blight** in tomato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days interval. To control **white fly** Spray Imidacloprid (18.7% S.L) 0.8 ml/lit of water or Acetamiprid 1 ml/lit of water or Diafenthiuron 12.5 gram/ 15 lit of water.

Potato: Farmers are advised to start earthing-up in potato if crop has attained a height of 15-22 cm. If so required, the practice may be repeated after 15 days. There are chances of **Potato Tuber Moth** infestation in potato. To manage this pest chemically spray Spinosad 45 % SC @ 75-ml/acre or Emamectin Benzoate 5% SG @ 80-gram/ acre or Chlorantraniliprole 18.5 % SC @80- ml/acre. To manage **late blight** in potato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days interval.

Onion: Do inter culture and hand weeding may be carried out at 45-50 DAT and application of 30-kg of urea and 15-kg of MOP fertilizer should be done at this stage. Onion being a shallow rooted crop, **needs frequent light irrigation to maintain optimum soil moisture for proper growth and bulb development**. Rabi crop needs 10-15 irrigations at 7-10 days interval depending upon the soil moisture condition. There are chances of **purple blotch and Stemphylium blight** disease incidence in onion crop. To manage purple blotch and Stemphylium leaf blight disease in onion spray Mancozeb 75% WP @ 600gram/acre or Difenconazole 25% E.C @ 200ml/acre or Tebuconazole 25.9% E.C @ 100ml/acre by mixing in 200 litre of water at 15 days interval

Fruits: To control **mealy bug infestation** in papaya, guava, mango wet the soil near the root zone of the plant with Chlorpyrifos @ 4 ml per litre of water and spray the same insecticide on plant @ 2 ml per litre of water; do not pluck fruits within 15 days of spraying

Mango: Present weather conditions are favourable for the incidence of **Mango hoppers**. To control go for spraying of phosphamidon @ 0.5ml/l (or) dimethoate 2ml/l (or) chlorpyrifos @2.5ml/l. during initial stages of flowering. Cloudy and foggy weather may aggravate the infestation of **powdery mildew** in Mango. Spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew before floral initiation.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA INDIA METEOROLOGICAL DEPARTMENT DISTRICT AGROMET UNIT KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

AGROMET ADVISORY BULLETIN FOR TANGI-CHOUDWAR



Bulletin No.-272

Date- 18/02/2022

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0.1 mm rainfall during the last week. The mean daily maximum and minimum temperature was 29.14°C (-2.6°C from normal) and 15.57°C (-2.3°C from normal), respectively. Mean relative humidity of the district was around 58-79%.

Forecasted period: Today(8.30IST) to 23rd February (8.30IST)2022

Weather advisory: Light rainfall is forecasted for Sunday and dry weather is forecasted for other 4 days in the block. The day temperature will be 29-32°C and night temperature will be 21-23°C for upcoming 5 days. Maximum and minimum relative humidity will be 74-90% and 34-44% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Tangi-Choudwar (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	18/02/2022	19/02/2022	20/02/2022	21/02/2022	22/02/2022
Rainfall(mm)	0	0	6	0	0
T-Max(°C)	29	30	32	32	30
T-Min(°C)	21	23	23	22	21
Cloud Cover(octa)	1	2	4	5	0
RH Max (%)	74	82	83	90	79
RH Min (%)	44	44	44	34	41
Wind Speed (Kmph)	11	13	22	12	12
Wind Direction (deg)	116	180	208	15	23

Agromet Advisory

General advisory: Farmers should undertake field preparation for cultivation of different summer vegetables like sponge gourd, bottle gourd, pumpkin, water melon, musk melon etc. If seedlings are already raised in polythene bags; transplant in main field @ 2-3 seedlings per pit. Irrigation should be paused on Sunday as there will be rainfall is forecasted.

Pulses:

Forecasted weather may aggravate the infestation of **Flea beetle and powdery mildew** in pulses. Spray neem oil 5 ml/L of water to control Flea beetle and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew. There may be chances of **leaf crinkle disease** infestation due to past foggy and cold conditions. To manage the insect vector of the disease like whitefly and aphid spray Thiamethoxam 25% W. G @ 40 gram/acre or Acetamiprid 20% S.P. @ 50 gram/acre or Imidacloprid 17.8% SL @ 60-ml acre by mixing in 200 litre of water in 15 days interval.

Rabi-Groundnut:

Farmers are advised to do intercultivation and apply Gypsum @200 kg/ acre in Groundnut which is at flowering stage. forecasted warm and dry days with low RH favours infestation of **thrips**, to control thrips spray Thiamethoxam 25% W.G @ 40 gram/acre or Imidacloprid 17.8% SL @ 50-ml/acre by mixing it in 200 litre of water. For control of **Tobacco caterpillar and spodoterago** for spraying of fipronil 5% SL 400ml (or) indoxacarb 14.5% SC 80gm or neem oil 1000ml @ 200 liter of water per acre.

Mustard: To meet the deficiency of Sulfur applies 200 gram sulfur per acre. **Powdery Mildew:** Spray Sulfex 2 gram/lit. To control **white rust disease** in mustard, spray Copper oxychloride @ 3g per litre of water. **Painted bug and Aphid attack** may take place in mustard crop, apply Thiamethoxam 25% WG @ 3g per 10 litres of water.

Vegetables:

Thrips and mites are prominent pests in vegetables at present. To control both, spray Bifenthrin @1g/litre or Profenophos @2ml/litre of water. To control the sucking pests like aphid & white fly, spray Imidacloprid @ 5ml/10 lit of water.

Tomato:

Transplanted seedlings are at 6-8 week stage in main field. Due to water stress in past, deficiency of boron or due to hormonal disbalance **flower drop occurs**. Apply timely and required amount of irrigation. Spray Boron 20% 2 gram/lit during early flowering and 15 days later. To manage **late blight** in tomato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days interval. To control **white fly** Spray Imidacloprid (18.7% S.L) 0.8 ml/lit of water or Acetamiprid 1 ml/lit of water or Diafenthiuron 12.5 gram/ 15 lit of water.

Potato: Farmers are advised to start earthing-up in potato if crop has attained a height of 15-22 cm. If so required, the practice may be repeated after 15 days. There are chances of **Potato Tuber Moth** infestation in potato. To manage this pest chemically spray Spinosad 45 % SC @ 75-ml/acre or Emamectin Benzoate 5% SG @ 80-gram/ acre or Chlorantraniliprole 18.5 % SC @80- ml/acre. To manage **late blight** in potato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days interval.

Onion: Do inter culture and hand weeding may be carried out at 45-50 DAT and application of 30-kg of urea and 15-kg of MOP fertilizer should be done at this stage. Onion being a shallow rooted crop, **needs frequent light irrigation to maintain optimum soil moisture for proper growth and bulb development**. Rabi crop needs 10-15 irrigations at 7-10 days interval depending upon the soil moisture condition. There are chances of **purple blotch and Stemphylium blight** disease incidence in onion crop. To manage purple blotch and Stemphylium leaf blight disease in onion spray Mancozeb 75% WP @ 600gram/acre or Difenoconazole 25% E.C @ 200ml/acre or Tebuconazole 25.9% E.C @ 100ml/acre by mixing in 200 litre of water at 15 days interval

Fruits: To control **mealy bug infestation** in papaya, guava, mango wet the soil near the root zone of the plant with Chlorpyrifos @ 4 ml per litre of water and spray the same insecticide on plant @ 2 ml per litre of water; do not pluck fruits within 15 days of spraying

Mango: Present weather conditions are favourable for the incidence of **Mango hoppers**. To control go for spraying of phosphamidon @ 0.5ml/l (or) dimethoate 2ml/l (or) chlorpyrifos @2.5ml/l. during initial stages of flowering. Cloudy and foggy weather may aggravate the infestation of **powdery mildew** in Mango. Spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew before floral initiation.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-272
Date- 18/02/2022

AGROMET ADVISORY BULLETIN FOR TIGIRIA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0.1 mm rainfall during the last week. The mean daily maximum and minimum temperature was 29.14°C (-2.6°C from normal) and 15.57°C (-2.3°C from normal), respectively. Mean relative humidity of the district was around 58-79%.

Forecasted period: Today(8.30IST) to 23th February (8.30IST)2022

Weather advisory: Very light rainfall is forecasted for Sunday and Dry weather is forecasted for other 4 days in the block. The day temperature will be 31-33°C and night temperature will be 17-19°C for upcoming 5 days. Maximum and minimum relative humidity will be 71-94% and 11-39% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Tigiria (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	18/02/2022	19/02/2022	20/02/2022	21/02/2022	22/02/2022
Rainfall(mm)	0	0	2	0	0
T-Max(°C)	31	31	33	33	31
T-Min(°C)	18	18	19	17	17
Cloud Cover(octa)	3	2	4	5	0
RH Max (%)	81	91	77	71	94
RH Min (%)	39	28	16	12	11
Wind Speed (Kmph)	7	8	12	9	7
Wind Direction (deg)	155	180	207	326	45

Agro met Advisory

General advisory: Farmers should undertake field preparation for cultivation of different summer vegetables like sponge gourd, bottle gourd, pumpkin, water melon, musk melon etc. If seedlings are already raised in polythene bags; transplant in main field @ 2-3 seedlings per pit. Irrigation should be paused on Sunday as there will be rainfall is forecasted.

Pulses:

Forecasted weather may aggravate the infestation of **Flea beetle and powdery mildew** in pulses. Spray neem oil 5 ml/L of water to control Flea beetle and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9% EC per litre water for controlling Powdery mildew. There may be chances of **leaf crinkle disease** infestation due to past foggy and cold conditions. To manage the insect vector of the disease like whitefly and aphid spray Thiamethoxam 25% W. G @ 40 gram/acre or Acetamiprid 20% S.P. @ 50 gram/acre or Imidacloprid 17.8% SL @ 60-ml acre by mixing in 200 litre of water in 15 days interval.

Rabi-Groundnut:

Farmers are advised to do intercultivation and apply Gypsum @200 kg/ acre in Groundnut which is at flowering stage. forecasted warm and dry days with low RH favours infestation of **thrips**, to control thrips spray Thiamethoxam 25% W.G @ 40 gram/acre or Imidacloprid 17.8% SL @ 50-ml/acre by mixing it in 200 litre of water. For control of **Tobacco caterpillar and spodotera** for spraying of fipronil 5% SL 400ml (or) indoxacarb 14.5% SC 80gm or neem oil 1000ml @ 200 liter of water per acre.

Mustard: To meet the deficiency of Sulfur applies 200 gram sulfur per acre. **Powdery Mildew:** Spray Sulfex 2 gram/lit. To control **white rust disease** in mustard, spray Copper oxychloride @ 3g per litre of water. **Painted bug and Aphid attack** may take place in mustard crop, apply Thiamethoxam 25% WG @ 3g per 10 litres of water.

Vegetables:

Thrips and mites are prominent pests in vegetables at present. To control both, spray Bifenthrin @ 1g/litre or Profenophos @ 2ml/litre of water. To control the sucking pests like aphid & white fly, spray Imidacloprid @ 5ml/10 lit of water.

Tomato:

Transplanted seedlings are at 6-8 week stage in main field. Due to water stress in past, deficiency of boron or due to hormonal disbalance **flower drop occurs**. Apply timely and required amount of irrigation. Spray Boron 20% 2 gram/lit during early flowering and 15 days later. To manage **late blight** in tomato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days interval. To control **white fly** Spray Imidacloprid (18.7% S.L) 0.8 ml/lit of water or Acetamiprid 1 ml/lit of water or Diafenthiuron 12.5 gram/ 15 lit of water.

Potato: Farmers are advised to start earthing-up in potato if crop has attained a height of 15-22 cm. If so required, the practice may be repeated after 15 days. There are chances of **Potato Tuber Moth** infestation in potato. To manage this pest chemically spray Spinosad 45 % SC @ 75-ml/acre or Emamectin Benzoate 5% SG @ 80-gram/ acre or Chlorantraniliprole 18.5 % SC @ 80- ml/acre. To manage **late blight** in potato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days interval.

Onion: Do inter culture and hand weeding may be carried out at 45-50 DAT and application of 30-kg of urea and 15-kg of MOP fertilizer should be done at this stage. Onion being a shallow rooted crop, **needs frequent light irrigation to maintain optimum soil moisture for proper growth and bulb development**. Rabi crop needs 10-15 irrigations at 7-10 days interval depending upon the soil moisture condition. There are chances of **purple blotch and Stemphylium blight** disease incidence in onion crop. To manage purple blotch and Stemphylium leaf blight disease in onion spray Mancozeb 75% WP @ 600gram/acre or Difenconazole 25% E.C @ 200ml/acre or Tebuconazole 25.9% E.C @ 100ml/acre by mixing in 200 litre of water at 15 days interval

Fruits: To control **mealy bug infestation in papaya, guava, mango** wet the soil near the root zone of the plant with Chlorpyrifos @ 4 ml per litre of water and spray the same insecticide on plant @ 2 ml per litre of water; do not pluck fruits within 15 days of spraying

Mango: Present weather conditions are favourable for the incidence of **Mango hoppers**. To control go for spraying of phosphamidon @ 0.5ml/l (or) dimethoate 2ml/l (or) chlorpyrifos @ 2.5ml/l. during initial stages of flowering. Cloudy and foggy weather may aggravate the infestation of **powdery mildew** in Mango. Spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9% EC per litre water for controlling Powdery mildew before floral initiation.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-272
 Date- 18/02/2022

				
MEGHDOOT APP	DAMINI APP	FACEBOOK (Damu Cuttack)	BLOCK SPECIFIC WHAT'S APP GROUPS	@DAMUCUTTACK

- Kindly install and use **“Meghdoot mobile App”** for Cuttack district weather forecast, warnings and weather based agromet advisory services.
<https://play.google.com/store/apps/details?id=com.aas.meghdoot>
- Kindly install and use **“Damini mobile App”** for location specific for lightening forecast and warnings.
<https://play.google.com/store/apps/details?id=com.lightening.live.damini>
- Follow the **block specific whats app groups** and join in your block wise groups for getting weather based agromet advisory services twice in a week (Tuesday and Friday) and other weather warning based farm management advisories.
- Follow the **Official Website links** to get the weather based agromet advisory service bulletins (**English & Odia**) twice in a week (Tuesday and Friday).

1	ICAR-NRRI, Cuttack, Odisha	https://icar-nrri.in/agromate/	
2	Agrimet division, IMD, Pune	https://imdagrimet.gov.in/imdproject/AGIndex.php	

Dr Sujata Sethy
 (Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
 SMS (Agrometeorology)
 Contact No-9861445080/7008453427