




GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com




Bulletin No.-289
Date- 19/04/2022

AGROCLIMATIC ZONES OF CUTTACK DISTRICT:



STANDARD PRECIPITATION INDEX AND NDVI MAP



LEGEND (Medium range block level rainfall forecast)

	No Rainfall	00 mm
	Very light rainfall	01-05 mm
	Light rainfall	06-10 mm
	Light to moderate rainfall	11-30 mm
	Moderate to heavy rainfall	31-65 mm
	Heavy to very heavy rainfall	66-115 mm
	Extremely heavy rainfall	>116 mm

Dr Sujata Sethy
 (Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com






Bulletin No.-289
Date- 19/04/2022

Past week (12-18 April, 2022) cumulative rainfall departure% from normal and (13 -19 April) average maximum temperature and its departure from normal for Cuttack district



Medium range (Today (8.30IST) to Sunday (8.30IST)) Rainfall (mm) forecast for Cuttack district



Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-289
Date- 19/04/2022

Met Sub division	Extended range outlook for subsequent 1 week (6 th to 12 th days) (24-30 April 2022)		
Odisha	<i>Parameter</i>	<i>Departure (%)</i>	<i>Category</i>
	Rainfall	-66.89	Above Normal
	<i>Parameter</i>	<i>Anomaly</i>	<i>Category</i>
	Maximum temperature	6.70	Moderately Above Normal
	Minimum temperature	2.93	Above Normal

General Agromet Advisory:- As There will be below normal rainfall and moderately above normal temperature is forecasted in subsequent week, after upcoming 5 days, so farmers are advised to go for irrigation activities in the subsequent week.

Long range forecast for the 2021 South-west Monsoon seasonal rainfall

- Southwest monsoon seasonal (June to September) rainfall over the country as a whole is most likely to be normal (96 to 104 % of Long Period Average (LPA)).
- Quantitatively, the monsoon seasonal (June to September) rainfall is likely to be 99% of the Long Period Average (LPA) with a model error of $\pm 5\%$. The LPA of the season rainfall over the country as a whole for the period 1971-2020 is 87 cm.
- *Hence Farmers should prepare the crop plan depending upon the land type, soil fertility and Variety availability, irrigation facility.*

Tercile probability rainfall forecast for 2022 southwest monsoon season




Fig.1.Probability forecast of tercile categories* (below normal, normal, and above normal) for the seasonal rainfall over India during the 2022 southwest monsoon season (June -September). The figure illustrates the most likely categories as well as their probabilities. The white shaded areas represent climatological probabilities for all the tercile categories. The probabilities were derived using the MME forecast prepared from a group of four best coupled climate models. (* Tercile categories have equal climatological probabilities, of 33.33% each).

Dr Sujata Sethy
(Nodal officer)



Follow Meghadoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427














GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-289
Date- 19/04/2022

Weather (Past and Forecasted) based Disease and pest in Major Crops and vegetables

Sl No	Crop	Stage and condition of the crop	Anticipated Disease and pest	
1	Pulses (Green gram and black gram)	Flowering to pod development		Cercospera Leaf spot
				Leaf blight
				Rust
				Wilting
2	Maize	Vegetative		Maize stalk rot
				Red hairy cattrpillar
3	Brinjal	Fruiting		Fruit and shoot borer
				Phomopsis Blight
				Epilachna Beetle/Hadda beetle
				Cercospera leaf spot
				Helicoverpa caterpillar

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area
















Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-289
Date- 19/04/2022

4	Tomato	Fruting		Bacterial wilt
5	Okra	Vegetative to fruiting		Yellow vein mosaiv(YMV)
6	Cucurbits	Flowering to Fruiting stage		Red Pumpkin bettle
				Leaf spot
				Late blight
				Stem Rot
				Downy Mildew
				Fruit rot
7	Banana			Banana leaf spot/sigatoka
				Pseudo stem weevil
8	Mango	Fruiting stage		Fruit borer
				Fruit drop
				Bacterial Canker

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA INDIA METEOROLOGICAL DEPARTMENT DISTRICT AGROMET UNIT KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-289

Date- 19/04/2022

AGROMET ADVISORY BULLETIN FOR ATHAGAD

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0mm rainfall during the last week. The mean daily maximum and minimum temperature was **38.2°C (1°C from normal)** and **25.5°C (+1.4°C from normal)**, respectively. Mean relative humidity of the district was around 66-76%.

Forecasted period: Today(8.30IST) to 24th April (8.30IST)2022

Weather advisory: *Very light to light rainfall / Thundershower is forecasted for Thursday and Friday and Dry weather is forecasted for other 3 days in the block.* The day temperature will be **32-37°C** and night temperature will be **25-28°C** for upcoming 5 days. Maximum and minimum relative humidity will be 80-89% and 19-38% respectively. **There will be clear to partly cloudy sky for next 5 days.**

Block	Athagad (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	19/04/2022	20/04/2022	21/04/2022	22/04/2022	23/04/2022
Rainfall (mm)	0	0	6	2	0
T-Max (°C)	37	36	32	36	35
T-Min (°C)	25	26	27	28	27
Cloud Cover (octa)	1	1	3	3	2
RH Max (%)	89	86	82	80	86
RH Min (%)	19	23	38	29	26
Wind Speed (kmph)	18	23	25	18	16
Wind Direction (deg)	198	198	198	198	198

Agromet Advisory

General advisory: After completion of harvesting of rabi crops and before attending ploughing operation, farmers are advised to collect the soil sample from the field & send the sample to nearest soil testing laboratory for nutrient analysis. Soil testing should be done once in three years in rainfed situation and every three cropping season in irrigated situation.

Pulses: Harvest matured pulses before rainy day. **Bacterial leaf blight** are likely to increase in present situation in pulses due to realized rainfall; if observe, spray Streptomycin or Plantomycin @ 2 g/litre of water as foliar treatment. **Rust infestation** likely to occur in pulses, if observe spray Mancozeb 35.0%SC 3gram/ltr of water. Present weather is congenial for **Cercospora leaf spot** infestation in black gram crop. On occurrence, immediately spray Agrimycin or Streptomycin 2 g/litre, at an interval of 12-15 days, to effectively reduce pest population.

Summer Maize:

If there was infestation of **Stem rot** at vegetative stage as rainfall was realized in past, farmers may drench the crop root with Mancozeb 35.0%SC 3gram/ltr of water or validamycin 2ml/litre of water and there may increase the infestation of **Red hairy caterpillar** in maize after rainfall, to minimize the infestation farmers may spray Neem oil 5ml/ltr of water or Chlorpyrifos 20 % EC @ 2ml/litre of water.

Vegetables:

Okra: To control **Yellow Vain Mosaic Virus (YVMV)** in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease.

Brinjal: There are chances of infestation of **Epilachna Beetle/Haddabeetle** in Brinjal and cucurbits crop; to manage Epilachna Beetle in cucurbits, spray neem-based pesticide (Azadirachtin) 1500 PPM @ 3ml/litre of water at early stage of infestation; to manage Epilachna Beetle chemically, spray Chlorpyrifos 20 % EC @ 2ml/litre of water or Profenophos 50 % EC @ 2ml/litre of water.

FRUIT AND SHOOT BORER IN BRINJAL: Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@250g/acre at 15 days interval. If **phomopsis blight** will occur in brinjal, then spray zineb75%WP 2gram/litre of water. **Cercospora leaf spot** may be found after rainfall, if observe, then spray Carbendazim or zineb75%WP 2gram/litre of water. If **helicoverpa** infestation will be found, spray Flunendiamide or indoxacarb 2ml/litre of water.

Cucurbits: To manage **Red Pumpkin Beetle** in Cucurbit crops spray Cyantranilprole 10.26% OD @ 400-ml/acre or Imidacloprid 17.8% SL @ 60-ml/acre or Chlorpyrifos 20% EC @ 400-ml/acre. **Late Blight** may happen due to rising day temperature. There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Don't go for weeding. The grasses and weeds will help in maintaining the humidity and it is safe for creeping of the vines and does not allow the fruit to come in contact with soil and avoid fruit rot. High temperature leads to fungal wilting in pointed gourd, spray 1.5 ml Carbendazim in 1 litre water. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxchloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Banana: For **leaf spot/sigatoka** as a precautionary measure spray carbendazim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalnil @ 2.0 g/ litre of water. For **pseudo stem weevil** Suggested to give stem injection with 5 ml Chlorpyrifos or Dimethoate or spray 2.5 ml of Chlorpyrifos per litre of water.

Mango: To control **fruit drop**, spray Borax @2gm per lit. of water and apply Planofix @2ml per 5 lit. of water. For control of mango **fruit borer** spray Neem oil 3ml + chlorpyrifos 1 ml perlitre of water along with stickers at marble stage of the fruit. The spraying should be preferably carried out during evening hours. To control **bacterial canker**, three sprays of streptomycin (100 ppm) or Agrimycin-100 (100 ppm) after first visual symptom at 10- days intervals is recommended.

Animal husbandry: Make necessary arrangement for clean and sufficient drinking water for the animals.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA INDIA METEOROLOGICAL DEPARTMENT DISTRICT AGROMET UNIT KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-289

Date- 19/04/2022

AGROMET ADVISORY BULLETIN FOR BADAMBA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0 mm rainfall during the last week. The mean daily maximum and minimum temperature was **38.2°C (1°C from normal)** and **25.5°C (+1.4°C from normal)**, respectively. Mean relative humidity of the district was around 66-76%.

Forecasted period: Today(8.30IST) to 24th April(8.30IST)2022

Weather advisory: *Very light to light rainfall / Thundershower is forecasted for Thursday and Friday and Dry weather is forecasted for other 3 days in the block.* The day temperature will be **34-39°C** and night temperature will be **25-28°C** for upcoming 5 days. Maximum and minimum relative humidity will be 74-87% and 17-34% respectively. **There will be clear to partly cloudy sky for next 5 days.**

Block	Badamba (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	19/04/2022	20/04/2022	21/04/2022	22/04/2022	23/04/2022
Rainfall(mm)	0	0	6	2	0
T-Max(°C)	39	38	34	37	37
T-Min(°C)	25	25	27	28	27
Cloud Cover(octa)	1	1	3	2	2
RH Max (%)	87	83	80	74	85
RH Min (%)	17	22	34	25	23
Wind Speed (Kmph)	14	19	20	16	13
Wind Direction (deg)	193	180	167	167	180

Agromet Advisory

General advisory: After completion of harvesting of rabi crops and before attending ploughing operation, farmers are advised to collect the soil sample from the field & send the sample to nearest soil testing laboratory for nutrient analysis. Soil testing should be done once in three years in rainfed situation and every three cropping season in irrigated situation.

Pulses: Harvest matured pulses before rainy day. Bacterial leaf blight are likely to increase in present situation in pulses due to realized rainfall; if observe, spray Streptomycin or Plantomycin @ 2 g/litre of water as foliar treatment. Rust infestation likely to occur in pulses, if observe spray Mancozeb 35.0%SC 3gram/ltr of water. Present weather is congenial for **Cercospora leaf spot** infestation in black gram crop. On occurrence, immediately spray Agrimycin or Streptomycin 2 g/litre, at an interval of 12-15 days, to effectively reduce pest population.

Summer Maize:

If there was infestation of **Stem rot** at vegetative stage as rainfall was realized in past, farmers may drench the crop root with Mancozeb 35.0%SC 3gram/ltr of water or validamycin 2ml/litre of water and there may increase the infestation of **Red hairy caterpillar** in maize after rainfall, to minimize the infestation farmers may spray Neem oil 5ml/ltr of water or Chlorpyrifos 20 % EC @ 2ml/litre of water.

Vegetables:

Okra: To control **Yellow Vain Mosaic Virus (YVMV)** in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease.

Brinjal: There are chances of infestation of **Epilachna Beetle/Haddabeetle** in Brinjal and cucurbits crop; to manage Epilachna Beetle in cucurbits, spray neem-based pesticide (Azadirachtin) 1500 PPM @ 3ml/litre of water at early stage of infestation; to manage Epilachna Beetle chemically, spray Chlorpyrifos 20 % EC @ 2ml/litre of water or Profenophos 50 % EC @ 2ml/litre of water.

FRUIT AND SHOOT BORER IN BRINJAL: Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@250g/acre at 15 days interval. If **phomopsis blight** will occur in brinjal, then spray zineb 75%WP 2gram/ltr of water. **Cercospora leaf spot** may be found after rainfall, if observe, then spray Carbendazim or zineb 75%WP 2gram/ltr of water. If **helicopter infestation** will be found, spray Flunendiamide or indoxacarb 2ml/ltr of water.

Cucurbits: To manage **Red Pumpkin Beetle** in Cucurbit crops spray Cyantraniliprole 10.26% OD @ 400-ml/acre or Imidacloprid 17.8% SL @ 60-ml/acre or Chlorpyrifos 20% EC @ 400-ml/acre. **Late Blight** may happen due to rising day temperature. There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Don't go for weeding. The grasses and weeds will help in maintaining the humidity and it is safe for creeping of the vines and does not allow the fruit to come in contact with soil and avoid fruit rot. High temperature leads to fungal wilting in pointed gourd, spray 1.5 ml Carbendazim in 1 litre water. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxychloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Banana: For **leaf spot/sigatoka** as a precautionary measure spray carbendazim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalanyl @ 2.0 g / litre of water. For **pseudo stem weevil** Suggested to give stem injection with 5 ml Chlorpyrifos or Dimethoate or spray 2.5 ml of Chlorpyrifos per litre of water.

Mango: To control **fruit drop**, spray Borax @ 2gm per lit. of water and apply Planofix @ 2ml per 5 lit. of water. For control of mango **fruit borer** spray Neem oil 3ml + chlorpyrifos 1 ml per litre of water along with stickers at marble stage of the fruit. The spraying should be preferably carried out during evening hours. To control **bacterial canker**, three sprays of streptomycin (100 ppm) or Agrimycin-100 (100 ppm) after first visual symptom at 10- days intervals is recommended.

Animal husbandry: Make necessary arrangement for clean and sufficient drinking water for the animals.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-289

Date- 19/04/2022

AGROMET ADVISORY BULLETIN FOR BANKI

District Past week average weather condition

The district received average 0 mm rainfall during the last week. The mean daily maximum and minimum temperature was **38.2°C (1°C from normal)** and 25.5°C (+1.4°C from normal), respectively. Mean relative humidity of the district was around 66-76%.

Forecasted period: Today(8.30IST) to 24th April(8.30IST)2022

Weather advisory: *Very light to light rainfall / Thundershower is forecasted for Thursday and Friday and Dry weather is forecasted for other 3 days in the block.* The day temperature will be 33-38°C and night temperature will be 25-27°C for upcoming 5 days. Maximum and minimum relative humidity will be 78-89% and 20-40% respectively. **There will be clear to partly cloudy sky for next 5 days.**

Block	Banki (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	19/04/2022	20/04/2022	21/04/2022	22/04/2022	23/04/2022
Rainfall(mm)	0	0	6	2	0
T-Max(°C)	38	37	33	37	37
T-Min(°C)	25	26	27	27	26
Cloud Cover(octa)	0	1	3	3	2
RH Max (%)	88	86	82	78	89
RH Min (%)	20	25	40	30	29
Wind Speed (Kmph)	18	23	24	19	16
Wind Direction (deg)	201	198	201	180	198

Agromet Advisory

General advisory: After completion of harvesting of rabi crops and before attending ploughing operation, farmers are advised to collect the soil sample from the field & send the sample to nearest soil testing laboratory for nutrient analysis. Soil testing should be done once in three years in rainfed situation and every three cropping season in irrigated situation.

Pulses: Harvest matured pulses before rainy day. Bacterial leaf blight are likely to increase in present situation in pulses due to realized rainfall; if observe, spray Streptomycin or Plantomycin @ 2 g/litre of water as foliar treatment. Rust infestation likely to occur in pulses, if observe spray Mancozeb 35.0%SC 3gram/ltr of water. Present weather is congenial for Cercospora leaf spot infestation in black gram crop. On occurrence, immediately spray Agrimycin or Streptomycin 2 g/litre, at an interval of 12-15 days, to effectively reduce pest population.

Summer Maize:

If there was infestation of Stem rot at vegetative stage as rainfall was realized in past, farmers may drench the crop root with Mancozeb 35.0%SC 3gram/ltr of water or validamycin 2ml/litre of water and there may increase the infestation of Red hairy caterpillar in maize after rainfall, to minimize the infestation farmers may spray Neem oil 5ml/ltr of water or Chlorpyrifos 20 % EC @ 2ml/litre of water.

Vegetables:

Okra: To control Yellow Vain Mosaic Virus (YVMV) in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease.

Brinjal: There are chances of infestation of Epilachna Beetle/Haddabeetle in Brinjal and cucurbits crop; to manage Epilachna Beetle in cucurbits, spray neem-based pesticide (Azadirachtin) 1500 PPM @ 3ml/litre of water at early stage of infestation; to manage Epilachna Beetle chemically, spray Chlorpyrifos 20 % EC @ 2ml/litre of water or Profenophos 50 % EC @ 2ml/litre of water.

FRUIT AND SHOOT BORER IN BRINJAL: Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@250g/acre at 15 days interval. If phomopsis blight will occur in brinjal, then spray zineb75%WP 2gram/ltr of water. Cercospora leaf spot may be found after rainfall, if observe, then spray Carbendazim or zineb75%WP 2gram/ltr of water. If helicoverpa infestation will be found, spray Flunendiamide or indoxacarb 2ml/ltr of water.

Cucurbits: To manage Red Pumpkin Beetle in Cucurbit crops spray Cyantraniliprole 10.26% OD @ 400-ml/acre or Imidacloprid 17.8% SL @ 60-ml/acre or Chlorpyrifos 20% EC @ 400-ml/acre. Late Blight may happen due to rising day temperature. There are chances of infestation of Downy Mildew in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Don't go for weeding. The grasses and weeds will help in maintaining the humidity and it is safe for creeping of the vines and does not allow the fruit to come in contact with soil and avoid fruit rot. High temperature leads to fungal wilting in pointed gourd, spray 1.5 ml Carbendazim in 1 litre water. Present weather may aggravate fruit and vine rot (symptoms: leaf spot, leaf blight, stem rot, fruit rot) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxychloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Banana: For leaf spot/sigatoka as a precautionary measure spray carbendazim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalnil @ 2.0 g / litre of water. For pseudo stem weevil Suggested to give stem injection with 5 ml Chlorpyrifos or Dimethoate or spray 2.5 ml of Chlorpyrifos per litre of water.

Mango: To control fruit drop, spray Borax @2gm per lit. of water and apply Planofix @2ml per 5 lit. of water. For control of mango fruit borer spray Neem oil 3ml + chlorpyrifos 1 ml per litre of water along with stickers at marble stage of the fruit. The spraying should be preferably carried out during evening hours. To control bacterial canker, three sprays of streptomycin (100 ppm) or Agrimycin-100 (100 ppm) after first visual symptom at 10- days intervals is recommended.

Animal husbandry: Make necessary arrangement for clean and sufficient drinking water for the animals.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA INDIA METEOROLOGICAL DEPARTMENT DISTRICT AGROMET UNIT KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-289

Date- 19/04/2022

AGROMET ADVISORY BULLETIN FOR BANKI-DAMPARA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0 mm rainfall during the last week. The mean daily maximum and minimum temperature was **38.2°C (1°C from normal)** and **25.5°C (+1.4°C from normal)**, respectively. Mean relative humidity of the district was around 66-76%.

Forecasted period: Today(8.30IST) to 24th April(8.30IST)2022

Weather Advisory: *Very light to light rainfall / Thundershower is forecasted for Thursday and Friday and Dry weather is forecasted for other 3 days in the block.* The day temperature will be **32-38°C** and night temperature will be **26-27°C** for upcoming 5 days. Maximum and minimum relative humidity will be 79-89% and 21-41% respectively. **There will be clear to partly cloudy sky for next 5 days.**

Block	Banki-Dampara (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	19/04/2022	20/04/2022	21/04/2022	22/04/2022	23/04/2022
Rainfall(mm)	0	0	6	2	0
T-Max(°C)	38	37	32	37	36
T-Min(°C)	26	26	27	27	26
Cloud Cover(octa)	0	1	4	3	1
RH Max (%)	88	86	82	79	89
RH Min (%)	21	25	41	31	30
Wind Speed (Kmph)	19	24	25	19	17
Wind Direction (deg)	201	198	201	180	198

Agromet Advisory

General advisory: After completion of harvesting of rabi crops and before attending ploughing operation, farmers are advised to collect the soil sample from the field & send the sample to nearest soil testing laboratory for nutrient analysis. Soil testing should be done once in three years in rainfed situation and every three cropping season in irrigated situation.

Pulses: Harvest matured pulses before rainy day. Bacterial leaf blight are likely to increase in present situation in pulses due to realized rainfall; if observe, spray Streptomycin or Plantomycin @ 2 g/litre of water as foliar treatment. Rust infestation likely to occur in pulses, if observe spray Mancozeb 35.0%SC 3gram/ltr of water. Present weather is congenial for **Cercospora leaf spot** infestation in black gram crop. On occurrence, immediately spray Agrimycin or Streptomycin 2 g/litre, at an interval of 12-15 days, to effectively reduce pest population.

Summer Maize:

If there was infestation of **Stem rot** at vegetative stage as rainfall was realized in past, farmers may drench the crop root with Mancozeb 35.0%SC 3gram/ltr of water or validamycin 2ml/litre of water and there may increase the infestation of **Red hairy caterpillar** in maize after rainfall, to minimize the infestation farmers may spray Neem oil 5ml/ltr of water or Chlorpyrifos 20 % EC @ 2ml/litre of water.

Vegetables:

Okra: To control **Yellow Vain Mosaic Virus (YVMV)** in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease.

Brinjal: There are chances of infestation of **Epilachna Beetle/Haddabeetle** in Brinjal and cucurbits crop; to manage Epilachna Beetle in cucurbits, spray neem-based pesticide (Azadirachtin) 1500 PPM @ 3ml/litre of water at early stage of infestation; to manage Epilachna Beetle chemically, spray Chlorpyrifos 20 % EC @ 2ml/litre of water or Profenophos 50 % EC @ 2ml/litre of water.

FRUIT AND SHOOT BORER IN BRINJAL: Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@250g/acre at 15 days interval. If **phomopsis blight** will occur in brinjal, then spray zineb75%WP 2gram/ltr of water. **Cercospora leaf spot** may be found after rainfall, if observe, then spray Carbendazim or zineb75%WP 2gram/ltr of water. If **helicoverpa** infestation will be found, spray Flunendiamide or indoxacarb 2ml/ltr of water.

Cucurbits: To manage **Red Pumpkin Beetle** in Cucurbit crops spray Cyantraniliprole 10.26% OD @ 400-ml/acre or Imidacloprid 17.8% SL @ 60-ml/acre or Chlorpyrifos 20% EC @ 400-ml/acre. **Late Blight** may happen due to rising day temperature. There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Don't go for weeding. The grasses and weeds will help in maintaining the humidity and it is safe for creeping of the vines and does not allow the fruit to come in contact with soil and avoid fruit rot. High temperature leads to fungal wilting in pointed gourd, spray 1.5 ml Carbendazim in 1 litre water. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxychloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Banana: For **leaf spot/sigatoka** as a precautionary measure spray carbendazim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalanil @ 2.0 g / litre of water. For **pseudo stem weevil** Suggested to give stem injection with 5 ml Chlorpyrifos or Dimethoate or spray 2.5 ml of Chlorpyrifos per litre of water.

Mango: To control **fruit drop**, spray Borax @2gm per lit. of water and apply Planofix @2ml per 5 lit. of water. For control of mango **fruit borer** spray Neem oil 3ml + chlorpyrifos 1 ml per litre of water along with stickers at marble stage of the fruit. The spraying should be preferably carried out during evening hours. To control **bacterial canker**, three sprays of streptomycin (100 ppm) or Agrimycin-100 (100 ppm) after first visual symptom at 10- days intervals is recommended.

Animal husbandry: Make necessary arrangement for clean and sufficient drinking water for the animals.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-289
Date- 19/04/2022

AGROMET ADVISORY BULLETIN FOR BARANGA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0 mm rainfall during the last week. The mean daily maximum and minimum temperature was **38.2°C (1°C from normal)** and **25.5°C (+1.4°C from normal)**, respectively. Mean relative humidity of the district was around 66-76%.

Forecasted period: Today(8.30IST) to 24thApril(8.30IST)2022

Weatheradvisory: Very light to light rainfall / Thundershower is forecasted for Thursday and Friday and Dry weather is forecasted for other 3days in the block. The day temperature will be **31-36°C** and night temperature will be **26-27°C** for upcoming 5 days. Maximum and minimum relative humidity will be 82-89% and 28-44% respectively. **There will be clear to partly cloudy sky for next 5days.**

Block	Baranga (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	19/04/2022	20/04/2022	21/04/2022	22/04/2022	23/04/2022
Rainfall(mm)	0	0	5	2	0
T-Max(°C)	36	34	31	35	35
T-Min(°C)	26	26	27	27	26
Cloud Cover	0	1	4	3	1
RH Max (%)	86	86	82	82	89
RH Min (%)	28	31	44	36	37
Wind Speed (Kmph)	24	28	28	24	22
Wind Direction (deg)	203	203	203	203	203

Agromet Advisory

General advisory: After completion of harvesting of rabi crops and before attending ploughing operation, farmers are advised to collect the soil sample from the field & send the sample to nearest soil testing laboratory for nutrient analysis. Soil testing should be done once in three years in rainfed situation and every three cropping season in irrigated situation.

Pulses: Harvest matured pulses before rainy day. **Bacterial leaf blight** are likely to increase in present situation in pulses due to realized rainfall; if observe, spray Streptomycin or Plantomycin @ 2 g/litre of water as foliar treatment. **Rust infestation** likely to occur in pulses, if observe spray Mancozeb 35.0%SC 3gram/ltr of water. Present weather is congenial for **Cercospora leaf spot** infestation in black gram crop. On occurrence, immediately spray Agrimycin or Streptomycin 2 g/litre, at an interval of 12-15 days, to effectively reduce pest population.

Summer Maize:

If there was infestation of **Stem rot** at vegetative stage as rainfall was realized in past, farmers may drench the crop root with Mancozeb 35.0%SC 3gram/ltr of water or validamycin 2ml/litre of water and there may increase the infestation of **Red hairy caterpillar** in maize after rainfall, to minimize the infestation farmers may spray Neem oil 5ml/ltr of water or Chlorpyrifos 20 % EC @ 2ml/litre of water.

Vegetables:

Okra: To control **Yellow Vain Mosaic Virus (YVMV)** in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease.

Brinjal: There are chances of infestation of **Epilachna Beetle/Haddabeetle** in Brinjal and cucurbits crop; to manage Epilachna Beetle in cucurbits, spray neem-based pesticide (Azadirachtin) 1500 PPM @ 3ml/litre of water at early stage of infestation; to manage Epilachna Beetle chemically, spray Chlorpyrifos 20 % EC @ 2ml/litre of water or Profenophos 50 % EC @ 2ml/litre of water.

FRUIT AND SHOOT BORER IN BRINJAL: Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@250g/acre at 15 days interval. If **phomopsis blight** will occur in brinjal, then spray zineb75%WP 2gram/litre of water. **Cercospora leaf spot** may be found after rainfall, if observe, then spray Carbendazim or zineb75%WP 2gram/litre of water. If **helicoverpa** infestation will be found, spray Flunendiamide or indoxacarb 2ml/litre of water.

Cucurbits: To manage **Red Pumpkin Beetle** in Cucurbit crops spray Cyantraniliprole 10.26% OD @ 400-ml/acre or Imidacloprid 17.8% SL @ 60-ml/acre or Chlorpyrifos 20% EC @ 400-ml/acre. **Late Blight** may happen due to rising day temperature. There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Don't go for weeding. The grasses and weeds will help in maintaining the humidity and it is safe for creeping of the vines and does not allow the fruit to come in contact with soil and avoid fruit rot. High temperature leads to fungal wilting in pointed gourd, spray 1.5 ml Carbendazim in 1 litre water. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxichloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Banana: For **leaf spot/sigatoka** as a precautionary measure spray carbendazim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalanyl @ 2.0 g/ litre of water. For **pseudo stem weevil** Suggested to give stem injection with 5 ml Chlorpyrifos or Dimethoate or spray 2.5 ml of Chlorpyrifos per litre of water.

Mango: To control **fruit drop**, spray Borax @2gm per lit. of water and apply Planofix @2ml per 5 lit. of water. For control of mango **fruit borer** spray Neem oil 3ml + chlorpyrifos 1 ml per litre of water along with stickers at marble stage of the fruit. The spraying should be preferably carried out during evening hours. To control **bacterial canker**, three sprays of streptomycin (100 ppm) or Agrimycin-100 (100 ppm) after first visual symptom at 10- days intervals is recommended.

Animal husbandry: Make necessary arrangement for clean and sufficient drinking water for the animals.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)



Bulletin No.-289
Date- 19/04/2022

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

AGROMET ADVISORY BULLETIN FOR CUTTACK SADAR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0 mm rainfall during the last week. The mean daily maximum and minimum temperature was **38.2°C (1°C from normal)** and **25.5°C (+1.4°C from normal)**, respectively. Mean relative humidity of the district was around 66-76%.

Forecasted period: Today(8.30IST) to 24th April (8.30IST)2022

Weather advisory: *Very light to light rainfall / Thundershower is forecasted for Thursday and Friday and Dry weather is forecasted for other 3 days in the block.* The day temperature will be **30-36°C** and night temperature will be **26-27°C** for upcoming 5 days. Maximum and minimum relative humidity will be 82-88% and 24-42% respectively. **There will be clear to partly cloudy sky for next 5 days.**

Block	Cuttack sadar (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	19/04/2022	20/04/2022	21/04/2022	22/04/2022	23/04/2022
Rainfall(mm)	0	0	4	2	0
T-Max(°C)	36	34	30	35	34
T-Min(°C)	26	26	27	27	27
Cloud Cover(octa)	1	1	3	3	1
RH Max (%)	88	86	82	82	87
RH Min (%)	24	27	42	33	32
Wind Speed (Kmph)	22	26	27	22	20
Wind Direction (deg)	201	203	203	203	201

Agromet Advisory

General advisory: After completion of harvesting of rabi crops and before attending ploughing operation, farmers are advised to collect the soil sample from the field & send the sample to nearest soil testing laboratory for nutrient analysis. Soil testing should be done once in three years in rainfed situation and every three cropping season in irrigated situation.

Pulses: Harvest matured pulses before rainy day. Bacterial leaf blight are likely to increase in present situation in pulses due to realized rainfall; if observe, spray Streptomycin or Plantomycin @ 2 g/litre of water as foliar treatment. Rust infestation likely to occur in pulses, if observe spray Mancozeb 35.0%SC 3gram/ltr of water. Present weather is congenial for **Cercospora leaf spot** infestation in black gram crop. On occurrence, immediately spray Agrimycin or Streptomycin 2 g/litre, at an interval of 12-15 days, to effectively reduce pest population.

Summer Maize:

If there was infestation of **Stem rot** at vegetative stage as rainfall was realized in past, farmers may drench the crop root with Mancozeb 35.0%SC 3gram/ltr of water or validamycin 2ml/litre of water and there may increase the infestation of **Red hairy caterpillar** in maize after rainfall, to minimize the infestation farmers may spray Neem oil 5ml/ltr of water or Chlorpyrifos 20 % EC @ 2ml/litre of water.

Vegetables:

Okra: To control **Yellow Vain Mosaic Virus (YVMV)** in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease.

Brinjal: There are chances of infestation of **Epilachna Beetle/Haddabeetle** in Brinjal and cucurbits crop; to manage Epilachna Beetle in cucurbits, spray neem-based pesticide (Azadirachtin) 1500 PPM @ 3ml/litre of water at early stage of infestation; to manage Epilachna Beetle chemically, spray Chlorpyrifos 20 % EC @ 2ml/litre of water or Profenophos 50 % EC @ 2ml/litre of water.

FRUIT AND SHOOT BORER IN BRINJAL: Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@250g/acre at 15 days interval. If **phomopsis blight** will occur in brinjal, then spray zineb 75%WP 2gram/ltr of water. **Cercospora leaf spot** may be found after rainfall, if observe, then spray Carbendazim or zineb 75%WP 2gram/ltr of water. If **helicoverpa** infestation will be found, spray Flunendiamide or indoxacarb 2ml/ltr of water.

Cucurbits: To manage **Red Pumpkin Beetle** in Cucurbit crops spray Cyantraniliprole 10.26% OD @ 400-ml/acre or Imidacloprid 17.8% SL @ 60-ml/acre or Chlorpyrifos 20% EC @ 400-ml/acre. **Late Blight** may happen due to rising day temperature. There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Don't go for weeding. The grasses and weeds will help in maintaining the humidity and it is safe for creeping of the vines and does not allow the fruit to come in contact with soil and avoid fruit rot. High temperature leads to fungal wilting in pointed gourd, spray 1.5 ml Carbendazim in 1 litre water. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxychloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Banana: For **leaf spot/sigatoka** as a precautionary measure spray carbendazim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalnil @ 2.0 g/ litre of water. For **pseudo stem weevil** Suggested to give stem injection with 5 ml Chlorpyrifos or Dimethoate or spray 2.5 ml of Chlorpyrifos per litre of water.

Mango: To control **fruit drop**, spray Borax @2gm per lit. of water and apply Planofix @2ml per 5 lit. of water. For control of mango **fruit borer** spray Neem oil 3ml + chlorpyrifos 1 ml per litre of water along with stickers at marble stage of the fruit. The spraying should be preferably carried out during evening hours. To control **bacterial canker**, three sprays of streptomycin (100 ppm) or Agrimycin-100 (100 ppm) after first visual symptom at 10- days intervals is recommended.

Animal husbandry: Make necessary arrangement for clean and sufficient drinking water for the animals.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA INDIA METEOROLOGICAL DEPARTMENT DISTRICT AGROMET UNIT KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

AGROMET ADVISORY BULLETIN FOR KANTAPADA



Bulletin No.-289

Date- 19/04/2022

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0 mm rainfall during the last week. The mean daily maximum and minimum temperature was **38.2°C (1°C from normal)** and **25.5°C (+1.4°C from normal)**, respectively. Mean relative humidity of the district was around 66-76%.

Forecasted period: Today(8.30IST) to 24th April (8.30IST)2022

Weather advisory: *Very light to light rainfall / Thundershower is forecasted for Thursday and Friday and Dry weather is forecasted for other 3days in the block.* The day temperature will be **32-35°C** and night temperature will be **26°C** for upcoming 5 days. Maximum and minimum relative humidity will be 81-89% and 39-48% respectively. **There will be clear to partly cloudy sky for next 5days.**

Block	Kantapada (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	19/04/2022	20/04/2022	21/04/2022	22/04/2022	23/04/2022
Rainfall(mm)	0	0	3	2	0
T-Max(°C)	34	34	32	35	34
T-Min(°C)	26	26	26	26	26
Cloud Cover(octa)	0	1	4	2	1
RH Max (%)	85	85	81	83	89
RH Min (%)	39	39	48	41	45
Wind Speed (Kmph)	25	28	28	27	24
Wind Direction (deg)	203	203	203	203	203

Agromet Advisory

General advisory: After completion of harvesting of rabi crops and before attending ploughing operation, farmers are advised to collect the soil sample from the field & send the sample to nearest soil testing laboratory for nutrient analysis. Soil testing should be done once in three years in rainfed situation and every three cropping season in irrigated situation.

Pulses: Harvest matured pulses before rainy day. Bacterial leaf blight are likely to increase in present situation in pulses due to realized rainfall; if observe, spray Streptomycin or Plantomycin @ 2 g/litre of water as foliar treatment. Rust infestation likely to occur in pulses, if observe spray Mancozeb 35.0%SC 3gram/ltr of water. Present weather is congenial for **Cercospora leaf spot** infestation in black gram crop. On occurrence, immediately spray Agrimycin or Streptomycin 2 g/litre, at an interval of 12-15 days, to effectively reduce pest population.

Summer Maize:

If there was infestation of **Stem rot** at vegetative stage as rainfall was realized in past, farmers may drench the crop root with Mancozeb 35.0%SC 3gram/ltr of water or validamycin 2ml/litre of water and there may increase the infestation of **Red hairy caterpillar** in maize after rainfall, to minimize the infestation farmers may spray Neem oil 5ml/ltr of water or Chlorpyrifos 20 % EC @ 2ml/litre of water.

Vegetables:

Okra: To control **Yellow Vain Mosaic Virus (YVMV)** in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease.

Brinjal: There are chances of infestation of **Epilachna Beetle/Haddabeetle** in Brinjal and cucurbits crop; to manage Epilachna Beetle in cucurbits, spray neem-based pesticide (Azadirachtin) 1500 PPM @ 3ml/litre of water at early stage of infestation; to manage Epilachna Beetle chemically, spray Chlorpyrifos 20 % EC @ 2ml/litre of water or Profenophos 50 % EC @ 2ml/litre of water.

FRUIT AND SHOOT BORER IN BRINJAL: Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@250g/acre at 15 days interval. If **phomopsis blight** will occur in brinjal, then spray zineb75%WP 2gram/ltr of water. **Cercospora leaf spot** may be found after rainfall, if observe, then spray Carbendazim or zineb75%WP 2gram/ltr of water. If **helicoverpa infestation** will be found, spray Flunendiamide or indoxacarb 2ml/ltr of water.

Cucurbits: To manage **Red Pumpkin Beetle** in Cucurbit crops spray Cyantraniliprole 10.26% OD @ 400-ml/acre or Imidacloprid 17.8% SL @ 60-ml/acre or Chlorpyrifos 20% EC @ 400-ml/acre. **Late Blight** may happen due to rising day temperature. There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Don't go for weeding. The grasses and weeds will help in maintaining the humidity and it is safe for creeping of the vines and does not allow the fruit to come in contact with soil and avoid fruit rot. High temperature leads to fungal wilting in pointed gourd, spray 1.5 ml Carbendazim in 1 litre water. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxychloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Banana: For **leaf spot/sigatoka** as a precautionary measure spray carbendazim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalanil @ 2.0 g / litre of water. For **pseudo stem weevil** Suggested to give stem injection with 5 ml Chlorpyrifos or Dimethoate or spray 2.5 ml of Chlorpyrifos per litre of water.

Mango: To control **fruit drop**, spray Borax @2gm per lit. of water and apply Planofix @2ml per 5 lit. of water. For control of mango **fruit borer** spray Neem oil 3ml + chlorpyrifos 1 ml per litre of water along with stickers at marble stage of the fruit. The spraying should be preferably carried out during evening hours. To control **bacterial canker**, three sprays of streptomycin (100 ppm) or Agrimycin-100 (100 ppm) after first visual symptom at 10- days intervals is recommended.

Animal husbandry: Make necessary arrangement for clean and sufficient drinking water for the animals.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

AGROMET ADVISORY BULLETIN FOR MAHANGA



Bulletin No.-289

Date- 19/04/2022

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0 mm rainfall during the last week. The mean daily maximum and minimum temperature was **38.2°C (1°C from normal)** and **25.5°C (+1.4°C from normal)**, respectively. Mean relative humidity of the district was around 66-76%.

Forecasted period: Today(8.30IST) to 24th April (8.30IST)2022

Weather advisory: *Very light to light rainfall / Thundershower is forecasted for Thursday and Friday and Dry weather is forecasted for other 3days in the block.* The day temperature will be and it will be **31-36°C** and night temperature will be **25-26°C** for upcoming 5 days. Maximum and minimum relative humidity will be 81-86% and 26-43% respectively. **There will be clear to partly cloudy sky for next 5days.**

Block	Mahanga (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	19/04/2022	20/04/2022	21/04/2022	22/04/2022	23/04/2022
Rainfall(mm)	0	0	3	2	0
T-Max(°C)	36	34	31	35	36
T-Min(°C)	25	26	26	26	26
Cloud Cover(octal)	0	0	3	2	1
RH Max (%)	86	85	81	82	86
RH Min (%)	26	32	43	36	37
Wind Speed (Kmph)	24	27	27	23	22
Wind Direction (deg)	201	203	203	203	203

Agromet Advisory

General advisory: After completion of harvesting of rabi crops and before attending ploughing operation, farmers are advised to collect the soil sample from the field & send the sample to nearest soil testing laboratory for nutrient analysis. **Soil testing should be done once in three years in rainfed situation and every three cropping season in irrigated situation.**

Pulses: Harvest matured pulses before rainy day. **Bacterial leaf blight** are likely to increase in present situation in pulses due to realized rainfall; if observe, spray Streptomycin or Plantomycin @ 2 g/litre of water as foliar treatment. **Rust infestation** likely to occur in pulses, if observe spray Mancozeb 35.0%SC 3gram/ltr of water. Present weather is congenial for **Cercospora leaf spot** infestation in black gram crop. On occurrence, immediately spray Agrimycin or Streptomycin 2 g/litre, at an interval of 12-15 days, to effectively reduce pest population.

Summer Maize:

If there was infestation of **Stem rot** at vegetative stage as rainfall was realized in past, farmers may drench the crop root with Mancozeb 35.0%SC 3gram/ltr of water or validamycin 2ml/litre of water and there may increase the infestation of **Red hairy caterpillar** in maize after rainfall, to minimize the infestation farmers may spray Neem oil 5ml/ltr of water or Chlorpyrifos 20 % EC @ 2ml/litre of water.

Vegetables:

Okra: To control **Yellow Vain Mosaic Virus (YVMV)** in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease.

Brinjal: There are chances of infestation of **Epilachna Beetle/Haddabeetle** in Brinjal and cucurbits crop; to manage Epilachna Beetle in cucurbits, spray neem-based pesticide (Azadirachtin) 1500 PPM @ 3ml/litre of water at early stage of infestation; to manage Epilachna Beetle chemically, spray Chlorpyrifos 20 % EC @ 2ml/litre of water or Profenophos 50 % EC @ 2ml/litre of water.

FRUIT AND SHOOT BORER IN BRINJAL: Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@250g/acre at 15 days interval. If **phomopsis blight** will occur in brinjal, then spray zineb75%WP 2gram/ltr of water. **Cercospora leaf spot** may be found after rainfall, if observe, then spray Carbendazim or zineb75%WP 2gram/ltr of water. If **helioverpa infestation** will be found, spray Flunendiamide or indoxacarb 2ml/ltr of water.

Cucurbits: To manage **Red Pumpkin Beetle** in Cucurbit crops spray Cyantraniliprole 10.26% OD @ 400-ml/acre or Imidacloprid 17.8% SL @ 60-ml/acre or Chlorpyrifos 20% EC @ 400-ml/acre. **Late Blight** may happen due to rising day temperature. There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Don't go for weeding. The grasses and weeds will help in maintaining the humidity and it is safe for creeping of the vines and does not allow the fruit to come in contact with soil and avoid fruit rot. High temperature leads to fungal wilting in pointed gourd, spray 1.5 ml Carbendazim in 1 litre water. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxychloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Banana: For **leaf spot/sigatoka** as a precautionary measure spray carbendazim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalnil @ 2.0 g / litre of water. For **pseudo stem weevil** Suggested to give stem injection with 5 ml Chlorpyrifos or Dimethoate or spray 2.5 ml of Chlorpyrifos per litre of water.

Mango: To control **fruit drop**, spray Borax @2gm per lit. of water and apply Planofix @2ml per 5 lit. of water. For control of mango **fruit borer** spray Neem oil 3ml + chlorpyrifos 1 ml per litre of water along with stickers at marble stage of the fruit. The spraying should be preferably carried out during evening hours. To control **bacterial canker**, three sprays of streptomycin (100 ppm) or Agrimycin-100 (100 ppm) after first visual symptom at 10- days intervals is recommended.

Dr Sujata Sathy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA INDIA METEOROLOGICAL DEPARTMENT DISTRICT AGROMET UNIT KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-289

Date- 19/04/2022

AGROMET ADVISORY BULLETIN FOR NARASINGHPUR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0 mm rainfall during the last week. The mean daily maximum and minimum temperature was **38.2°C (1°C from normal)** and **25.5°C (+1.4°C from normal)**, respectively. Mean relative humidity of the district was around 66-76%.

Forecasted period: Today(8.30IST) to 24th April (8.30IST)2022

Weather advisory: *Very light to light rainfall / Thundershower is forecasted for Thursday and Friday and Dry weather is forecasted for other 3 days in the block.* The day temperature will be **36-39°C** and night temperature will be **24-27°C** for upcoming 5 days. Maximum and minimum relative humidity will be 73-86% and 10-26% respectively. **There will be clear to partly cloudy sky for next 5 days.**

Block	Narasinghpur (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	19/04/2022	20/04/2022	21/04/2022	22/04/2022	23/04/2022
Rainfall(mm)	0	0	6	2	0
T-Max(°C)	39	39	36	37	39
T-Min(°C)	24	25	26	27	27
Cloud Cover(octa)	1	2	2	2	3
RH Max (%)	86	79	79	73	80
RH Min (%)	10	14	26	20	17
Wind Speed (Kmph)	9	11	14	10	8
Wind Direction (deg)	189	189	189	169	180

Agromet Advisory

General advisory: After completion of harvesting of rabi crops and before attending ploughing operation, farmers are advised to collect the soil sample from the field & send the sample to nearest soil testing laboratory for nutrient analysis. Soil testing should be done once in three years in rainfed situation and every three cropping season in irrigated situation.

Pulses: Harvest matured pulses before rainy day. Bacterial leaf blight are likely to increase in present situation in pulses due to realized rainfall; if observe, spray Streptomycin or Plantomycin @ 2 g/litre of water as foliar treatment. Rust infestation likely to occur in pulses, if observe spray Mancozeb 35.0%SC 3gram/ltr of water. Present weather is congenial for **Cercospora leaf spot** infestation in black gram crop. On occurrence, immediately spray Agrimycin or Streptomycin 2 g/litre, at an interval of 12-15 days, to effectively reduce pest population.

Summer Maize:

If there was infestation of **Stem rot** at vegetative stage as rainfall was realized in past, farmers may drench the crop root with Mancozeb 35.0%SC 3gram/ltr of water or validamycin 2ml/litre of water and there may increase the infestation of **Red hairy caterpillar** in maize after rainfall, to minimize the infestation farmers may spray Neem oil 5ml/ltr of water or Chlorpyrifos 20 % EC @ 2ml/litre of water.

Vegetables:

Okra: To control **Yellow Vain Mosaic Virus (YVMV)** in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease.

Brinjal: There are chances of infestation of **Epilachna Beetle/Haddabeetle** in Brinjal and cucurbits crop; to manage Epilachna Beetle in cucurbits, spray neem-based pesticide (Azadirachtin) 1500 PPM @ 3ml/litre of water at early stage of infestation; to manage Epilachna Beetle chemically, spray Chlorpyrifos 20 % EC @ 2ml/litre of water or Profenophos 50 % EC @ 2ml/litre of water.

FRUIT AND SHOOT BORER IN BRINJAL: Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@250g/acre at 15 days interval. If **phomopsis blight** will occur in brinjal, then spray zineb75%WP 2gram/ltr of water. **Cercospora leaf spot** may be found after rainfall, if observe, then spray Carbendazim or zineb75%WP 2gram/ltr of water. If **helicopter infestation** will be found, spray Flunendiamide or indoxacarb 2ml/ltr of water.

Cucurbits: To manage **Red Pumpkin Beetle** in Cucurbit crops spray Cyantraniliprole 10.26% OD @ 400-ml/acre or Imidacloprid 17.8% SL @ 60-ml/acre or Chlorpyrifos 20% EC @ 400-ml/acre. **Late Blight** may happen due to rising day temperature. There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Don't go for weeding. The grasses and weeds will help in maintaining the humidity and it is safe for creeping of the vines and does not allow the fruit to come in contact with soil and avoid fruit rot. High temperature leads to fungal wilting in pointed gourd, spray 1.5 ml Carbendazim in 1 litre water. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxychloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Banana: For **leaf spot/sigatoka** as a precautionary measure spray carbendazim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalanyl @ 2.0 g / litre of water. For **pseudo stem weevil** Suggested to give stem injection with 5 ml Chlorpyrifos or Dimethoate or spray 2.5 ml of Chlorpyrifos per litre of water.

Mango: To control **fruit drop**, spray Borax @2gm per lit. of water and apply Planofix @2ml per 5 lit. of water. For control of mango **fruit borer** spray Neem oil 3ml + chlorpyrifos 1 ml per litre of water along with stickers at marble stage of the fruit. The spraying should be preferably carried out during evening hours. To control **bacterial canker**, three sprays of streptomycin (100 ppm) or Agrimycin-100 (100 ppm) after first visual symptom at 10- days intervals is recommended.

Animal husbandry: Make necessary arrangement for clean and sufficient drinking water for the animals.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-289
Date- 19/04/2022

AGROMET ADVISORY BULLETIN FOR NIALI

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0 mm rainfall during the last week. The mean daily maximum and minimum temperature was **38.2°C (1°C from normal)** and **25.5°C (+1.4°C from normal)**, respectively. Mean relative humidity of the district was around 66-76%.

Forecasted period: Today(8.30IST) to 24th April (8.30IST)2022

Weather advisory: *Very light to light rainfall / Thundershower is forecasted for Thursday and Friday and Dry weather is forecasted for other 3 days in the block.* The day temperature will be **33-35°C** and night temperature will be **26°C** for upcoming 5 days. Maximum and minimum relative humidity will be 81-88% and 45-52% respectively. **There will be clear to partly cloudy sky for next 5 days.**

Block	Niali (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	19/04/2022	20/04/2022	21/04/2022	22/04/2022	23/04/2022
Rainfall(mm)	0	0	3	2	0
T-Max(°C)	34	34	33	35	34
T-Min(°C)	26	26	26	26	26
Cloud Cover(octa)	0	1	4	1	0
RH Max (%)	86	85	81	83	88
RH Min (%)	48	45	52	46	52
Wind Speed(Kmph)	26	28	29	28	26
Wind Direction(deg)	203	203	203	203	203

Agromet Advisory

General advisory: After completion of harvesting of rabi crops and before attending ploughing operation, farmers are advised to collect the soil sample from the field & send the sample to nearest soil testing laboratory for nutrient analysis. Soil testing should be done once in three years in rainfed situation and every three cropping season in irrigated situation.

Pulses: Harvest matured pulses before rainy day. Bacterial leaf blight are likely to increase in present situation in pulses due to realized rainfall; if observe, spray Streptomycin or Plantomycin @ 2 g/litre of water as foliar treatment. Rust infestation likely to occur in pulses, if observe spray Mancozeb 35.0%SC 3gram/ltr of water. Present weather is congenial for Cercospora leaf spot infestation in black gram crop. On occurrence, immediately spray Agrimycin or Streptomycin 2 g/litre, at an interval of 12-15 days, to effectively reduce pest population.

Summer Maize:

If there was infestation of Stem rot at vegetative stage as rainfall was realized in past, farmers may drench the crop root with Mancozeb 35.0%SC 3gram/ltr of water or validamycin 2ml/ltr of water and there may increase the infestation of Red hairy caterpillar in maize after rainfall, to minimize the infestation farmers may spray Neem oil 5ml/ltr of water or Chlorpyrifos 20 % EC @ 2ml/litre of water.

Vegetables:

Okra: To control Yellow Vain Mosaic Virus (YVMV) in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease.

Brinjal: There are chances of infestation of Epilachna Beetle/Haddabeetle in Brinjal and cucurbits crop; to manage Epilachna Beetle in cucurbits, spray neem-based pesticide (Azadirachtin) 1500 PPM @ 3ml/litre of water at early stage of infestation; to manage Epilachna Beetle chemically, spray Chlorpyrifos 20 % EC @ 2ml/litre of water or Profenophos 50 % EC @ 2ml/litre of water.

FRUIT AND SHOOT BORER IN BRINJAL: Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@250g/acre at 15 days interval. If phomopsis blight will occur in brinjal, then spray zineb75%WP 2gram/ltr of water. Cercospora leaf spot may be found after rainfall, if observe, then spray Carbendazim or zineb75%WP 2gram/ltr of water. If helicoverpa infestation will be found, spray Flunendiamide or indoxacarb 2ml/ltr of water.

Cucurbits: To manage Red Pumpkin Beetle in Cucurbit crops spray Cyantraniliprole 10.26% OD @ 400-ml/acre or Imidacloprid 17.8% SL @ 60-ml/acre or Chlorpyrifos 20% EC @ 400-ml/acre. Late Blight may happen due to rising day temperature. There are chances of infestation of Downy Mildew in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Don't go for weeding. The grasses and weeds will help in maintaining the humidity and it is safe for creeping of the vines and does not allow the fruit to come in contact with soil and avoid fruit rot. High temperature leads to fungal wilting in pointed gourd, spray 1.5 ml Carbendazim in 1 litre water. Present weather may aggravate fruit and vine rot (symptoms: leaf spot, leaf blight, stem rot, fruit rot) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxychloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Banana: For leaf spot/sigatoka as a precautionary measure spray carbendazim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalnil@ 2.0 g/ litre of water. For pseudo stem weevil Suggested to give stem injection with 5 ml Chloropyrifos or Dimethoate or spray 2.5 ml of Chloropyrifos per litre of water.

Mango: To control fruit drop, spray Borax @2gm per lit. of water and apply Planofix @2ml per 5 lit. of water. For control of mango fruit borer spray Neem oil 3ml + chlorpyrifos 1 ml per litre of water along with stickers at marble stage of the fruit. The spraying should be preferably carried out during evening hours. To control bacterial canker, three sprays of streptomycin (100 ppm) or Agrimycin-100 (100 ppm) after first visual symptom at 10- days intervals is recommended.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-289

Date- 19/04/2022

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0 mm rainfall during the last week. The mean daily maximum and minimum temperature was **38.2°C (1°C from normal)** and **25.5°C (+1.4°C from normal)**, respectively. Mean relative humidity of the district was around 66-76%.

Forecasted period: Today(8.30IST) to 24thApril (8.30IST)2022

Weather advisory: *Very light to light rainfall / Thundershower is forecasted for Thursday and Friday and Dry weather is forecasted for other 3days in the block.* The day temperature will be **31-35°C** and night temperature will be **26°C** for upcoming 5 days. Maximum and minimum relative humidity will be **81-87%** and **30-44%** respectively. **There will be clear to partly cloudy sky for next 5days.**

Block	Nischintakoili (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	19/04/2022	20/04/2022	21/04/2022	22/04/2022	23/04/2022
Rainfall(mm)	0	0	3	2	0
T-Max(°C)	35	34	31	35	35
T-Min(°C)	26	26	26	26	26
Cloud Cover(octa)	0	1	3	2	1
RH Max (%)	86	85	81	82	87
RH Min (%)	30	34	44	38	41
Wind Speed (Kmph)	25	27	27	24	23
Wind Direction (deg)	201	203	203	203	203

Agromet Advisory

General advisory: After completion of harvesting of rabi crops and before attending ploughing operation, farmers are advised to collect the soil sample from the field & send the sample to nearest soil testing laboratory for nutrient analysis. Soil testing should be done once in three years in rainfed situation and every three cropping season in irrigated situation.

Pulses: Harvest matured pulses before rainy day. Bacterial leaf blight are likely to increase in present situation in pulses due to realized rainfall; if observe, spray Streptomycin or Plantomycin @ 2 g/litre of water as foliar treatment. Rust infestation likely to occur in pulses, if observe spray Mancozeb 35.0%SC 3gram/ltr of water. Present weather is congenial for Cercospora leaf spot infestation in black gram crop. On occurrence, immediately spray Agrimycin or Streptomycin 2 g/litre, at an interval of 12-15 days, to effectively reduce pest population.

Summer Maize:

If there was infestation of Stem rot at vegetative stage as rainfall was realized in past, farmers may drench the crop root with Mancozeb 35.0%SC 3gram/ltr of water or validamycin 2ml/litre of water and there may increase the infestation of Red hairy caterpillar in maize after rainfall, to minimize the infestation farmers may spray Neem oil 5ml/ltr of water or Chlorpyrifos 20 % EC @ 2ml/litre of water.

Vegetables:

Okra: To control Yellow Vain Mosaic Virus (YVMV) in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease.

Brinjal: There are chances of infestation of Epilachna Beetle/Haddabeetle in Brinjal and cucurbits crop; to manage Epilachna Beetle in cucurbits, spray neem-based pesticide (Azadirachtin) 1500 PPM @ 3ml/litre of water at early stage of infestation; to manage Epilachna Beetle chemically, spray Chlorpyrifos 20 % EC @ 2ml/litre of water or Profenophos 50 % EC @ 2ml/litre of water.

FRUIT AND SHOOT BORER IN BRINJAL: Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@250g/acre at 15 days interval. If phomopsis blight will occur in brinjal, then spray zineb75%WP 2gram/ltr of water. Cercospora leaf spot may be found after rainfall, if observe, then spray Carbendazim or zineb75%WP 2gram/ltr of water. If helicoverpa infestation will be found, spray Flunendiamide or indoxacarb 2ml/ltr of water.

Cucurbits: To manage Red Pumpkin Beetle in Cucurbit crops spray Cyantraniliprole 10.26% OD @ 400-ml/acre or Imidacloprid 17.8% SL @ 60-ml/acre or Chlorpyrifos 20% EC @ 400-ml/acre. Late Blight may happen due to rising day temperature. There are chances of infestation of Downy Mildew in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Don't go for weeding. The grasses and weeds will help in maintaining the humidity and it is safe for creeping of the vines and does not allow the fruit to come in contact with soil and avoid fruit rot. High temperature leads to fungal wilting in pointed gourd, spray 1.5 ml Carbendazim in 1 litre water. Present weather may aggravate fruit and vine rot (symptoms: leaf spot, leaf blight, stem rot, fruit rot) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxychloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Banana: For leaf spot/sigatoka as a precautionary measure spray carbendazim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalnil @ 2.0 g / litre of water. For pseudo stem weevil Suggested to give stem injection with 5 ml Chlorpyrifos or Dimethoate or spray 2.5 ml of Chlorpyrifos per litre of water.

Mango: To control fruit drop, spray Borax @2gm per lit. of water and apply Planofix @2ml per 5 lit. of water. For control of mango fruit borer spray Neem oil 3ml + chlorpyrifos 1 ml per litre of water along with stickers at marble stage of the fruit. The spraying should be preferably carried out during evening hours. To control bacterial canker, three sprays of streptomycin (100 ppm) or Agrimycin-100 (100 ppm) after first visual symptom at 10- days intervals is recommended.

Animal husbandry: Make necessary arrangement for clean and sufficient drinking water for the animals.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-289
Date- 19/04/2022

AGROMET ADVISORY BULLETIN FOR SALEPUR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0 mm rainfall during the last week. The mean daily maximum and minimum temperature was **38.2°C (1°C from normal)** and **25.5°C (+1.4°C from normal)**, respectively. Mean relative humidity of the district was around 66-76%.

Forecasted period: Today(8.30IST) to 24th April (8.30IST)2022

Weather advisory: *Very light to light rainfall / Thundershower is forecasted for Thursday and Friday and Dry weather is forecasted for other 3days in the block.* The day temperature will be **31-35°C** and night temperature will be **26°C** for upcoming 5 days. Maximum and minimum relative humidity will be 81-87% and 28-44% respectively. **There will be clear to partly cloudy sky for next 5days.**

Block	Salepur (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	19/04/2022	20/04/2022	21/04/2022	22/04/2022	23/04/2022
Rainfall(mm)	0	0	3	2	0
T-Max(°C)	35	34	31	34	35
T-Min(°C)	26	26	26	26	26
Cloud Cover(octa)	0	1	3	2	1
RH Max (%)	86	85	81	82	87
RH Min (%)	28	33	44	37	39
Wind Speed (Kmph)	25	27	27	24	22
Wind Direction (deg)	201	203	203	203	203

Agromet Advisory

General advisory: After completion of harvesting of rabi crops and before attending ploughing operation, farmers are advised to collect the soil sample from the field & send the sample to nearest soil testing laboratory for nutrient analysis. Soil testing should be done once in three years in rainfed situation and every three cropping season in irrigated situation.

Pulses: Harvest matured pulses before rainy day. Bacterial leaf blight are likely to increase in present situation in pulses due to realized rainfall; if observe, spray Streptomycin or Plantomycin @ 2 g/litre of water as foliar treatment. Rust infestation likely to occur in pulses, if observe spray Mancozeb 35.0%SC 3gram/ltr of water. Present weather is congenial for Cercospora leaf spot infestation in black gram crop. On occurrence, immediately spray Agrimycin or Streptomycin 2 g/litre, at an interval of 12-15 days, to effectively reduce pest population.

Summer Maize:

If there was infestation of Stem rot at vegetative stage as rainfall was realized in past, farmers may drench the crop root with Mancozeb 35.0%SC 3gram/ltr of water or validamycin 2ml/litre of water and there may increase the infestation of Red hairy caterpillar in maize after rainfall, to minimize the infestation farmers may spray Neem oil 5ml/ltr of water or Chlorpyrifos 20 % EC @ 2ml/litre of water.

Vegetables:

Okra: To control Yellow Vain Mosaic Virus (YVMV) in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease.

Brinjal: There are chances of infestation of Epilachna Beetle/Haddabeetle in Brinjal and cucurbits crop; to manage Epilachna Beetle in cucurbits, spray neem-based pesticide (Azadirachtin) 1500 PPM @ 3ml/litre of water at early stage of infestation; to manage Epilachna Beetle chemically, spray Chlorpyrifos 20 % EC @ 2ml/litre of water or Profenophos 50 % EC @ 2ml/litre of water.

FRUIT AND SHOOT BORER IN BRINJAL: Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@250g/acre at 15 days interval. If phomopsis blight will occur in brinjal, then spray zineb 75%WP 2gram/ltr of water. Cercospora leaf spot may be found after rainfall, if observe, then spray Carbendazim or zineb 75%WP 2gram/ltr of water. If helicoverpa infestation will be found, spray Flunendiamide or indoxacarb 2ml/ltr of water.

Cucurbits: To manage Red Pumpkin Beetle in Cucurbit crops spray Cyantraniliprole 10.26% OD @ 400-ml/acre or Imidacloprid 17.8% SL @ 60-ml/acre or Chlorpyrifos 20% EC @ 400-ml/acre. Late Blight may happen due to rising day temperature. There are chances of infestation of Downy Mildew in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Don't go for weeding. The grasses and weeds will help in maintaining the humidity and it is safe for creeping of the vines and does not allow the fruit to come in contact with soil and avoid fruit rot. High temperature leads to fungal wilting in pointed gourd, spray 1.5 ml Carbendazim in 1 litre water. Present weather may aggravate fruit and vine rot (symptoms: leaf spot, leaf blight, stem rot, fruit rot) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxychloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Banana: For leaf spot/sigatoka as a precautionary measure spray carbendazim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalanil @ 2.0 g / litre of water. For pseudo stem weevil Suggested to give stem injection with 5 ml Chlorpyrifos or Dimethoate or spray 2.5 ml of Chlorpyrifos per litre of water.

Mango: To control fruit drop, spray Borax @ 2gm per lit. of water and apply Planofix @ 2ml per 5 lit. of water. For control of mango fruit borer spray Neem oil 3ml + chlorpyrifos 1 ml per litre of water along with stickers at marble stage of the fruit. The spraying should be preferably carried out during evening hours. To control bacterial canker, three sprays of streptomycin (100 ppm) or Agrimycin-100 (100 ppm) after first visual symptom at 10- days intervals is recommended.

Animal husbandry: Make necessary arrangement for clean and sufficient drinking water for the animals.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA INDIA METEOROLOGICAL DEPARTMENT DISTRICT AGROMET UNIT KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

AGROMET ADVISORY BULLETIN FOR TANGI-CHOUDWAR



Bulletin No.-289

Date- 19/04/2022

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0 mm rainfall during the last week. The mean daily maximum and minimum temperature was **38.2°C (1°C from normal)** and **25.5°C (+1.4°C from normal)**, respectively. Mean relative humidity of the district was around 66-76%.

Forecasted period: Today(8.30IST) to 24th April (8.30IST)2022

Weatheradvisory: *Very light to light rainfall / Thundershower is forecasted for Thursday and Friday and Dry weather is forecasted for other 3days in the block.* The day temperature will be **34-35°C** and night temperature will be **27-28°C** for upcoming 5 days. Maximum and minimum relative humidity will be **82-88%** and **52-59%** respectively. **There will be clear to partly cloudy sky for next 5days.**

Block	Tangi-Choudwar(VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	19/04/2022	20/04/2022	21/04/2022	22/04/2022	23/04/2022
Rainfall(mm)	0	0	6	2	0
T-Max(°C)	35	35	34	35	34
T-Min(°C)	27	28	28	28	28
Cloud Cover(octa)	0	1	4	1	0
RH Max (%)	88	86	82	84	86
RH Min (%)	59	52	59	53	59
Wind Speed (Kmph)	27	29	29	28	28
Wind Direction (deg)	203	203	203	203	203

Agromet Advisory

General advisory: After completion of harvesting of rabi crops and before attending ploughing operation, farmers are advised to collect the soil sample from the field & send the sample to nearest soil testing laboratory for nutrient analysis. Soil testing should be done once in three years in rainfed situation and every three cropping season in irrigated situation.

Pulses: Harvest matured pulses before rainy day. Bacterial leaf blight are likely to increase in present situation in pulses due to realized rainfall; if observe, spray Streptomycin or Plantomycin @ 2 g/litre of water as foliar treatment. Rust infestation likely to occur in pulses, if observe spray Mancozeb 35.0%SC 3gram/ltr of water. Present weather is congenial for **Cercospora leaf spot** infestation in black gram crop. On occurrence, immediately spray Agrimycin or Streptomycin 2 g/litre, at an interval of 12-15 days, to effectively reduce pest population.

Summer Maize:

If there was infestation of **Stem rot** at vegetative stage as rainfall was realized in past, farmers may drench the crop root with Mancozeb 35.0%SC 3gram/ltr of water or validamycin 2ml/litre of water and there may increase the infestation of **Red hairy caterpillar** in maize after rainfall, to minimize the infestation farmers may spray Neem oil 5ml/ltr of water or Chlorpyrifos 20 % EC @ 2ml/litre of water.

Vegetables:

Okra: To control **Yellow Vain Mosaic Virus (YVMV)** in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease.

Brinjal: There are chances of infestation of **Epilachna Beetle/Haddabeetle** in Brinjal and cucurbits crop; to manage Epilachna Beetle in cucurbits, spray neem-based pesticide (Azadirachtin) 1500 PPM @ 3ml/litre of water at early stage of infestation; to manage Epilachna Beetle chemically, spray Chlorpyrifos 20 % EC @ 2ml/litre of water or Profenophos 50 % EC @ 2ml/litre of water.

FRUIT AND SHOOT BORER IN BRINJAL: Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@250g/acre at 15 days interval. If **phomopsis blight** will occur in brinjal, then spray zineb75%WP 2gram/ltr of water. **Cercospora leaf spot** may be found after rainfall, if observe, then spray Carbendazim or zineb75%WP 2gram/ltr of water. If **helicoverpa infestation** will be found, spray Flunendiamide or indoxacarb 2ml/ltr of water.

Cucurbits: To manage **Red Pumpkin Beetle** in Cucurbit crops spray Cyantraniliprole 10.26% OD @ 400-ml/acre or Imidacloprid 17.8% SL @ 60-ml/acre or Chlorpyrifos 20% EC @ 400-ml/acre. **Late Blight** may happen due to rising day temperature. There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Don't go for weeding. The grasses and weeds will help in maintaining the humidity and it is safe for creeping of the vines and does not allow the fruit to come in contact with soil and avoid fruit rot. High temperature leads to fungal wilting in pointed gourd, spray 1.5 ml Carbendazim in 1 litre water. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxychloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Banana: For **leaf spot/sigatoka** as a precautionary measure spray carbendazim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalnil @ 2.0 g / litre of water. For **pseudo stem weevil** Suggested to give stem injection with 5 ml Chloropyrifos or Dimethoate or spray 2.5 ml of Chloropyrifos per litre of water.

Mango: To control **fruit drop**, spray Borax @2gm per lit. of water and apply Planofix @2ml per 5 lit. of water. For control of mango **fruit borer** spray Neem oil 3ml + chlorpyrifos 1 ml per litre of water along with stickers at marble stage of the fruit. The spraying should be preferably carried out during evening hours. To control **bacterial canker**, three sprays of streptomycin (100 ppm) or Agrimycin-100 (100 ppm) after first visual symptom at 10- days intervals is recommended.

Animal husbandry: Make necessary arrangement for clean and sufficient drinking water for the animals.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-289

Date- 19/04/2022

AGROMET ADVISORY BULLETIN FOR TIGIRIA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0 mm rainfall during the last week. The mean daily maximum and minimum temperature was **38.2°C (1°C from normal)** and **25.5°C (+1.4°C from normal)**, respectively. Mean relative humidity of the district was around 66-76%.

Forecasted period: Today(8.30IST) to 24th April (8.30IST)2022

Weather advisory: *Very light to light rainfall / Thundershower is forecasted for Thursday and Friday and Dry weather is forecasted for other 3 days in the block.* The day temperature will be **33-38°C** and night temperature will be **25-28°C** for upcoming 5 days. Maximum and minimum relative humidity will be 77-89% and 18-36% respectively. **There will be clear to partly cloudy sky for next 5 days.**

Block	Tigiria (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	19/04/2022	20/04/2022	21/04/2022	22/04/2022	23/04/2022
Rainfall(mm)	0	0	6	3	0
T-Max(°C)	38	37	33	37	36
T-Min(°C)	25	26	27	28	27
Cloud Cover(octa)	1	1	3	3	2
RH Max (%)	89	86	82	77	85
RH Min (%)	18	22	36	26	23
Wind Speed (Kmph)	16	21	23	17	14
Wind Direction (deg)	180	180	193	162	193

Agro met Advisory

General advisory: After completion of harvesting of rabi crops and before attending ploughing operation, farmers are advised to collect the soil sample from the field & send the sample to nearest soil testing laboratory for nutrient analysis. Soil testing should be done once in three years in rainfed situation and every three cropping season in irrigated situation.

Pulses: Harvest matured pulses before rainy day. Bacterial leaf blight are likely to increase in present situation in pulses due to realized rainfall; if observe, spray Streptomycin or Plantomycin @ 2 g/litre of water as foliar treatment. Rust infestation likely to occur in pulses, if observe spray Mancozeb 35.0%SC 3gram/ltr of water. Present weather is congenial for **Cercospora leaf spot** infestation in black gram crop. On occurrence, immediately spray Agrimycin or Streptomycin 2 g/litre, at an interval of 12-15 days, to effectively reduce pest population.

Summer Maize:

If there was infestation of **Stem rot** at vegetative stage as rainfall was realized in past, farmers may drench the crop root with Mancozeb 35.0%SC 3gram/ltr of water or validamycin 2ml/litre of water and there may increase the infestation of **Red hairy caterpillar** in maize after rainfall, to minimize the infestation farmers may spray Neem oil 5ml/ltr of water or Chlorpyrifos 20 % EC @ 2ml/litre of water.

Vegetables:

Okra: To control **Yellow Vain Mosaic Virus (YVMV)** in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease.

Brinjal: There are chances of infestation of **Epilachna Beetle/Haddabeetle** in Brinjal and cucurbits crop; to manage Epilachna Beetle in cucurbits, spray neem-based pesticide (Azadirachtin) 1500 PPM @ 3ml/litre of water at early stage of infestation; to manage Epilachna Beetle chemically, spray Chlorpyrifos 20 % EC @ 2ml/litre of water or Profenophos 50 % EC @ 2ml/litre of water.

FRUIT AND SHOOT BORER IN BRINJAL: Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@250g/acre at 15 days interval. If **phomopsis blight** will occur in brinjal, then spray zineb75%WP 2gram/ltr of water. **Cercospora leaf spot** may be found after rainfall, if observe, then spray Carbendazim or zineb75%WP 2gram/ltr of water. If **helicoverpa** infestation will be found, spray Flunendiamide or indoxacarb 2ml/ltr of water.

Cucurbits: To manage **Red Pumpkin Beetle** in Cucurbit crops spray Cyantraniliprole 10.26% OD @ 400-ml/acre or Imidacloprid 17.8% SL @ 60-ml/acre or Chlorpyrifos 20% EC @ 400-ml/acre. **Late Blight** may happen due to rising day temperature. There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Don't go for weeding. The grasses and weeds will help in maintaining the humidity and it is safe for creeping of the vines and does not allow the fruit to come in contact with soil and avoid fruit rot. High temperature leads to fungal wilting in pointed gourd, spray 1.5 ml Carbendazim in 1 litre water. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxychloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Banana: For **leaf spot/sigatoka** as a precautionary measure spray carbendazim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalnil @ 2.0 g / litre of water. For **pseudo stem weevil** Suggested to give stem injection with 5 ml Chlorpyrifos or Dimethoate or spray 2.5 ml of Chlorpyrifos per litre of water.

Mango: To control **fruit drop**, spray Borax @2gm per lit. of water and apply Planofix @2ml per 5 lit. of water. For control of mango **fruit borer** spray Neem oil 3ml + chlorpyrifos 1 ml per litre of water along with stickers at marble stage of the fruit. The spraying should be preferably carried out during evening hours. To control **bacterial canker**, three sprays of streptomycin (100 ppm) or Agrimycin-100 (100 ppm) after first visual symptom at 10- days intervals is recommended.

Animal husbandry: Make necessary arrangement for clean and sufficient drinking water for the animals.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-289
 Date- 19/04/2022

				
MEGHDOOT APP	DAMINI APP	FACEBOOK (Damu Cuttack)	BLOCK SPECIFIC WHAT'S APP GROUPS	@DAMUCUTTACK



- Kindly install and use **“Meghdoot mobile App”** for Cuttack district weather forecast, warnings and weather based agromet advisory services.

<https://play.google.com/store/apps/details?id=com.aas.meghdoot>

- Kindly install and use **“Damini mobile App”** for location specific for lightening forecast and warnings.

<https://play.google.com/store/apps/details?id=com.lightening.live.damini>

- Follow the **block specific whats app groups** and join in your block wise groups for getting weather based agromet advisory services twice in a week (Tuesday and Friday) and other weather warning based farm management advisories.
- Follow the **Official Website links** to get the weather based agromet advisory service bulletins (**English & Odia**) twice in a week (Tuesday and Friday).

1	ICAR-NRRI, Cuttack, Odisha	https://icar-nrri.in/agromate/	
2	Agromet division, IMD, Pune	https://imdagrmet.gov.in/imdproject/AGIndex.php	

Dr Sujata Sethy
 (Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
 SMS (Agrometeorology)
 Contact No-9861445080/7008453427