



Cuttack District Block Wise Agro-met Advisory Service Bulletin

*Friday 13th May, 2022
(For the period 13th to 17th May, 2022)*



*Issued by
District Agromet Unit,
KrishiVigyan Kendra Cuttack*





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-295
Date- 13/05/2022

STANDARD PRECIPITATION INDEX AND NDVI MAP

Standard Precipitation Index (SPI)	Normalized difference vegetation Index (NDVI)
<p>Standardised Precipitation Index for the period 14th April 2022 to 11th May 2022</p> <p>MINISTRY OF EARTH SCIENCES INDIA METEOROLOGICAL DEPARTMENT HYDROMET SECTION, CRS PUNE</p> <p>STANDARDIZED PRECIPITATION INDEX FOR THE PERIOD 14 APR 2022 TO 11 MAY 2022</p> <p>SPI VALUES</p> <ul style="list-style-type: none"> < -2 (Extremely Dry) -1.50 to -1.99 (Extremely Dry) -1.00 to -1.49 (Moderately Dry) -0.50 to -0.99 (Mildly Dry) 0 to 0.49 (Near Normal) 0.50 to 0.99 (Moderately Wet) 1.00 to 1.49 (Extremely Wet) 1.50 to 1.99 (Extremely Wet) > 2 (Extremely Wet) No Data 	<p>NOAA/VIIIRS/BLENDED NDVI Composite ending on Week no 18 (30.04.2022 to 06.05.2022) over Agricultural region</p> <p style="text-align: center;">Odisha</p> <p>Agriculture vigour is moderate over the state.</p> <p>Legend:</p> <ul style="list-style-type: none"> -0.9999 - 0 0 - 0.05 0.05 - 0.1 0.1 - 0.15 0.15 - 0.2 0.2 - 0.25 0.25 - 0.3 0.3 - 0.35 0.35 - 0.4 0.4 - 0.45 0.45 - 0.5 0.5 - 0.55 0.55 - 0.6 > 0.6 <p>Bare soil, rocks clouds, rain, snow, etc.</p> <p>Moderate</p> <p>Good</p> <p>Very good</p>
<p>As per SPI value (cumulative 4weeks -14 April 2022 – 11 May 2022), mildly wet condition was there in Cuttack district.</p>	<p>According to NDVI map (30.04.2022 to 06.05.2022) the agriculture vigor was very light to light over most places of Cuttack district</p>

LEGEND (Medium range block level rainfall forecast)		
	No Rainfall	00 mm
	Very light rainfall	01-05 mm
	Light rainfall	06-10 mm
	Light to moderate rainfall	11-30 mm
	Moderate to heavy rainfall	31-65 mm
	Heavy to very heavy rainfall	66-115 mm
	Extremely heavy rainfall	>116 mm

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com





Bulletin No.-295
Date- 13/05/2022

Past week (06-12 May, 2022) cumulative rainfall departure% from normal and (07 -13 May) average maximum temperature and its departure from normal for Cuttack district



Medium range (Today (8.30IST) to Wednesday (8.30IST)) Rainfall (mm) forecast for Cuttack district



Dr Sujata Sethy
 (Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-295
Date- 13/05/2022

Met Sub division	Extended range outlook for subsequent 1 week (6 th to 12 th days) (18 th May to 24 th May 2022)		
Odisha	<i>Parameter</i>	<i>Departure (%)</i>	<i>Category</i>
	Rainfall	56.37	Below Normal
	<i>Parameter</i>	<i>Anomaly</i>	<i>Category</i>
	Maximum temperature	0.91	Below Normal
	Minimum temperature	0.94	Normal

General Agromet Advisory:- As there will be **below normal rainfall is forecasted in the subsequent week after 5 days in Odisha condition** so farmers are advised to go for summer ploughing in the dry days and may go for application of fertilizer and plant protection chemical in the subsequent week to control disease and pest as rainfall and humid weather was realized in the past week.

Long range forecast for the 2021 South-west Monsoon seasonal rainfall and Monthly Rainfall Forecast for May 2022

- Southwest monsoon seasonal (June to September) rainfall over the country **as a whole is most likely to be normal (96 to 104 % of Long Period Average (LPA))**.
- Quantitatively, the monsoon seasonal (June to September) rainfall is likely to be 99% of the Long Period Average (LPA) with a model error of $\pm 5\%$. The LPA of the season rainfall over the country as a whole for the period 1971-2020 is 87 cm.
- As per monthly outlook for May month there will **be above normal rainfall likely to occur at Odisha condition.**
- Hence Farmers should prepare the crop plan depending upon the land type, soil fertility and Variety availability, irrigation facility.*




Fig.1. Probability forecast of tercile categories (below normal, normal, and above normal) for the seasonal rainfall over India during the 2022 southwest monsoon season (June-September). The figure illustrates the most likely categories as well as their probabilities. The white shaded areas represent climatological probabilities for all the tercile categories. The probabilities were derived using the MME forecast prepared from a group of four best coupled climate models. (Tercile categories have equal climatological probabilities, of 33.33% each).




Fig.2. Probability forecast of tercile categories (below normal, normal, and above normal) for the rainfall over India during May 2022. The figure illustrates the most likely categories as well as their probabilities. The dotted area shown in the map climatologically receives very less rainfall and the white shaded areas within the land areas represent climatological probabilities. The probabilities were derived using the MME forecast prepared from a group of coupled climate models. (Tercile categories have equal climatological probabilities, of 33.33% each).

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-295
Date- 13/05/2022

Weather (Past and Forecasted) based Disease and pest in Major Crops and vegetables

Sl No	Crop	Stage and condition of the crop	Anticipated Disease and pest	
1	Maize	Vegetative to tasseling/silking		Maize stalk rot
				Fall army worm
				Bnaded Leaf blight
2	Sugarcane	Vegetative		Red rot
				Wilt
3	Jute	Sowing		Weed
4	Brinjal	Fruiting		Phomopsis Blight
				Cercopora leaf spot
5	Tomato	Fruiting		Bacterial wilt
6	Okra	Flowering to fruiting		fusarium and phytopthera
				Cercospora leaf spot

Dr Sujata Sethy
 (Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area













Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-295
Date- 13/05/2022

7	Cucurbits	Flowering to Fruiting stage		Late blight
				Downy Mildew
				Leaf spot
				Stem Rot
				Fruit rot
8	Banana			Banana leaf spot/sigatoka
				Panama wilt
9	Mango	Fruiting stage		Fruit fly
				Anthracnose
10	Papaya	Fruting		phytophthora crown rot or foot rot

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-295

Date- 13/05/2022

AGROMET ADVISORY BULLETIN FOR ATHAGAD

District Past week average weather condition and Block past week average Rainfall condition

The block received average 7mm and district received average 5mm rainfall during the last week. The mean daily maximum and minimum temperature was 35°C (-2°C from normal) and 25°C (0.4°C from normal), respectively. Mean relative humidity of the district was around 70-85%.

Forecasted period: Today(8.30IST) to 18th May(8.30IST)2022

Weather advisory: Light rainfall / Thundershower is forecasted for Monday and Tuesday, dry weather is forecasted for other 3 days in the block. The day temperature will be 34-37°C and night temperature will be 27°C for upcoming 5 days. Maximum and minimum relative humidity will be 75-83% and 34-45% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Athagad (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	13/05/2022	14/05/2022	15/05/2022	16/05/2022	17/05/2022
Rainfall (mm)	0	0	0	6	7
T-Max (°C)	35	37	34	35	34
T-Min (°C)	27	27	27	27	27
Cloud Cover (octa)	1	2	1	1	2
RH Max (%)	75	83	81	81	83
RH Min (%)	38	34	42	45	43
Wind Speed (kmph)	26	26	27	28	26
Wind Direction (deg)	203	198	198	198	193

Agromet Advisory

General advisory: Farmers are advised to drain out excess water from their field. They may go for plant protection measures and fertilizer application as dry weather is forecasted in upcoming 3 days.

SUMMER PLOUGHING: In dry sunny days after Friday, go for summer ploughing at places receiving soaking rainfall to reduce weed infestation, to destroy eggs and larvae of hibernated soil insects and to increase soil profile recharge.

GREEN MANURING: Optimum time for sowing of green manuring crops like Sesbania and sunhemp in the areas receiving convectional rain. Green manuring crops are sown in medium and low land condition at 6-8 weeks before transplanting of rice. Use 10 kg seeds of Sesbania per acre.. Apply all the recommended dose of phosphorous of rice (100 kg of SSP) to the field during sowing time. Incorporate the green manuring plant at 6-8 week stage for better decomposition before transplanting. It adds 6-7 tons of green matter and in turn 25-30 kg of N to the succeeding rice crop per ha.

Maize:

Prolonged dry spell followed by rain may increase the infestation of fall army worm in maize. To control spray emamectin benzoate (5% S.G.) @ 0.4g per litre of water. If there will be infestation of Stem rot or banded leaf blight at vegetative stage as rainfall is forecasted, farmers may drench the crop root with Mancozeb 35.0%SC 3gram/ltr of water or validamycin 2ml/litre of water.

Vegetables: After rain there may be chances of increase in infestation of wilting in vegetables. To prevent from wilting or partial wilting in brinjal, Chilli and Okra apply Neem or Karanjaa cake and spray (basal + foliar) Tebuconazole fungicide @ 1ml/litre. Foliar spray of 2% DAP + 1% KCl (MOP) is recommended in vegetables.

Okra: Spray Metalaxyl+Mancozeb/Carboxin+Thiram@2g/litre to control rot or soil borne fungal disease. Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenconazole @ 1ml/lit at 15 days interval to control Cercospora leaf spot.

Brinjal:

If phomopsis blight will occur in brinjal, then spray zineb 75%WP 2gram/litre of water. Cercospora leaf spot may be found after rainfall, if observe, then spray Carbendazim or zineb 75%WP 2gram/litre of water.

Cucurbits: Spray of 40 ppm NAA for controlling excessive pre-mature fall of flowering/buds/young developing fruits and pods. There are chances of infestation of Downy Mildew in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Present weather may aggravate fruit and vine rot (symptoms: leaf spot, leaf blight, stem rot, fruit rot) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxychloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit: Banana: In Banana Drain the excess water as soon as possible. Spray 1% KNO₃ or Urea 2% solution 2-3 times during vegetative to flowering stage. Topdressing of booster dose of 80 g MOP + 100 g Urea per plant in two to three splits at monthly intervals. For leaf spot/sigatoka as a precautionary measure spray carbendazim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalnil @ 2.0 g / litre of water. Drenching soil with Metiram 75% WG @ 1.5g/4 lit or carbendazim 12%+ mancozeb 63% @ 2g/lit to prevent Panama wilt.

Mango: Control measures should be taken against fruit fly. Spray the foliage with Tebuconazole 50% + Trifloxystrobin 25% WG @ 1g/lit to control Anthracnose. Spray Deltamethrin 2.8% EC to control fruit fly.

Papaya: After rain due to water logging there may be increase in the infestation of phytophthora crown rot or foot rot in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-295

Date- 13/05/2022

AGROMET ADVISORY BULLETIN FOR BADAMBA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0.4mm and district received average 5mm rainfall during the last week. The mean daily maximum and minimum temperature was **35°C (-2°C from normal)** and 25°C (0.4°C from normal), respectively. Mean relative humidity of the district was around 70-85%.

Forecasted period: Today(8.30IST) to 18th May(8.30IST)2022

Weather advisory: *Light rainfall / Thundershower is forecasted for Monday and Tuesday, dry weather is forecasted for other 3 days in the block.* The day temperature will be 36-39°C and night temperature will be 26°C for upcoming 5 days. Maximum and minimum relative humidity will be 75-84% and 31-40% respectively. **There will be clear to partly cloudy sky for next 5 days.**

Block	Badamba (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	13/05/2022	14/05/2022	15/05/2022	16/05/2022	17/05/2022
Rainfall(mm)	0	0	0	6	7
T-Max(°C)	37	39	36	36	37
T-Min(°C)	26	26	26	26	26
Cloud Cover(octa)	1	1	1	1	2
RH Max (%)	75	81	79	80	84
RH Min (%)	36	31	37	40	39
Wind Speed (Kmph)	21	21	23	24	21
Wind Direction (deg)	198	180	167	167	193

Agromet Advisory

General advisory: Farmers are advised to drain out excess water from their field. They may go for plant protection measures and fertilizer application as dry weather is forecasted in upcoming 3 days.

SUMMER PLOUGHING: In dry sunny days after Friday, go for summer ploughing at places receiving soaking rainfall to reduce weed infestation, to destroy eggs and larvae of hibernated soil insects and to increase soil profile recharge.

GREEN MANURING: Optimum time for sowing of green manuring crops like Sesbania and sunhemp in the areas receiving convectional rain. Green manuring crops are sown in medium and low land condition at 6-8 weeks before transplanting of rice. Use 10 kg seeds of Sesbania per acre.. Apply all the recommended dose of phosphorous of rice (100 kg of SSP) to the field during sowing time. Incorporate the green manuring plant at 6-8 week stage for better decomposition before transplanting. It adds 6-7 tons of green matter and in turn 25-30 kg of N to the succeeding rice crop per ha.

Maize:

Prolonged dry spell followed by rain may increase the infestation of fall army worm in maize. To control spray emamectin benzoate (5% S.G.) @ 0.4g per litre of water. If there will be infestation of **Stem rot or banded leaf blight** at vegetative stage as rainfall is forecasted, farmers may drench the crop root with Mancozeb 35.0%SC 3gram/ltr of water or validamycin 2ml/litre of water.

Vegetables: After rain there may be chances of increase in infestation of wilting in vegetables. To prevent from wilting or partial wilting in brinjal, Chilli and Okra apply Neem or Karanjaa cake and spray (basal + foliar) Tebuconazole fungicide @ 1ml/litre. Foliar spray of 2% DAP + 1% KCl (MOP) is recommended in vegetables.

Okra: Spray Metalaxyl+Mancozeb/Carboxin+Thiram@2g/litre to control **rot or soil borne fungal disease**. Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenoconazole @ 1ml/lit at 15 days interval to control **Cercospora leaf spot**.

Brinjal:

If **phomopsis blight** will occur in brinjal, then spray zineb 75%WP 2gram/litre of water. **Cercospora leaf spot** may be found after rainfall, if observe, then spray Carbendazim or zineb 75%WP 2gram/litre of water.

Cucurbits: Spray of 40 ppm NAA for controlling excessive pre-mature fall of flowering/buds/young developing fruits and pods. There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxychloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit: Banana: In Banana Drain the excess water as soon as possible. Spray 1% KNO₃ or Urea 2% solution 2-3 times during vegetative to flowering stage. Topdressing of booster dose of 80 g MOP + 100 g Urea per plant in two to three splits at monthly intervals. For **leaf spot/sigatoka** as a precautionary measure spray carbendazim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalnil @ 2.0 g / litre of water. Drenching soil with Metiram 75% WG @ 1.5g/4 lit or carbendazim 12%+ mancozeb 63% @ 2g/lit to prevent **Panama wilt**.

Mango: Control measures should be taken against fruit fly. Spray the foliage with Tebuconazole 50% + Trifloxystrobin 25% WG @ 1g/lit to control **Anthracnose**. Spray Deltamethrin 2.8% EC to control **fruit fly**.

Papaya: After rain due to water logging there may be increase in the infestation of **phytophthora crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-295
Date- 13/05/2022

AGROMET ADVISORY BULLETIN FOR BANKI

District Past week average weather condition

The district received average 5mm rainfall during the last week. The mean daily maximum and minimum temperature was **35°C (-2°C from normal)** and 25°C (0.4°C from normal), respectively. Mean relative humidity of the district was around 70-85%.

Forecasted period: Today(8.30IST) to 18th May(8.30IST)2022

Weather advisory:: *Very light rainfall / Thundershower is forecasted for Monday and Tuesday, dry weather is forecasted for other 3 days in the block.* The day temperature will be 34-37°C and night temperature will be 26-27°C for upcoming 5 days. Maximum and minimum relative humidity will be 75-84% and 35-45% respectively. **There will be clear to partly cloudy sky for next 5 days.**

Block	Banki (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	13/05/2022	14/05/2022	15/05/2022	16/05/2022	17/05/2022
Rainfall(mm)	0	0	0	4	5
T-Max(°C)	35	37	34	35	35
T-Min(°C)	27	27	27	27	26
Cloud Cover(octa)	1	2	1	1	2
RH Max (%)	75	83	81	81	84
RH Min (%)	39	35	43	45	43
Wind Speed (Kmph)	25	26	27	28	25
Wind Direction (deg)	201	198	193	180	193

Agromet Advisory

General advisory: Farmers are advised to drain out excess water from their field. They may go for plant protection measures and fertilizer application as dry weather is forecasted in upcoming 3 days.

SUMMER PLOUGHING: In dry sunny days after Friday, go for summer ploughing at places receiving soaking rainfall to reduce weed infestation, to destroy eggs and larvae of hibernated soil insects and to increase soil profile recharge.

GREEN MANURING: Optimum time for sowing of green manuring crops like Sesbania and sunhemp in the areas receiving convectional rain. Green manuring crops are sown in medium and low land condition at 6-8 weeks before transplanting of rice. Use 10 kg seeds of Sesbania per acre.. Apply all the recommended dose of phosphorous of rice (100 kg of SSP) to the field during sowing time. Incorporate the green manuring plant at 6-8 week stage for better decomposition before transplanting. It adds 6-7 tons of green matter and in turn 25-30 kg of N to the succeeding rice crop per ha.

Maize: Prolonged dry spell followed by rain may increase the infestation of fall army worm in maize. To control spray emamectin benzoate (5% S.G.) @ 0.4g per litre of water. If there will be infestation of **Stem rot or banded leaf blight** at vegetative stage as rainfall is forecasted, farmers may drench the crop root with Mancozeb 35.0%SC 3gram/ltr of water or validamycin 2ml/litre of water.

Vegetables: After rain there may be chances of increase in infestation of wilting in vegetables. To prevent from wilting or partial wilting in brinjal, Chilli and Okra apply Neem or Karanjaa cake and spray (basal + foliar) Tebuconazole fungicide @ 1ml/litre. Foliar spray of 2% DAP + 1% KCl (MOP) is recommended in vegetables.

Okra: Spray Metalaxyl+Mancozeb/Carboxin+Thiram@2g/litre to control **rot or soil borne fungal disease**. Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenconazole @ 1ml/lit at 15 days interval to control **Cercospora leaf spot**.

Brinjal:

If **phomopsis blight** will occur in brinjal, then spray zineb 75% WP 2gram/litre of water. **Cercospora leaf spot** may be found after rainfall, if observe, then spray Carbendazim or zineb 75% WP 2gram/litre of water.

Cucurbits: Spray of 40 ppm NAA for controlling excessive pre-mature fall of flowering/buds/young developing fruits and pods. There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxichloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit: Banana: In Banana Drain the excess water as soon as possible. Spray 1% KNO₃ or Urea 2% solution 2-3 times during vegetative to flowering stage. Topdressing of booster dose of 80 g MOP + 100 g Urea per plant in two to three splits at monthly intervals. For **leaf spot/sigatoka** as a precautionary measure spray carbendazim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalnil @ 2.0 g / litre of water. Drenching soil with Metiram 75% WG @ 1.5g/4 lit or carbendazim 12%+ mancozeb 63% @ 2g/lit to prevent **Panama wilt**.

Mango: Control measures should be taken against fruit fly. Spray the foliage with Tebuconazole 50% + Trifloxystrobin 25% WG @ 1g/lit to control **Anthraxnose**. Spray Deltamethrin 2.8% EC to control **fruit fly**.

Papaya: After rain due to water logging there may be increase in the infestation of **phytophthora crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA INDIA METEOROLOGICAL DEPARTMENT DISTRICT AGROMET UNIT KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-295

Date- 13/05/2022

AGROMET ADVISORY BULLETIN FOR BANKI-DAMPARA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 8mm and district received average 5mm rainfall during the last week. The mean daily maximum and minimum temperature was 35°C (-2°C from normal) and 25°C (0.4°C from normal), respectively. Mean relative humidity of the district was around 70-85%.

Forecasted period: Today(8.30IST) to 18th May(8.30IST)2022

Weather Advisory: Very light rainfall / Thundershower is forecasted for Monday and Tuesday, dry weather is forecasted for other 3 days in the block. The day temperature will be 34-37°C and night temperature will be 27°C for upcoming 5 days. Maximum and minimum relative humidity will be 75-84% and 36-46% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Banki-Dampara (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	13/05/2022	14/05/2022	15/05/2022	16/05/2022	17/05/2022
Rainfall(mm)	0	0	0	5	5
T-Max(°C)	35	37	34	35	34
T-Min(°C)	27	27	27	27	27
Cloud Cover(octa)	1	2	1	1	2
RH Max (%)	75	83	82	81	84
RH Min (%)	39	36	44	46	44
Wind Speed (Kmph)	25	26	27	28	25
Wind Direction (deg)	203	198	193	193	193

Agromet Advisory

General advisory: Farmers are advised to drain out excess water from their field. They may go for plant protection measures and fertilizer application as dry weather is forecasted in upcoming 3 days.

SUMMER PLOUGHING: In dry sunny days after Friday, go for summer ploughing at places receiving soaking rainfall to reduce weed infestation, to destroy eggs and larvae of hibernated soil insects and to increase soil profile recharge.

GREEN MANURING: Optimum time for sowing of green manuring crops like Sesbania and sunhemp in the areas receiving convectional rain. Green manuring crops are sown in medium and low land condition at 6-8 weeks before transplanting of rice. Use 10 kg seeds of Sesbania per acre.. Apply all the recommended dose of phosphorous of rice (100 kg of SSP) to the field during sowing time. Incorporate the green manuring plant at 6-8 week stage for better decomposition before transplanting. It adds 6-7 tons of green matter and in turn 25-30 kg of N to the succeeding rice crop per ha.

Maize: Prolonged dry spell followed by rain may increase the infestation of fall army worm in maize. To control spray emamectin benzoate (5% S.G.) @ 0.4g per litre of water. If there will be infestation of **Stem rot or banded leaf blight** at vegetative stage as rainfall is forecasted, farmers may drench the crop root with Mancozeb 35.0%SC 3gram/ltr of water or validamycin 2ml/litre of water.

Vegetables: After rain there may be chances of increase in infestation of wilting in vegetables. To prevent from wilting or partial wilting in brinjal, Chilli and Okra apply Neem or Karanjaa cake and spray (basal + foliar) Tebuconazole fungicide @ 1ml/litre. Foliar spray of 2% DAP + 1% KCl (MOP) is recommended in vegetables.

Okra: Spray Metalaxyl+Mancozeb/Carboxin+Thiram@2g/litre to control **rot or soil borne fungal disease**. Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenconazole @ 1ml/lit at 15 days interval to control **Cercospora leaf spot**.

Brinjal:

If **phomopsis blight** will occur in brinjal, then spray zineb 75% WP 2gram/litre of water. **Cercospora leaf spot** may be found after rainfall, if observe, then spray Carbendazim or zineb 75% WP 2gram/litre of water.

Cucurbits: Spray of 40 ppm NAA for controlling excessive pre-mature fall of flowering/buds/young developing fruits and pods. There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxichloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit: Banana: In Banana Drain the excess water as soon as possible. Spray 1% KNO₃ or Urea 2% solution 2-3 times during vegetative to flowering stage. Topdressing of booster dose of 80 g MOP + 100 g Urea per plant in two to three splits at monthly intervals. For **leaf spot/sigatoka** as a precautionary measure spray carbendazim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalnil @ 2.0 g / litre of water. Drenching soil with Metiram 75% WG @ 1.5g/4 lit or carbendazim 12%+ mancozeb 63% @ 2g/lit to prevent **Panama wilt**.

Mango: Control measures should be taken against fruit fly. Spray the foliage with Tebuconazole 50% + Trifloxystrobin 25% WG @ 1g/lit to control **Anthraxnose**. Spray Deltamethrin 2.8% EC to control **fruit fly**.

Papaya: After rain due to water logging there may be increase in the infestation of **phytophthora crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-295
Date- 13/05/2022

AGROMET ADVISORY BULLETIN FOR BARANGA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 12mm and district received average 5mm rainfall during the last week. The mean daily maximum and minimum temperature was 35°C (-2°C from normal) and 25°C (0.4°C from normal), respectively. Mean relative humidity of the district was around 70-85%.

Forecasted period: Today(8.30IST) to 18th May (8.30IST)2022

Weatheradvisory: Very light rainfall / Thundershower is forecasted for Monday and Tuesday, dry weather is forecasted for other 3days in the block. The day temperature will be 33-34°C and night temperature will be 27-28°C for upcoming 5 days. Maximum and minimum relative humidity will be 78-83% and 43-50% respectively. There will clear to partly cloudy sky for next 5days.

Block	Baranga (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	13/05/2022	14/05/2022	15/05/2022	16/05/2022	17/05/2022
Rainfall(mm)	0	0	0	4	4
T-Max(°C)	33	34	33	33	33
T-Min(°C)	27	27	27	27	28
Cloud Cover	1	2	2	1	3
RH Max (%)	78	83	83	81	81
RH Min (%)	43	44	48	50	48
Wind Speed (Kmph)	29	29	28	29	27
Wind Direction (deg)	203	203	203	203	203

Agromet Advisory

General advisory: Farmers are advised to drain out excess water from their field. They may go for plant protection measures and fertilizer application as dry weather is forecasted in upcoming 3days.

SUMMER PLOUGHING: In dry sunny days after Friday, go for summer ploughing at places receiving soaking rainfall to reduce weed infestation, to destroy eggs and larvae of hibernated soil insects and to increase soil profile recharge.

GREEN MANURING: Optimum time for sowing of green manuring crops like Sesbania and sunhemp in the areas receiving convectional rain. Green manuring crops are sown in medium and low land condition at 6-8 weeks before transplanting of rice. Use 10 kg seeds of Sesbania per acre.. Apply all the recommended dose of phosphorous of rice (100 kg of SSP) to the field during sowing time. Incorporate the green manuring plant at 6-8 week stage for better decomposition before transplanting. It adds 6-7 tons of green matter and in turn 25-30 kg of N to the succeeding rice crop per ha.

Maize: Prolonged dry spell followed by rain may increase the infestation of fall army worm in maize. To control spray emamectin benzoate (5% S.G.) @ 0.4g per litre of water. If there will be infestation of **Stem rot or banded leaf blight** at vegetative stage as rainfall is forecasted, farmers may drench the crop root with Mancozeb 35.0%SC 3gram/ltr of water or validamycin 2ml/litre of water.

Vegetables:

After rain there may be chances of increase in infestation of wilting in vegetables. To prevent from wilting or partial wilting in brinjal, Chilli and Okra apply Neem or Karanjaa cake and spray (basal + foliar) Tebuconazole fungicide @ 1ml/litre. Foliar spray of 2% DAP + 1% KCl (MOP) is recommended in vegetables .

Okra: Spray Metalaxyl+Mancozeb/Carboxin+Thiram@2g/litre to control **rot or soil borne fungal disease**.Foliar spraying of Trifloxystrobin25% + tebuconazole 50% WG @ 1g/lit or Difenconazole @ 1ml/lit at 15 days interval to control **Cercospora leaf spot**.

Brinjal:

If **phomopsis blight** will occur in brinjal, then spray zineb75%WP 2gram/litre of water.**Cercospora leaf spot** may be found after rainfall, if observe, then spray Carbendazim or zineb75%WP 2gram/litre of water.

Cucurbits: Spray of 40 ppm NAA for controlling excessive pre-mature fall of flowering/buds/young developing fruits and pods There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxychloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit: Banana: In Banana Drain the excess water as soon as possible. Spray 1% KNO₃ or Urea 2% solution 2-3 times during vegetative to flowering stage. Topdressing of booster dose of 80 g MOP + 100 g Urea per plant in two to three splits at monthly intervals. For **leaf spot/sigatoka** as a precautionary measure spray carbendazim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalnil @ 2.0 g/ litre of water. Drenching soil with Metiram 75% WG @ 1.5g/4 lit or carbendazim 12%+ mancozeb 63% @ 2g/lit to prevent **Panama wilt**.

Mango: Control measures should be taken against fruit fly. Spray the foliage with Tebuconazole 50% + Trifloxystrobin 25% WG @ 1g/lit to control **Anthraxnose**. Spray Deltamethrin 2.8% EC to control **fruit fly**.

Papaya: After rain due to water logging there may be increase in the infestation of **phytophthora crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)



Bulletin No.-295
Date- 13/05/2022

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

AGROMET ADVISORY BULLETIN FOR CUTTACK SADAR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 2mm and district received average 5mm rainfall during the last week. The mean daily maximum and minimum temperature was **35°C (-2°C from normal)** and **25°C (0.4°C from normal)**, respectively. Mean relative humidity of the district was around 70-85%.

Forecasted period: Today(8.30IST) to 18th May (8.30IST)2022

Weather advisory: *Very light rainfall / Thundershower is forecasted for Monday and Tuesday, dry weather is forecasted for other 3 days in the block.* The day temperature will be **33-35°C** and night temperature will be **27°C** for upcoming 5 days. Maximum and minimum relative humidity will be 77-83% and 40-47% respectively. **There will be clear to partly cloudy sky for next 5 days.**

Block	Cuttack sadar (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	13/05/2022	14/05/2022	15/05/2022	16/05/2022	17/05/2022
Rainfall(mm)	0	0	0	3	4
T-Max(°C)	34	35	33	34	33
T-Min(°C)	27	27	27	27	27
Cloud Cover(octa)	1	2	2	1	3
RH Max (%)	77	83	82	81	81
RH Min (%)	41	40	46	47	46
Wind Speed (Kmph)	28	28	28	29	27
Wind Direction (deg)	203	203	201	203	198

Agromet Advisory

General advisory: Farmers are advised to drain out excess water from their field. They may go for plant protection measures and fertilizer application as dry weather is forecasted in upcoming 3 days.

SUMMER PLOUGHING: In dry sunny days after Friday, go for summer ploughing at places receiving soaking rainfall to reduce weed infestation, to destroy eggs and larvae of hibernated soil insects and to increase soil profile recharge.

GREEN MANURING: Optimum time for sowing of green manuring crops like Sesbania and sunhemp in the areas receiving convectional rain. Green manuring crops are sown in medium and low land condition at 6-8 weeks before transplanting of rice. Use 10 kg seeds of Sesbania per acre.. Apply all the recommended dose of phosphorous of rice (100 kg of SSP) to the field during sowing time. Incorporate the green manuring plant at 6-8 week stage for better decomposition before transplanting. It adds 6-7 tons of green matter and in turn 25-30 kg of N to the succeeding rice crop per ha.

Suagrcane: For **Red rot and wilt disease**, spray the crop with Mancozeb 0.3% or carbendazim 0.1% or Azoxystrobin 18.2%SC/ Difenconazole 11.45SC @ 1ml per liter water in dry days.

Vegetables:

After rain there may be chances of increase in infestation of wilting in vegetables. To prevent from wilting or partial wilting in brinjal, Chilli and Okra apply Neem or Karanjaa cake and spray (basal + foliar) Tebuconazole fungicide @ 1ml/liter. Foliar spray of 2% DAP + 1% KCl (MOP) is recommended in vegetables .

Okra: Spray Metalxyl+Mancozeb/Carboxin+Thiram@2g/litre to control **rot or soil borne fungal disease**.Foliar spraying of Trifloxystrobin25% + tebuconazole 50% WG @ 1g/lit or Difenconazole @ 1ml/lit at 15 days interval to control **Cercospora leaf spot**.

Brinjal:If phomopsis blight will occur in brinjal, then spray zineb75%WP 2gram/litr of water.**Cercopspera leaf spot** may be found after rainfall, if observe, then spray Carbendazim or zineb75%WP 2gram/litr of water.

Cucurbits: Spray of 40 ppm NAA for controlling excessive pre-mature fall of flowering/buds/young developing fruits and pods There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre. Pointed gourd is at flowering and fruiting stage. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxylchloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Banana: In Banana Drain the excess water as soon as possible. Spray 1% KNO₃ or Urea 2% solution 2-3 times during vegetative to flowering stage. Topdressing of booster dose of 80 g MOP + 100 g Urea per plant in two to three splits at monthly intervals. For **leaf spot/sigatoka** as a precautionary measure spray carbendizim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalanil@ 2.0 g/ litre of water. Drenching soil with Metiram 75% WG @ 1.5g/4 lit or carbendazim 12%+ mancozeb 63% @ 2g/lit to prevent **Panama wilt**.

Mango: Control measures should be taken against fruit fly. Spray the foliage with Tebuconazole 50% + Trifloxystrobin 25% WG @ 1g/lit to control **Anthracnose**. Spray Deltamethrin 2.8% EC to control **fruit fly**.

Papaya: After rain due to water logging there may be increase in the infestation of **phytopthera crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

Dr Sujata Sathy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)



Bulletin No.-295
Date- 13/05/2022

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

AGROMET ADVISORY BULLETIN FOR KANTAPADA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 5mm and district received average 5mm rainfall during the last week. The mean daily maximum and minimum temperature was **35°C (-2°C from normal)** and **25°C (0.4°C from normal)**, respectively. Mean relative humidity of the district was around 70-85%.

Forecasted period: Today(8.30IST) to 18th May(8.30IST)2022

Weather advisory: *Very light rainfall / Thundershower is forecasted for Monday and Tuesday, dry weather is forecasted for other 3 days in the block.* The day temperature will be 32-34°C and night temperature will be 27-28°C for upcoming 5 days. Maximum and minimum relative humidity will be 79-83% and 47-52% respectively. **There will be clear to partly cloudy sky for next 5 days.**

Block	Kantapada (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	13/05/2022	14/05/2022	15/05/2022	16/05/2022	17/05/2022
Rainfall(mm)	0	0	0	2	3
T-Max(°C)	33	34	32	33	33
T-Min(°C)	27	27	27	27	28
Cloud Cover(octa)	0	2	1	1	3
RH Max (%)	79	82	83	81	80
RH Min (%)	47	49	52	52	52
Wind Speed (Kmph)	29	29	28	28	27
Wind Direction (deg)	203	203	203	203	203

Agromet Advisory

General advisory: Farmers are advised to drain out excess water from their field. They may go for plant protection measures and fertilizer application as dry weather is forecasted in upcoming 3 days.

SUMMER PLOUGHING: In dry sunny days after Friday, go for summer ploughing at places receiving soaking rainfall to reduce weed infestation, to destroy eggs and larvae of hibernated soil insects and to increase soil profile recharge.

GREEN MANURING: Optimum time for sowing of green manuring crops like Sesbania and sunhemp in the areas receiving convectional rain. Green manuring crops are sown in medium and low land condition at 6-8 weeks before transplanting of rice. Use 10 kg seeds of Sesbania per acre.. Apply all the recommended dose of phosphorous of rice (100 kg of SSP) to the field during sowing time. Incorporate the green manuring plant at 6-8 week stage for better decomposition before transplanting. It adds 6-7 tons of green matter and in turn 25-30 kg of N to the succeeding rice crop per ha.

Suagrcane: For **Red rot and wilt disease**, spray the crop with Mancozeb 0.3% or carbendazim 0.1% or Azoxystrobin 18.2%SC/ Difenconazole 11.45SC @ 1ml per liter water in dry days.

Vegetables:

After rain there may be chances of increase in infestation of wilting in vegetables. To prevent from wilting or partial wilting in brinjal, Chilli and Okra apply Neem or Karanjaa cake and spray (basal + foliar) Tebuconazole fungicide @ 1ml/liter. Foliar spray of 2% DAP + 1% KCl (MOP) is recommended in vegetables.

Okra: Spray Metalaxyl+Mancozeb/Carboxin+Thiram@2g/litre to control **rot or soil borne fungal disease**. Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenconazole @ 1ml/lit at 15 days interval to control **Cercospora leaf spot**.

Brinjal: If phomopsis blight will occur in brinjal, then spray zineb 75% WP 2gram/litre of water. **Cercospora leaf spot** may be found after rainfall, if observe, then spray Carbendazim or zineb 75% WP 2gram/litre of water.

Cucurbits: Spray of 40 ppm NAA for controlling excessive pre-mature fall of flowering/buds/young developing fruits and pods. There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre. Pointed gourd is at flowering and fruiting stage. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxychloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Banana: In Banana Drain the excess water as soon as possible. Spray 1% KNO₃ or Urea 2% solution 2-3 times during vegetative to flowering stage. Topdressing of booster dose of 80 g MOP + 100 g Urea per plant in two to three splits at monthly intervals. For **leaf spot/sigatoka** as a precautionary measure spray carbendazim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalnil @ 2.0 g/ litre of water. Drenching soil with Metiram 75% WG @ 1.5g/4 lit or carbendazim 12%+ mancozeb 63% @ 2g/lit to prevent **Panama wilt**.

Mango: Control measures should be taken against fruit fly. Spray the foliage with Tebuconazole 50% + Trifloxystrobin 25% WG @ 1g/lit to control **Anthraco-nose**. Spray Deltamethrin 2.8% EC to control **fruit fly**.

Papaya: After rain due to water logging there may be increase in the infestation of **phytophthora crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

AGROMET ADVISORY BULLETIN FOR MAHANGA



Bulletin No.-295

Date- 13/05/2022

District Past week average weather condition and Block past week average Rainfall condition

The block received average 3mm and district received average 5mm rainfall during the last week. The mean daily maximum and minimum temperature was **35°C (-2°C from normal)** and **25°C (0.4°C from normal)**, respectively. Mean relative humidity of the district was around 70-85%.

Forecasted period: Today(8.30IST) to 18th May(8.30IST)2022

Weather advisory: *Light rainfall / Thundershower is forecasted for Monday and Tuesday, dry weather is forecasted for other 3 days in the block.* The day temperature will be and it will be **34-35°C** and night temperature will be **27°C** for upcoming 5 days. Maximum and minimum relative humidity will be 77-82% and 43-48% respectively. **There will be clear to partly cloudy sky for next 5 days.**

Block	Mahanga (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	13/05/2022	14/05/2022	15/05/2022	16/05/2022	17/05/2022
Rainfall(mm)	0	0	0	6	7
T-Max(°C)	34	35	34	34	34
T-Min(°C)	27	27	27	27	27
Cloud Cover(octal)	1	2	2	1	2
RH Max (%)	77	82	81	80	78
RH Min (%)	43	43	48	48	47
Wind Speed (Kmph)	28	28	27	28	25
Wind Direction (deg)	203	203	203	203	201

Agromet Advisory

General advisory: Farmers are advised to drain out excess water from their field. They may go for plant protection measures and fertilizer application as dry weather is forecasted in upcoming 3 days.

SUMMER PLOUGHING: In dry sunny days after Friday, go for summer ploughing at places receiving soaking rainfall to reduce weed infestation, to destroy eggs and larvae of hibernated soil insects and to increase soil profile recharge.

GREEN MANURING: Optimum time for sowing of green manuring crops like Sesbania and sunhemp in the areas receiving convectional rain. Green manuring crops are sown in medium and low land condition at 6-8 weeks before transplanting of rice. Use 10 kg seeds of Sesbania per acre.. Apply all the recommended dose of phosphorous of rice (100 kg of SSP) to the field during sowing time. Incorporate the green manuring plant at 6-8 week stage for better decomposition before transplanting. It adds 6-7 tons of green matter and in turn 25-30 kg of N to the succeeding rice crop per ha.

Jute:

The weather is optimum for Jute sowing. Before sowing seeds are to be treated with 2 g Carbendazim or 3 g Thiram/kg seeds to avoid root and stem rot. In Jute field, spray neem oil 10,000ppm @ 3ml/litre of water with sticker or Chlorpyrifos 50 EC @ 2ml/litre of water in the evening hour for preventive measure; **drain out excess water from field.** Realised rain will aggravate **weed problem** in early sown crop. To control grass weeds, apply Quizalofop-pethyl @ 2ml/litre of water at 2-3 leaf stage of weeds

Vegetables:

After rain there may be chances of increase in infestation of wilting in vegetables. To prevent from wilting or partial wilting in brinjal, Chilli and Okra apply Neem or Karanjaa cake and spray (basal + foliar) Tebuconazole fungicide @ 1ml/liter. Foliar spray of 2% DAP + 1% KCl (MOP) is recommended in vegetables.

Okra: Spray Metalyxyl+Mancozeb/Carboxin+Thiram @ 2g/litre to control **rot or soil borne fungal disease**. Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenconazole @ 1ml/lit at 15 days interval to control **Cercospora leaf spot**.

Brinjal: If phomopsis blight will occur in brinjal, then spray zineb 75% WP 2gram/lit of water. **Cercospora leaf spot** may be found after rainfall, if observe, then spray Carbendazim or zineb 75% WP 2gram/lit of water.

Cucurbits: Spray of 40 ppm NAA for controlling excessive pre-mature fall of flowering/buds/young developing fruits and pods. There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Fruit: Banana: In Banana Drain the excess water as soon as possible. Spray 1% KNO₃ or Urea 2% solution 2-3 times during vegetative to flowering stage. Topdressing of booster dose of 80 g MOP + 100 g Urea per plant in two to three splits at monthly intervals. For **leaf spot/sigatoka** as a precautionary measure spray carbendazim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalnil @ 2.0 g / litre of water. Drenching soil with Metiram 75% WG @ 1.5g/4 lit or carbendazim 12% + mancozeb 63% @ 2g/lit to prevent **Panama wilt**.

Mango: Control measures should be taken against fruit fly. Spray the foliage with Tebuconazole 50% + Trifloxystrobin 25% WG @ 1g/lit to control **Anthraco-nose**. Spray Deltamethrin 2.8% EC to control **fruit fly**.

Papaya: After rain due to water logging there may be increase in the infestation of **phytophthora crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-295

Date- 13/05/2022

District Past week average weather condition and Block past week average Rainfall condition

The block received average 2mm and district received average 5mm rainfall during the last week. The mean daily maximum and minimum temperature was **35°C (-2°C from normal)** and **25°C (0.4°C from normal)**, respectively. Mean relative humidity of the district was around 70-85%.

Forecasted period: Today(8.30IST) to 18th May(8.30IST)2022

Weather advisory: *Light rainfall / Thundershower is forecasted for Monday and Tuesday, dry weather is forecasted for other 3 days in the block.* The day temperature will be **38-40°C** and night temperature will be **25-26°C** for upcoming 5 days. Maximum and minimum relative humidity will be 76-82% and 25-30% respectively. **There will be clear to partly cloudy sky for next 5 days.**

Block	Narasinghpur (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	13/05/2022	14/05/2022	15/05/2022	16/05/2022	17/05/2022
Rainfall(mm)	0	0	0	7	8
T-Max(°C)	38	40	38	38	38
T-Min(°C)	25	26	26	26	26
Cloud Cover(octa)	1	1	1	1	3
RH Max (%)	76	80	78	81	82
RH Min (%)	28	25	27	30	28
Wind Speed (Kmph)	13	14	14	16	13
Wind Direction (deg)	180	180	167	167	161

Agromet Advisory

General advisory: Farmers are advised to drain out excess water from their field. They may go for plant protection measures and fertilizer application as dry weather is forecasted in upcoming 3 days.

SUMMER PLOUGHING: In dry sunny days after Friday, go for summer ploughing at places receiving soaking rainfall to reduce weed infestation, to destroy eggs and larvae of hibernated soil insects and to increase soil profile recharge.

GREEN MANURING: Optimum time for sowing of green manuring crops like Sesbania and sunhemp in the areas receiving convectional rain. Green manuring crops are sown in medium and low land condition at 6-8 weeks before transplanting of rice. Use 10 kg seeds of Sesbania per acre.. Apply all the recommended dose of phosphorous of rice (100 kg of SSP) to the field during sowing time. Incorporate the green manuring plant at 6-8 week stage for better decomposition before transplanting. It adds 6-7 tons of green matter and in turn 25-30 kg of N to the succeeding rice crop per ha.

Suagrcane: For **Red rot and wilt disease**, spray the crop with Mancozeb 0.3% or carbendazim 0.1% or Azoxystrobin 18.2%SC/ Difenconazole 11.45SC @ 1ml per liter water in dry days.

Vegetables:

After rain there may be chances of increase in infestation of wilting in vegetables. To prevent from wilting or partial wilting in brinjal, Chilli and Okra apply Neem or Karanjaa cake and spray (basal + foliar) Tebuconazole fungicide @ 1ml/liter. Foliar spray of 2% DAP + 1% KCl (MOP) is recommended in vegetables .

Okra: Spray Metalxyl+Mancozeb/Carboxin+Thiram@2g/litre to control **rot or soil borne fungal disease**.Foliar spraying of Trifloxystrobin25% + tebuconazole 50% WG @ 1g/lit or Difenconazole @ 1ml/lit at 15 days interval to control **Cercospora leaf spot**.

Brinjal:If phomopsis blight will occur in brinjal, then spray zineb75%WP 2gram/litr of water.**Cercopspera leaf spot** may be found after rainfall, if observe, then spray Carbendazim or zineb75%WP 2gram/litr of water.

Cucurbits: Spray of 40 ppm NAA for controlling excessive pre-mature fall of flowering/buds/young developing fruits and pods There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre. Pointed gourd is at flowering and fruiting stage. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxychloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Banana: In Banana Drain the excess water as soon as possible. Spray 1% KNO₃ or Urea 2% solution 2-3 times during vegetative to flowering stage. Topdressing of booster dose of 80 g MOP + 100 g Urea per plant in two to three splits at monthly intervals. For **leaf spot/sigatoka** as a precautionary measure spray carbendizim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalanil@ 2.0 g/ litre of water. Drenching soil with Metiram 75% WG @1.5g/4 lit or carbendazim 12%+ mancozeb 63% @ 2g/lit to prevent **Panama wilt**.

Mango: Control measures should be taken against fruit fly. Spray the foliage with Tebuconazole 50% + Trifloxystrobin 25% WG @ 1g/lit to control **Anthracnose**. Spray Deltamethrin 2.8% EC to control **fruit fly**.

Papaya: After rain due to water logging there may be increase in the infestation of **phytophthora crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-295
Date- 13/05/2022

AGROMET ADVISORY BULLETIN FOR NIALI

District Past week average weather condition and Block past week average Rainfall condition

The block received average 1mm and district received average 5mm rainfall during the last week. The mean daily maximum and minimum temperature was **35°C (-2°C from normal)** and **25°C (0.4°C from normal)**, respectively. Mean relative humidity of the district was around 70-85%.

Forecasted period: Today(8.30IST) to 18th May(8.30IST)2022

Weather advisory: *Very light rainfall / Thundershower is forecasted for Monday and Tuesday, dry weather is forecasted for other 3 days in the block.* The day temperature will be **32-34°C** and night temperature will be **27-28°C** for upcoming 5 days. Maximum and minimum relative humidity will be 79-83% and 50-56% respectively. **There will be clear to partly cloudy sky for next 5 days.**

Block	Niali (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	13/05/2022	14/05/2022	15/05/2022	16/05/2022	17/05/2022
Rainfall(mm)	0	0	0	2	2
T-Max(°C)	33	34	32	33	33
T-Min(°C)	27	27	27	27	28
Cloud Cover(octa)	0	2	2	1	3
RH Max (%)	79	82	83	81	80
RH Min (%)	50	53	55	56	55
Wind Speed(Kmph)	30	29	28	28	27
Wind Direction(deg)	203	203	203	203	203

Agromet Advisory

General advisory: Farmers are advised to drain out excess water from their field. They may go for plant protection measures and fertilizer application as dry weather is forecasted in upcoming 3 days.

SUMMER PLOUGHING: In dry sunny days after Friday, go for summer ploughing at places receiving soaking rainfall to reduce weed infestation, to destroy eggs and larvae of hibernated soil insects and to increase soil profile recharge.

GREEN MANURING: Optimum time for sowing of green manuring crops like Sesbania and sunhemp in the areas receiving convectional rain. Green manuring crops are sown in medium and low land condition at 6-8 weeks before transplanting of rice. Use 10 kg seeds of Sesbania per acre.. Apply all the recommended dose of phosphorous of rice (100 kg of SSP) to the field during sowing time. Incorporate the green manuring plant at 6-8 week stage for better decomposition before transplanting. It adds 6-7 tons of green matter and in turn 25-30 kg of N to the succeeding rice crop per ha.

Suagrcane: For **Red rot and wilt disease**, spray the crop with Mancozeb 0.3% or carbendazim 0.1% or Azoxystrobin 18.2%SC/ Difenconazole 11.45SC @ 1ml per liter water in dry days.

Vegetables:

After rain there may be chances of increase in infestation of wilting in vegetables. To prevent from wilting or partial wilting in brinjal, Chilli and Okra apply Neem or Karanjaa cake and spray (basal + foliar) Tebuconazole fungicide @ 1ml/liter. Foliar spray of 2% DAP + 1% KCl (MOP) is recommended in vegetables .

Okra: Spray Metalaxyl+Mancozeb/Carboxin+Thiram@2g/litre to control **rot or soil borne fungal disease**.Foliar spraying of Trifloxystrobin25% + tebuconazole 50% WG @ 1g/lit or Difenconazole @ 1ml/lit at 15 days interval to control **Cercospora leaf spot**.

Brinjal:If **phomopsis blight** will occur in brinjal, then spray zineb75%WP 2gram/litr of water.**Cercospora leaf spot** may be found after rainfall, if observe, then spray Carbendazim or zineb75%WP 2gram/litr of water.

Cucurbits: Spray of 40 ppm NAA for controlling excessive pre-mature fall of flowering/buds/young developing fruits and pods There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre. Pointed gourd is at flowering and fruiting stage. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxichloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Banana: In Banana Drain the excess water as soon as possible. Spray 1% KNO₃ or Urea 2% solution 2-3 times during vegetative to flowering stage. Topdressing of booster dose of 80 g MOP + 100 g Urea per plant in two to three splits at monthly intervals. For **leaf spot/sigatoka** as a precautionary measure spray carbendazim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalnil@ 2.0 g/ litre of water. Drenching soil with Metiram 75% WG @ 1.5g/4 lit or carbendazim 12%+ mancozeb 63% @ 2g/lit to prevent **Panama wilt**.

Mango: Control measures should be taken against fruit fly. Spray the foliage with Tebuconazole 50% + Trifloxystrobin 25% WG @ 1g/lit to control **Anthracnose**. Spray Deltamethrin 2.8% EC to control **fruit fly**.

Papaya: After rain due to water logging there may be increase in the infestation of **phytophthora crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)



Bulletin No.-295
Date- 13/05/2022

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

AGROMET ADVISORY BULLETIN FOR NISCHINTAKOILI

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0.4mm and district received average 5mm rainfall during the last week. The mean daily maximum and minimum temperature was **35°C (-2°C from normal)** and 25°C (0.4°C from normal), respectively. Mean relative humidity of the district was around 70-85%.

Forecasted period: Today(8.30IST) to 18th May(8.30IST)2022

Weatheradvisory: *Very light rainfall / Thundershower is forecasted for Monday and Tuesday, dry weather is forecasted for other 3 days in the block.* The day temperature will be 33-35°C and night temperature will be 27-28°C for upcoming 5 days. Maximum and minimum relative humidity will be 77-82% and 44-49% respectively. **There will be clear to partly cloudy sky for next 5 days.**

Block	Nischintakoili (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	13/05/2022	14/05/2022	15/05/2022	16/05/2022	17/05/2022
Rainfall(mm)	0	0	0	4	5
T-Max(°C)	34	35	33	34	34
T-Min(°C)	27	27	27	27	28
Cloud Cover(octa)	1	2	2	1	3
RH Max (%)	77	82	82	80	78
RH Min (%)	44	45	49	49	48
Wind Speed (Kmph)	28	28	27	28	26
Wind Direction (deg)	203	203	203	203	201

Agromet Advisory

General advisory: Farmers are advised to drain out excess water from their field. They may go for plant protection measures and fertilizer application as dry weather is forecasted in upcoming 3 days.

SUMMER PLOUGHING: In dry sunny days after Friday, go for summer ploughing at places receiving soaking rainfall to reduce weed infestation, to destroy eggs and larvae of hibernated soil insects and to increase soil profile recharge.

GREEN MANURING: Optimum time for sowing of green manuring crops like Sesbania and sunhemp in the areas receiving convectional rain. Green manuring crops are sown in medium and low land condition at 6-8 weeks before transplanting of rice. Use 10 kg seeds of Sesbania per acre.. Apply all the recommended dose of phosphorous of rice (100 kg of SSP) to the field during sowing time. Incorporate the green manuring plant at 6-8 week stage for better decomposition before transplanting. It adds 6-7 tons of green matter and in turn 25-30 kg of N to the succeeding rice crop per ha.

Suagrcane: For **Red rot and wilt disease**, spray the crop with Mancozeb 0.3% or carbendazim 0.1% or Azoxystrobin 18.2%SC/ Difenconazole 11.45SC @ 1ml per liter water in dry days.

Vegetables:

After rain there may be chances of increase in infestation of wilting in vegetables. To prevent from wilting or partial wilting in brinjal, Chilli and Okra apply Neem or Karanjaa cake and spray (basal + foliar) Tebuconazole fungicide @ 1ml/liter. Foliar spray of 2% DAP + 1% KCl (MOP) is recommended in vegetables .

Okra: Spray Metalaxyl+Mancozeb/Carboxin+Thiram@2g/litre to control **rot or soil borne fungal disease**.Foliar spraying of Trifloxystrobin25% + tebuconazole 50% WG @ 1g/lit or Difenconazole @ 1ml/lit at 15 days interval to control **Cercospora leaf spot**.

Brinjal:If phomopsis blight will occur in brinjal, then spray zineb75%WP 2gram/litr of water.**Cercopspera leaf spot** may be found after rainfall, if observe, then spray Carbendazim or zineb75%WP 2gram/litr of water.

Cucurbits: Spray of 40 ppm NAA for controlling excessive pre-mature fall of flowering/buds/young developing fruits and pods There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre. Pointed gourd is at flowering and fruiting stage. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxychloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Banana: In Banana Drain the excess water as soon as possible. Spray 1% KNO₃ or Urea 2% solution 2-3 times during vegetative to flowering stage. Topdressing of booster dose of 80 g MOP + 100 g Urea per plant in two to three splits at monthly intervals. For **leaf spot/sigatoka** as a precautionary measure spray carbendizim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalnil@ 2.0 g/ litre of water. Drenching soil with Metiram 75% WG @ 1.5g/4 lit or carbendazim 12%+ mancozeb 63% @ 2g/lit to prevent **Panama wilt**.

Mango: Control measures should be taken against fruit fly. Spray the foliage with Tebuconazole 50% + Trifloxystrobin 25% WG @ 1g/lit to control **Anthracoise**. Spray Deltamethrin 2.8% EC to control **fruit fly**.

Papaya: After rain due to water logging there may be increase in the infestation of **phytopthera crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-295
Date- 13/05/2022

AGROMET ADVISORY BULLETIN FOR SALEPUR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0.3mm and district received average 5mm rainfall during the last week. The mean daily maximum and minimum temperature was **35°C (-2°C from normal)** and **25°C (0.4°C from normal)**, respectively. Mean relative humidity of the district was around 70-85%.

Forecasted period: Today(8.30IST) to 18th May(8.30IST)2022

Weather advisory: *Very light rainfall / Thundershower is forecasted for Monday and Tuesday, dry weather is forecasted for other 3days in the block.* The day temperature will be **33-35°C** and night temperature will be **27-28°C** for upcoming 5 days. Maximum and minimum relative humidity will be **77-82%** and **44-49%** respectively. **There will be clear to partly cloudy sky for next 5days.**

Block	Salepur(VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	13/05/2022	14/05/2022	15/05/2022	16/05/2022	17/05/2022
Rainfall(mm)	0	0	0	4	5
T-Max(°C)	34	35	33	34	33
T-Min(°C)	27	27	27	27	28
Cloud Cover(octa)	1	2	2	1	3
RH Max (%)	77	82	82	81	78
RH Min (%)	44	44	48	49	48
Wind Speed (Kmph)	28	28	27	28	26
Wind Direction (deg)	203	203	203	203	201

Agromet Advisory

General advisory: Farmers are advised to drain out excess water from their field. They may go for plant protection measures and fertilizer application as dry weather is forecasted in upcoming 3days.

SUMMER PLOUGHING: In dry sunny days after Friday, go for summer ploughing at places receiving soaking rainfall to reduce weed infestation, to destroy eggs and larvae of hibernated soil insects and to increase soil profile recharge.

GREEN MANURING: Optimum time for sowing of green manuring crops like Sesbania and sunhemp in the areas receiving convectional rain. Green manuring crops are sown in medium and low land condition at 6-8 weeks before transplanting of rice. Use 10 kg seeds of Sesbania per acre.. Apply all the recommended dose of phosphorous of rice (100 kg of SSP) to the field during sowing time. Incorporate the green manuring plant at 6-8 week stage for better decomposition before transplanting. It adds 6-7 tons of green matter and in turn 25-30 kg of N to the succeeding rice crop per ha.

Maize: Prolonged dry spell followed by rain may increase the infestation of fall army worm in maize. To control spray emamectin benzoate (5% S.G.) @ 0.4g per litre of water. If there will be infestation of Stem rot or banded leaf blight at vegetative stage as rainfall is forecasted, farmers may drench the crop root with Mancozeb 35.0%SC 3gram/ltr of water or validamycin 2ml/litre of water.

Jute: The weather is optimum for Jute sowing. Before sowing seeds are to be treated with 2 g Carbendazim or 3 g Thiram/kg seeds to avoid root and stem rot. In Jute field, spray neem oil 10,000ppm @ 3ml/litre of water with sticker or Chlorpyrifos 50 EC @ 2ml/litre of water in the evening hour for preventive measure; drain out excess water from field. Realised rain will aggravate weed problem in early sown crop. To control grass weeds, apply Quizalofop-pethyl@2ml/litre of water at 2-3 leaf stage of weeds

Vegetables:

After rain there may be chances of increase in infestation of wilting in vegetables. To prevent from wilting or partial wilting in brinjal, Chilli and Okra apply Neem or Karanjaa cake and spray (basal + foliar) Tebuconazole fungicide @ 1ml/liter. Foliar spray of 2% DAP + 1% KCl (MOP) is recommended in vegetables .

Okra: Spray Metalxyl+Mancozeb/Carboxin+Thiram@2g/litre to control rot or soil borne fungal disease. Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenconazole @ 1ml/lit at 15 days interval to control Cercospora leaf spot.

Brinjal: If phomopsis blight will occur in brinjal, then spray zineb 75% WP 2gram/litre of water. Cercospora leaf spot may be found after rainfall, if observe, then spray Carbendazim or zineb 75% WP 2gram/litre of water.

Cucurbits: Spray of 40 ppm NAA for controlling excessive pre-mature fall of flowering/buds/young developing fruits and pods. There are chances of infestation of Downy Mildew in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Fruit: Banana: In Banana Drain the excess water as soon as possible. Spray 1% KNO₃ or Urea 2% solution 2-3 times during vegetative to flowering stage. Topdressing of booster dose of 80 g MOP + 100 g Urea per plant in two to three splits at monthly intervals. For leaf spot/sigatoka as a precautionary measure spray carbendazim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalnil @ 2.0 g / litre of water. Drenching soil with Metiram 75% WG @ 1.5g/4 lit or carbendazim 12%+ mancozeb 63% @ 2g/lit to prevent Panama wilt.

Mango: Control measures should be taken against fruit fly. Spray the foliage with Tebuconazole 50% + Trifloxystrobin 25% WG @ 1g/lit to control Anthracnose. Spray Deltamethrin 2.8% EC to control fruit fly.

Papaya: After rain due to water logging there may be increase in the infestation of phytophthora crown rot or foot rot in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-295

Date- 13/05/2022

AGROMET ADVISORY BULLETIN FOR TANGI-CHOUDWAR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 1.3mm and district received average 5mm rainfall during the last week. The mean daily maximum and minimum temperature was **35°C (-2°C from normal)** and 25°C (0.4°C from normal), respectively. Mean relative humidity of the district was around 70-85%.

Forecasted period: Today(8.30IST) to 18th May(8.30IST)2022

Weather advisory: Light rainfall / Thundershower is forecasted for Monday and Tuesday, dry weather is forecasted for other 3 days in the block. The day temperature will be 33-35°C and night temperature will be 29°C for upcoming 5 days. Maximum and minimum relative humidity will be 80-83% and 56-65% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Tangi-Choudwar (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	13/05/2022	14/05/2022	15/05/2022	16/05/2022	17/05/2022
Rainfall(mm)	0	0	0	6	7
T-Max(°C)	34	35	33	34	34
T-Min(°C)	29	29	29	29	29
Cloud Cover(octa)	1	2	4	3	2
RH Max (%)	80	82	83	82	82
RH Min (%)	56	58	65	63	60
Wind Speed (Kmph)	31	30	28	30	26
Wind Direction (deg)	203	203	203	203	203

Agromet Advisory

General advisory: Farmers are advised to drain out excess water from their field. They may go for plant protection measures and fertilizer application as dry weather is forecasted in upcoming 3 days.

SUMMER PLOUGHING: In dry sunny days after Friday, go for summer ploughing at places receiving soaking rainfall to reduce weed infestation, to destroy eggs and larvae of hibernated soil insects and to increase soil profile recharge.

GREEN MANURING: Optimum time for sowing of green manuring crops like Sesbania and sunhemp in the areas receiving convectional rain. Green manuring crops are sown in medium and low land condition at 6-8 weeks before transplanting of rice. Use 10 kg seeds of Sesbania per acre.. Apply all the recommended dose of phosphorous of rice (100 kg of SSP) to the field during sowing time. Incorporate the green manuring plant at 6-8 week stage for better decomposition before transplanting. It adds 6-7 tons of green matter and in turn 25-30 kg of N to the succeeding rice crop per ha.

Maize: Prolonged dry spell followed by rain may increase the infestation of fall army worm in maize. To control spray emamectin benzoate (5% S.G.) @ 0.4g per litre of water. If there will be infestation of Stem rot or banded leaf blight at vegetative stage as rainfall is forecasted, farmers may drench the crop root with Mancozeb 35.0%SC 3gram/ltr of water or validamycin 2ml/litre of water.

Vegetables:

After rain there may be chances of increase in infestation of wilting in vegetables. To prevent from wilting or partial wilting in brinjal, Chilli and Okra apply Neem or Karanjaa cake and spray (basal + foliar) Tebuconazole fungicide @ 1ml/liter. Foliar spray of 2% DAP + 1% KCl (MOP) is recommended in vegetables.

Okra: Spray Metalaxyl+Mancozeb/Carboxin+Thiram@2g/litre to control rot or soil borne fungal disease. Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenconazole @ 1ml/lit at 15 days interval to control Cercospora leaf spot.

Brinjal:

If phomopsis blight will occur in brinjal, then spray zineb 75% WP 2gram/litre of water. Cercospora leaf spot may be found after rainfall, if observe, then spray Carbendazim or zineb 75% WP 2gram/litre of water.

Cucurbits:

Spray of 40 ppm NAA for controlling excessive pre-mature fall of flowering/buds/young developing fruits and pods. There are chances of infestation of Downy Mildew in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Fruit:

Banana: In Banana Drain the excess water as soon as possible. Spray 1% KNO₃ or Urea 2% solution 2-3 times during vegetative to flowering stage. Topdressing of booster dose of 80 g MOP + 100 g Urea per plant in two to three splits at monthly intervals. For leaf spot/sigatoka as a precautionary measure spray carbendazim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalnil @ 2.0 g/ litre of water. Drenching soil with Metiram 75% WG @ 1.5g/4 lit or carbendazim 12%+ mancozeb 63% @ 2g/lit to prevent Panama wilt.

Mango: Control measures should be taken against fruit fly. Spray the foliage with Tebuconazole 50% + Trifloxystrobin 25% WG @ 1g/lit to control Anthracnose. Spray Deltamethrin 2.8% EC to control fruit fly.

Papaya: After rain due to water logging there may be increase in the infestation of phytophthora crown rot or foot rot in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-295
Date- 13/05/2022

AGROMET ADVISORY BULLETIN FOR TIGIRIA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 7mm and district received average 5mm rainfall during the last week. The mean daily maximum and minimum temperature was 35°C (-2°C from normal) and 25°C (0.4°C from normal), respectively. Mean relative humidity of the district was around 70-85%.

Forecasted period: Today(8.30IST) to 18th May(8.30IST)2022

Weather advisory: Light rainfall / Thundershower is forecasted for Monday and Tuesday, dry weather is forecasted for other 3 days in the block. The day temperature will be 35-38°C and night temperature will be 26-27°C for upcoming 5 days. Maximum and minimum relative humidity will be 74-84% and 32-42% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Tigiria (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	13/05/2022	14/05/2022	15/05/2022	16/05/2022	17/05/2022
Rainfall(mm)	0	0	0	7	7
T-Max(°C)	36	38	35	36	36
T-Min(°C)	27	27	27	27	26
Cloud Cover(octa)	1	2	1	1	2
RH Max (%)	74	82	80	81	84
RH Min (%)	37	32	40	42	40
Wind Speed (Kmph)	24	24	26	27	24
Wind Direction (deg)	201	193	180	180	193

Agro met Advisory

General advisory: Farmers are advised to drain out excess water from their field. They may go for plant protection measures and fertilizer application as dry weather is forecasted in upcoming 3 days.

SUMMER PLOUGHING: In dry sunny days after Friday, go for summer ploughing at places receiving soaking rainfall to reduce weed infestation, to destroy eggs and larvae of hibernated soil insects and to increase soil profile recharge.

GREEN MANURING: Optimum time for sowing of green manuring crops like Sesbania and sunhemp in the areas receiving convectional rain. Green manuring crops are sown in medium and low land condition at 6-8 weeks before transplanting of rice. Use 10 kg seeds of Sesbania per acre. Apply all the recommended dose of phosphorous of rice (100 kg of SSP) to the field during sowing time. Incorporate the green manuring plant at 6-8 week stage for better decomposition before transplanting. It adds 6-7 tons of green matter and in turn 25-30 kg of N to the succeeding rice crop per ha.

Maize: Prolonged dry spell followed by rain may increase the infestation of fall army worm in maize. To control spray emamectin benzoate (5% S.G.) @ 0.4g per litre of water. If there will be infestation of Stem rot or banded leaf blight at vegetative stage as rainfall is forecasted, farmers may drench the crop root with Mancozeb 35.0%SC 3gram/ltr of water or validamycin 2ml/litre of water.

Vegetables:

After rain there may be chances of increase in infestation of wilting in vegetables. To prevent from wilting or partial wilting in brinjal, Chilli and Okra apply Neem or Karanjaa cake and spray (basal + foliar) Tebuconazole fungicide @ 1ml/litre. Foliar spray of 2% DAP + 1% KCl (MOP) is recommended in vegetables.

Okra:

Spray Metalaxyl+Mancozeb/Carboxin+Thiram@2g/litre to control rot or soil borne fungal disease. Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenoconazole @ 1ml/lit at 15 days interval to control Cercospora leaf spot.

Brinjal:

If phomopsis blight will occur in brinjal, then spray zineb 75% WP 2gram/litre of water. Cercospora leaf spot may be found after rainfall, if observe, then spray Carbendazim or zineb 75% WP 2gram/litre of water.

Cucurbits: Spray of 40 ppm NAA for controlling excessive pre-mature fall of flowering/buds/young developing fruits and pods. There are chances of infestation of Downy Mildew in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Fruit:

Banana: In Banana Drain the excess water as soon as possible. Spray 1% KNO₃ or Urea 2% solution 2-3 times during vegetative to flowering stage. Topdressing of booster dose of 80 g MOP + 100 g Urea per plant in two to three splits at monthly intervals. For leaf spot/sigatoka as a precautionary measure spray carbendazim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalnil @ 2.0 g/ litre of water. Drenching soil with Metiram 75% WG @ 1.5g/4 lit or carbendazim 12%+ mancozeb 63% @ 2g/lit to prevent Panama wilt.

Mango: Control measures should be taken against fruit fly. Spray the foliage with Tebuconazole 50% + Trifloxystrobin 25% WG @ 1g/lit to control Anthracnose. Spray Deltamethrin 2.8% EC to control fruit fly.

Papaya: After rain due to water logging there may be increase in the infestation of phytophthora crown rot or foot rot in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-295
 Date- 13/05/2022

				
MEGHDOOT APP	DAMINI APP	FACEBOOK (Damu Cuttack)	BLOCK SPECIFIC WHAT'S APP GROUPS	@DAMUCUTTACK



- Kindly install and use **“Meghdoot mobile App”** for Cuttack district weather forecast, warnings and weather based agromet advisory services.

<https://play.google.com/store/apps/details?id=com.aas.meghdoot>

- Kindly install and use **“Damini mobile App”** for location specific for lightening forecast and warnings.

<https://play.google.com/store/apps/details?id=com.lightening.live.damini>

- Follow the **block specific whats app groups** and join in your block wise groups for getting weather based agromet advisory services twice in a week (Tuesday and Friday) and other weather warning based farm management advisories.
- Follow the **Official Website links** to get the weather based agromet advisory service bulletins (**English & Odia**) twice in a week (Tuesday and Friday).

1	ICAR-NRRI, Cuttack, Odisha	https://icar-nrri.in/agromate/	
2	Agromet division, IMD, Pune	https://imdagrmet.gov.in/imdproject/AGIndex.php	

Dr Sujata Sethy
 (Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
 SMS (Agrometeorology)
 Contact No-9861445080/7008453427