




Cuttack District Block Wise Agro-met Advisory Service Bulletin

Tuesday 19th July, 2022
(For the period 19th to 23rd July, 2022)



Issued by
District Agromet Unit,
KrishiVigyan Kendra Cuttack







GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com








Bulletin No.-314
Date- 19/07/2022

Past week (12-18 July, 2022) cumulative rainfall departure% from normal and (13-19 July) average maximum and minimum temperature for Cuttack district



Medium range (Today (8.30IST) to Sunday (8.30IST)) Rainfall (mm) forecast for Cuttack district



Dr Sujata Sethy
 (Nodal officer)



**Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area**



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-314
Date- 19/07/2022

Met Sub division	Extended range outlook for subsequent 1 week (6 th to 12 th days) (24-30 July 2022)		
Odisha	<i>Parameter</i>	<i>Departure (%)</i>	<i>Category</i>
	Rainfall	19.80	Normal
	<i>Parameter</i>	<i>Anomaly</i>	<i>Category</i>
	Maximum temperature	0.44	Normal
	Minimum temperature	0.15	Normal

General Agromet Advisory:- There will be normal rainfall in subsequent week after 5 days in Odisha condition, so farmers are advised to drain out excess water from their rice nurseries and also should complete direct as well as nursery sowing (dry nursery) of kharif rice in that period. They may go for transplanting activity as there will be above normal rainfall condition in subsequent week.

Updated Long range forecast for the 2021 South-west Monsoon seasonal rainfall and Monthly Rainfall Forecast for July 2022

- Southwest monsoon seasonal (June to September) rainfall over the country as a whole is most likely to be normal (96 to 104 % of Long Period Average (LPA)).
- Quantitatively, the monsoon seasonal (June to September) rainfall over the country as a whole is likely to be 103% of the Long Period Average (LPA) with a model error of $\pm 4\%$. The LPA of the seasonal rainfall over the country as a whole based on data of 1971-2020 is 87 cm.
- As per probabilistic forecast of rainfall, there will be below normal rainfall during Monsoon season and below normal rainfall during July month in Odisha condition.
- Hence Farmers should prepare the crop plan depending upon the land type, soil fertility and Variety availability, irrigation facility.
- Farmers may prefer short to medium duration rice variety for this kharif season and may avoid upland rice cultivation.

Tercile probability rainfall forecast for 2022 southwest monsoon season




Fig.1. Updated Probability forecast of tercile categories* (below normal, normal, and above normal) of rainfall over India during monsoon season (June -September), 2022. The figure illustrates the most likely categories as well as their probabilities. The white shaded areas within the land area represent climatological probabilities.

*Tercile categories have equal climatological probabilities of 33.33% each.

probability rainfall forecast for 2022 JUL




Fig.1. Probability forecast of tercile categories* (below normal, normal and above normal) for the rainfall over India during July 2022. The figure illustrates the most likely categories as well as their probabilities. The white shaded areas within the land area represent climatological probabilities. The probabilities were derived using the MME forecast prepared from a group of coupled climate models. (Tercile categories have equal climatological probabilities of 33.33% each).

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427













GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-314
Date- 19/07/2022

Weather (Past and Forecasted) based Disease and pest in Major Crops and vegetables

Sl No	Crop	Stage and condition of the crop	Anticipated Disease and pest	
1	Rice	Nursery stage		Thrips
				Seedling blight
				Sheath blight
2	Sugarcane	Vegetative to Tillering		Early shot borer
				Top borer
				Smut
				Inter node borer
3	Maize	5 week stage		Stem fly
				Leaf blight
3	Jute	30-50days		Stem rot

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427














GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT

KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-314
Date- 19/07/2022

				Indigo caterpillar
4	Brinjal, tomato, chilli	seedling		Seedling rot
				Stem rot
5	Tomato and brinjal	Fruting		Bacterial wilt
6	chilli	fruting		thrips
				Leaf curl
7	okra	Fruting		Fruit and shoot borer
				White fly
				Jassid
8	Banana			Sigatoka
9	Papaya			Yellow mosaic virus

Dr Sujata Sethy
(Nodal officer)



**Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area**



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-314
Date- 19/07/2022

AGROMET ADVISORY BULLETIN FOR ATHAGAD

District Past week average weather condition and Block past week average Rainfall condition

The block received average 17mm and district received average 21mm rainfall during the last week. The mean daily maximum and minimum temperature was 32°C (-0.7°C from normal) and 25°C (+1.1°C from normal), respectively. Mean relative humidity was around 88-95%.

Forecasted period: Today(8.30IST) to 24th July(8.30IST)2022

Weather advisory: Moderate with Isolated heavy rainfall is forecasted for Friday and Saturday and light to moderate rainfall is forecasted for other 3 days in the block. The day temperature will be 31-33°C and night temperature will be 25-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 81-85% and 50-56% respectively. **There will be partly cloudy to overcast sky for next 5 days.**

Block	Athagad (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	19/07/2022	20/07/2022	21/07/2022	22/07/2022	23/07/2022
Rainfall (mm)	2	9	18	37	38
T-Max (°C)	32	31	32	33	32
T-Min (°C)	25	25	26	26	25
Cloud Cover (octa)	7	6	8	8	8
RH Max (%)	81	83	85	84	85
RH Min (%)	51	50	56	55	56
Wind Speed (kmph)	15	12	11	8	9
Wind Direction (deg)	225	212	203	203	199

Agromet Advisory

General advisory: Drain out excess water from rice nurseries, orchards, maize and jute fields. Postpone irrigation, intercultural operations, application of plant protection measures and fertilizers to the standing crops from Thursday to Saturday and they may carry out weeding, pesticide and fertilizer application in upcoming 2 days. Keep the harvested produce at safer places. Cover the vegetable seedling with polythene to protect them from heavy shower.

Kharifrice:(DIRECT SEEDED RICE):

Direct seeded rice are at three to four week stage. As light to moderate amount of rainfall is forecasted in the upcoming days, farmers should go for beushanning. Apply 35 kg urea/ac at beushanning followed by weeding and gap filling. Drain out the excess water at the time of fertilizer application and pond it after one day. If basal fertilizer not applied, apply 18 kg urea, 100kg ssp and 26 kg MOP while beushanning followed by top dressing of 35 kg urea after 7-10 days.

TRANSPLANTED RICE:

Considering the anticipated rainfall, paddy transplanting may be completed before 30th July. Apply 400-gram Chlorantraniliprole 0.4 % G or 800-gram Cartap Hydrochloride 4% GR in the 10 decimal nursery area 7 days before transplanting to manage gall midge, stem borer, caseworm, leaf folder and root knot nematode up to 3 weeks after transplanting. Intermittent dry and wet weather condition, is favourable for incidence of **BLB** and **Sheath blight** in early sown nursery seedlings. Go for foliar spray of Hexaconazole with Plantomycin@1.0 g/l to prevent the spread of the disease. Prevailing high temperature condition may lead to infestation of **rice thrips** in rice nursery. Apply Fipronil 5SC@2ml/l of water or Thiomethoxam@0.4g/ litre of water.

Maize:

Stem fly in Maize infects the plants from sowing to 5 weeks after sowing, temperature between 25-30 deg. celsius and humidity above 60% favour pest incidence. In vegetative stage to control stem fly spray Methyl demeton @1 ml/lit or Dimethoate @ 2ml/lit or apply Carbofuran 3G granuals @ 14 kg/acre. High humid conditions, temperature between 18-27 Deg. Celsius favours the occurrence of **leaf blight** in Maize. To control spray Azoxystrobin + Mancozeb (Pluton) @ 300 g/ac.

ARHAR: In arhar crop, for early variety apply 35 kg DAP, 4 kg Urea and 14 kg MOP as basal fertilizer per acre and for late variety apply 52 kg DAP, 6 kg Urea and 14 kg MOP as basal fertilizer per acre in dry days.

VEGETABLES:

June-july is the best time for sowing of tomato in raised bed for nursery seedlings. Do seed treatment with Trichoderma viride @ 4g/kg seeds. The seed treatment must be done 24 hours before sowing of seeds.

- Due to forecast of rain, Chances of **fruit rot** infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever **sucking pests** incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. **To prevent of soil borne diseases like seedling rot, stem rot and wilt diseases apply 3g of Copper oxychloride per litre of water around the main stem.**
- Incidence of **thrips** may observe in chilli, if observe spray Spinosad 45% SC @ 0.4 ml/litre of water or Imidacloprid 17.8% SL @ 0.3 ml/litre of water in clean and dry weather. The present weather conditions are favourable for Upward and downward leafcurl in chilli. To control spray phosalon (Zolone) 3 ml. or diphenthiuron @ 1g./L of water.
- To control **fruit and shoot borer** in okra, Spray Bacillus thuringiensis 5% W.P @1.5 gram/lit of water. Present weather condition favours incidence of white fly and jassid. To control this apply Thiomethoxam@0.4g/ litre of water.

Fruits: Spray mancozeb 2g/litre, to control sigatoka leaf spot in **banana**. To control **papaya yellow mosaic virus**, spray Imidachloprid @ 0.5 ml / litre.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-314
Date- 19/07/2022

AGROMET ADVISORY BULLETIN FOR BADAMBA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 22mm and district received average 21mm rainfall during the last week. The mean daily maximum and minimum temperature was 32°C (-0.7°C from normal) and 25°C (+1.1°C from normal), respectively. Mean relative humidity was around 88-95%.

Forecasted period: Today(8.30IST) to 24th July(8.30IST)2022

Weather advisory: Moderate with isolated heavy rainfall is forecasted for Friday and Saturday and light to moderate rainfall is forecasted for other 3 days in the block. The day temperature will be 31-32°C and night temperature will be 24-25°C for upcoming 5 days. Maximum and minimum relative humidity will be 82-86% and 51-59% respectively. **There will be partly cloudy to overcast sky for next 5 days.**

Block	Badamba (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	19/07/2022	20/07/2022	21/07/2022	22/07/2022	23/07/2022
Rainfall(mm)	2	10	18	38	36
T-Max(°C)	32	31	32	32	32
T-Min(°C)	24	25	25	25	25
Cloud Cover(octa)	6	6	8	8	8
RH Max (%)	82	84	86	83	86
RH Min (%)	54	51	59	56	53
Wind Speed (Kmph)	12	10	9	7	9
Wind Direction (deg)	234	225	205	207	225

Agromet Advisory

General advisory: Drain out excess water from rice nurseries, orchards, maize and jute fields. Postpone irrigation, intercultural operations, application of plant protection measures and fertilizers to the standing crops from Thursday to Saturday and they may carry out weeding, pesticide and fertilizer application in upcoming 2 days. Keep the harvested produce at safer places. Cover the vegetable seedling with polythene to protect them from heavy shower.

Khari rice: (DIRECT SEEDED RICE):

Direct seeded rice are at three to four week stage. As light to moderate amount of rainfall is forecasted in the upcoming days, farmers should go for beushanning. Apply 35 kg urea/ac at beushanning followed by weeding and gap filling. Drain out the excess water at the time of fertilizer application and pond it after one day. If basal fertilizer not applied, apply 18 kg urea, 100kg ssp and 26 kg MOP while beushanning followed by top dressing of 35 kg urea after 7-10 days.

TRANSPLANTED RICE:

Considering the anticipated rainfall, paddy transplanting may be completed before 30th July. Apply 400-gram Chlorantraniliprole 0.4 % G or 800-gram Cartap Hydrochloride 4% GR in the 10 decimal nursery area 7 days before transplanting to manage gall midge, stem borer, caseworm, leaf folder and root knot nematode up to 3 weeks after transplanting. Intermittent dry and wet weather condition, is favourable for incidence of **BLB** and **Sheath blight** in early sown nursery seedlings. Go for foliar spray of Hexaconazole with Plantomycin @ 1.0 g/l to prevent the spread of the disease. Prevailing high temperature condition may lead to infestation of **rice thrips** in rice nursery. Apply Fipronil 5SC @ 2ml/l of water or Thiomethoxam @ 0.4g/ litre of water.

Maize:

Stem fly in Maize infects the plants from sowing to 5 weeks after sowing, temperature between 25-30 deg. celsius and humidity above 60% favour pest incidence. In vegetative stage to control stem fly spray Methyl demeton @ 1 ml/l or Dimethoate @ 2ml/l or apply Carbofuran 3G granuals @ 14 kg/acre. High humid conditions, temperature between 18-27 Deg. Celsius favours the occurrence of **leaf blight** in Maize. To control spray Azoxystrobin + Mancozeb (Pluton) @ 300 g /ac.

ARHAR: In arhar crop, for early variety apply 35 kg DAP, 4 kg Urea and 14 kg MOP as basal fertilizer per acre and for late variety apply 52 kg DAP, 6 kg Urea and 14 kg MOP as basal fertilizer per acre in dry days.

VEGETABLES:

June-july is the best time for sowing of tomato in raised bed for nursery seedlings. Do seed treatment with Trichoderma viride @ 4g/kg seeds. The seed treatment must be done 24 hours before sowing of seeds.

- Due to forecast of rain, Chances of **fruit rot** infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever **sucking pests** incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. To prevent of soil borne diseases like seedling rot, stem rot and wilt diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Incidence of **thrips** may observe in chilli, if observe spray Spinosad 45% SC @ 0.4 ml/litre of water or Imidacloprid 17.8% SL @ 0.3 ml/litre of water in clean and dry weather. The present weather conditions are favourable for Upward and downward leafcurl in chilli. To control spray phosalon (Zolone) 3 ml. or diphenhiuron @ 1g./L of water.
- To control **fruit and shoot borer** in okra, Spray Bacillus thuringiensis 5% W.P @ 1.5 gram/lit of water. Present weather condition favours incidence of white fly and jassid. To control this apply Thiomethoxam @ 0.4g/ litre of water.

Fruits:

Spray mancozeb 2g/litre, to control sigatoka leaf spot in **banana**. To control **papaya yellow mosaic virus**, spray Imidachloprid @ 0.5 ml / litre.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-314
Date- 19/07/2022

AGROMET ADVISORY BULLETIN FOR BANKI

District Past week average weather condition

The district received average 21mm rainfall during the last week. The mean daily maximum and minimum temperature was 32°C (-0.7°C from normal) and 25°C (+1.1°C from normal), respectively. Mean relative humidity was around 88-95%.

Forecasted period: Today(8.30IST) to 24th July(8.30IST)2022

Weather advisory: Moderate with isolated heavy rainfall is forecasted for Friday and Saturday and very light to light rainfall is forecasted for other 3 days in the block. The day temperature will be 31-33°C and night temperature will be 25-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 82-84% and 49-55% respectively. **There will be partly cloudy to overcast sky for next 5 days.**

Block	Banki (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	19/07/2022	20/07/2022	21/07/2022	22/07/2022	23/07/2022
Rainfall(mm)	3	7	10	28	32
T-Max(°C)	32	31	32	33	33
T-Min(°C)	25	25	25	26	25
Cloud Cover(octa)	6	6	8	8	8
RH Max (%)	82	82	84	83	84
RH Min (%)	52	49	55	54	51
Wind Speed (Kmph)	14	13	11	8	10
Wind Direction (deg)	225	212	203	207	210

Agromet Advisory

General advisory: Drain out excess water from rice nurseries, orchards, maize and jute fields. Postpone irrigation, intercultural operations, application of plant protection measures and fertilizers to the standing crops from Thursday to Saturday and they may carry out weeding, pesticide and fertilizer application in upcoming 2 days. Keep the harvested produce at safer places. Cover the vegetable seedling with polythene to protect them from heavy shower.

Kharifrice:(DIRECT SEEDED RICE):

Direct seeded rice are at three to four week stage. As light to moderate amount of rainfall is forecasted in the upcoming days, farmers should go for beushanning. Apply 35 kg urea/ac at beushanning followed by weeding and gap filling. Drain out the excess water at the time of fertilizer application and pond it after one day. If basal fertilizer not applied, apply 18 kg urea, 100kg ssp and 26 kg MOP while beushanning followed by top dressing of 35 kg urea after 7-10 days.

TRANSPLANTED RICE:

Considering the anticipated rainfall, paddy transplanting may be completed before 30th July. Apply 400-gram Chlorantraniliprole 0.4 % G or 800-gram Cartap Hydrochloride 4% GR in the 10 decimal nursery area 7 days before transplanting to manage gall midge, stem borer, caseworm, leaf folder and root knot nematode up to 3 weeks after transplanting. Intermittent dry and wet weather condition, is favourable for incidence of **BLB** and **Sheath blight** in early sown nursery seedlings. Go for foliar spray of Hexaconazole with Plantomycin@1.0 g/l to prevent the spread of the disease. Prevailing high temperature condition may lead to infestation of **rice thrips** in rice nursery. Apply Fipronil 5SC@2ml/l of water or Thiomethoxam@0.4g/ litre of water.

Maize:

Stem fly in Maize infects the plants from sowing to 5 weeks after sowing, temperature between 25-30 deg. celsius and humidity above 60% favour pest incidence. In vegetative stage to control stem fly spray Methyl demeton @1 ml/l or Dimethoate @ 2ml/l or apply Carbofuran 3G granuals @ 14 kg/acre. High humid conditions, temperature between 18-27 Deg. Celsius favours the occurrence of **leaf blight** in Maize. To control spray Azoxystrobin + Mancozeb (Pluton) @ 300 g /ac.

ARHAR: In arhar crop, for early variety apply 35 kg DAP, 4 kg Urea and 14 kg MOP as basal fertilizer per acre and for late variety apply 52 kg DAP, 6 kg Urea and 14 kg MOP as basal fertilizer per acre in dry days.

VEGETABLES:

June-july is the best time for sowing of tomato in raised bed for nursery seedlings. Do seed treatment with Trichoderma viride @ 4g/kg seeds. The seed treatment must be done 24 hours before sowing of seeds.

- Due to forecast of rain, Chances of **fruit rot** infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever **sucking pests** incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. **To prevent of soil borne diseases like seedling rot, stem rot and wilt diseases apply 3g of Copper oxychloride per litre of water around the main stem.**
- Incidence of **thrips** may observe in chilli, if observe spray Spinosad 45% SC @ 0.4 ml/litre of water or Imidacloprid 17.8% SL @ 0.3 ml/litre of water in clean and dry weather. The present weather conditions are favourable for Upward and downward leafcurl in chilli. To control spray phosalon (Zolone) 3 ml. or diphenthiuron @ 1g./L of water.
- To control **fruit and shoot borer** in okra, Spray Bacillus thuringiensis 5% W.P @1.5 gram/lit of water. Present weather condition favours incidence of white fly and jassid. To control this apply Thiomethoxam@0.4g/ litre of water.

Fruits:

Spray mancozeb 2g/litre, to control sigatoka leaf spot in **banana**. To control **papaya yellow mosaic virus**, spray Imidachloprid @ 0.5 ml / litre.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-314
Date- 19/07/2022

AGROMET ADVISORY BULLETIN FOR BANKI-DAMPARA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 21mm and district received average 21mm rainfall during the last week. The mean daily maximum and minimum temperature was 32°C (-0.7°C from normal) and 25°C (+1.1°C from normal), respectively. Mean relative humidity was around 88-95%.

Forecasted period: Today(8.30IST) to 24th July(8.30IST)2022

Weather Advisory:: Moderate with Isolated heavy rainfall is forecasted for Friday and Saturday and very light to light rainfall is forecasted for other 3 days in the block. The day temperature will be 31-33°C and night temperature will be 25-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 82-84% and 49-55% respectively. **There will be partly cloudy to overcast sky for next 5 days.**

Block	Banki-Dampara (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	19/07/2022	20/07/2022	21/07/2022	22/07/2022	23/07/2022
Rainfall(mm)	4	9	10	29	35
T-Max(°C)	32	31	32	33	33
T-Min(°C)	25	25	25	26	25
Cloud Cover(octa)	6	6	8	8	8
RH Max (%)	82	82	84	83	84
RH Min (%)	52	49	55	54	52
Wind Speed (Kmph)	15	13	11	8	10
Wind Direction (deg)	225	212	203	207	210

Agromet Advisory

General advisory: Drain out excess water from rice nurseries, orchards, maize and jute fields. Postpone irrigation, intercultural operations, application of plant protection measures and fertilizers to the standing crops from Thursday to Saturday and they may carry out weeding, pesticide and fertilizer application in upcoming 2 days. Keep the harvested produce at safer places. Cover the vegetable seedling with polythene to protect them from heavy shower.

Kharifrice:(DIRECT SEEDED RICE):

Direct seeded rice are at three to four week stage. As light to moderate amount of rainfall is forecasted in the upcoming days, farmers should go for beushanning. Apply 35 kg urea/ac at beushanning followed by weeding and gap filling. Drain out the excess water at the time of fertilizer application and pond it after one day. If basal fertilizer not applied, apply 18 kg urea, 100kg ssp and 26 kg MOP while beushanning followed by top dressing of 35 kg urea after 7-10 days.

TRANSPLANTED RICE:

Considering the anticipated rainfall, paddy transplanting may be completed before 30th July. Apply 400-gram Chlorantraniliprole 0.4 % G or 800-gram Cartap Hydrochloride 4% GR in the 10 decimal nursery area 7 days before transplanting to manage gall midge, stem borer, caseworm, leaf folder and root knot nematode up to 3 weeks after transplanting. Intermittent dry and wet weather condition, is favourable for incidence of **BLB** and **Sheath blight** in early sown nursery seedlings. Go for foliar spray of Hexaconazole with Plantomycin@1.0 g/l to prevent the spread of the disease. Prevailing high temperature condition may lead to infestation of **rice thrips** in rice nursery. Apply Fipronil 5SC@2ml/l of water or Thiomethoxam@0.4g/ litre of water.

Maize:

Stem fly in Maize infects the plants from sowing to 5 weeks after sowing, temperature between 25-30 deg. celsius and humidity above 60% favour pest incidence. In vegetative stage to control stem fly spray Methyl demeton @1 ml/l or Dimethoate @ 2ml/l or apply Carbofuran 3G granuals @ 14 kg/acre. High humid conditions, temperature between 18-27 Deg. Celsius favours the occurrence of **leaf blight** in Maize. To control spray Azoxystrobin + Mancozeb (Pluton) @ 300 g /ac.

ARHAR: In arhar crop, for early variety apply 35 kg DAP, 4 kg Urea and 14 kg MOP as basal fertilizer per acre and for late variety apply 52 kg DAP, 6 kg Urea and 14 kg MOP as basal fertilizer per acre in dry days.

VEGETABLES:

June-july is the best time for sowing of tomato in raised bed for nursery seedlings. Do seed treatment with Trichoderma viride @ 4g/kg seeds. The seed treatment must be done 24 hours before sowing of seeds.

- Due to forecast of rain, Chances of **fruit rot** infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever **sucking pests** incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. To prevent of soil borne diseases like seedling rot, stem rot and wilt diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Incidence of **thrips** may observe in chilli, if observe spray Spinosad 45% SC @ 0.4 ml/litre of water or Imidacloprid 17.8% SL @ 0.3 ml/litre of water in clean and dry weather. The present weather conditions are favourable for Upward and downward leafcurl in chilli. To control spray phosalon (Zolone) 3 ml. or diphenhiuron @ 1g./L of water.
- To control **fruit and shoot borer** in okra, Spray Bacillus thuringiensis 5% W.P @1.5 gram/lit of water. Present weather condition favours incidence of white fly and jassid. To control this apply Thiomethoxam@0.4g/ litre of water.

Fruits:

Spray mancozeb 2g/litre, to control sigatoka leaf spot in **banana**. To control **papaya yellow mosaic virus**, spray Imidachloprid @ 0.5 ml / litre.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-314
Date- 19/07/2022

AGROMET ADVISORY BULLETIN FOR BARANGA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 18mm and district received average 21mm rainfall during the last week. The mean daily maximum and minimum temperature was 32°C (-0.7°C from normal) and 25°C (+1.1°C from normal), respectively. Mean relative humidity was around 88-95%.

Forecasted period: Today(8.30IST) to 24th July(8.30IST)2022

Weather advisory: *Moderate with Isolated heavy rainfall is forecasted for Friday and Saturday and very light to light rainfall is forecasted for other 3 days in the block.* The day temperature will be 32-34°C and night temperature will be 26°C for upcoming 5 days. Maximum and minimum relative humidity will be 78-81% and 46-54% respectively. **There will be partly cloudy to overcast sky for next 5 days.**

Block	Baranga (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	19/07/2022	20/07/2022	21/07/2022	22/07/2022	23/07/2022
Rainfall(mm)	5	4	10	28	32
T-Max(°C)	34	32	33	33	33
T-Min(°C)	26	26	26	26	26
Cloud Cover	7	6	8	8	8
RH Max (%)	78	79	79	81	81
RH Min (%)	48	46	49	54	54
Wind Speed (Kmph)	18	15	13	10	11
Wind Direction (deg)	225	207	205	203	205

Agromet Advisory

General advisory: *Drain out excess water from rice nurseries, orchards, maize and jute fields. Postpone irrigation, intercultural operations, application of plant protection measures and fertilizers to the standing crops from Thursday to Saturday and they may carry out weeding, pesticide and fertilizer application in upcoming 2 days. Keep the harvested produce at safer places. Cover the vegetable seedling with polythene to protect them from heavy shower.*

Kharifrice:(DIRECT SEEDED RICE):

Direct seeded rice are at three to four week stage. As light to moderate amount of rainfall is forecasted in the upcoming days, farmers should go for beushanning. Apply 35 kg urea/ac at beushanning followed by weeding and gap filling. Drain out the excess water at the time of fertilizer application and pond it after one day. If basal fertilizer not applied, apply 18 kg urea, 100kg ssp and 26 kg MOP while beushanning followed by top dressing of 35 kg urea after 7-10 days.

TRANSPLANTED RICE:

Considering the anticipated rainfall, paddy transplanting may be completed before 30th July. Apply 400-gram Chlorantraniliprole 0.4 % G or 800-gram Cartap Hydrochloride 4% GR in the 10 decimal nursery area 7 days before transplanting to manage gall midge, stem borer, caseworm, leaf folder and root knot nematode up to 3 weeks after transplanting. Intermittent dry and wet weather condition, is favourable for incidence of **BLB** and **Sheath blight** in early sown nursery seedlings. Go for foliar spray of Hexaconazole with Plantomycin@1.0 g/l to prevent the spread of the disease. Prevailing high temperature condition may lead to infestation of **rice thrips** in rice nursery. Apply Fipronil 5SC@2ml/l of water or Thiomethoxam@0.4g/ litre of water.

Maize:

Stem fly in Maize infects the plants from sowing to 5 weeks after sowing, temperature between 25-30 deg. celsius and humidity above 60% favour pest incidence. In vegetative stage to control stem fly spray Methyl demeton @1 ml/lit or Dimethoate @ 2ml/lit or apply Carbofuran 3G granuals @ 14 kg/acre. High humid conditions, temperature between 18-27 Deg. Celsius favours the occurrence of **leaf blight** in Maize. To control spray Azoxystrobin + Mancozeb (Pluton) @ 300 g /ac.

ARHAR: In arhar crop, for early variety apply 35 kg DAP, 4 kg Urea and 14 kg MOP as basal fertilizer per acre and for late variety apply 52 kg DAP, 6 kg Urea and 14 kg MOP as basal fertilizer per acre in dry days.

VEGETABLES:

June-july is the best time for sowing of tomato in raised bed for nursery seedlings. Do seed treatment with Trichoderma viride @ 4g/kg seeds. The seed treatment must be done 24 hours before sowing of seeds.

- Due to forecast of rain, Chances of **fruit rot** infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever **sucking pests** incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. **To prevent of soil borne diseases like seedling rot, stem rot and wilt diseases apply 3g of Copper oxychloride per litre of water around the main stem.**
- Incidence of **thrips** may observe in chilli, if observe spray Spinosad 45% SC @ 0.4 ml/litre of water or Imidacloprid 17.8% SL @ 0.3 ml/litre of water in clean and dry weather. The present weather conditions are favourable for Upward and downward leafcurl in chilli. To control spray phosalon (Zolone) 3 ml. or diphenthiuron @ 1g./L of water.
- To control **fruit and shoot borer** in okra, Spray Bacillus thuringiensis 5% W.P @1.5 gram/lit of water. Present weather condition favours incidence of white fly and jassid. To control this apply Thiomethoxam@0.4g/ litre of water.

Fruits: Spray mancozeb 2g/litre, to control sigatoka leaf spot in banana. To control papaya yellow mosaic virus, spray Imidachloprid @ 0.5 ml / litre.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-314
Date- 19/07/2022

AGROMET ADVISORY BULLETIN FOR CUTTACK SADAR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 19mm and district received average 21mm rainfall during the last week. The mean daily maximum and minimum temperature was 32°C (-0.7°C from normal) and 25°C (+1.1°C from normal), respectively. Mean relative humidity was around 88-95%.

Forecasted period: Today(8.30IST) to 24th July (8.30IST)2022

Weather advisory: Moderate with Isolated heavy rainfall is forecasted for Friday and Saturday and very light to light rainfall is forecasted for other 3 days in the block. The day temperature will be 32-33°C and night temperature will be 26°C for upcoming 5 days. Maximum and minimum relative humidity will be 78-83% and 47-57% respectively. **There will be partly cloudy to overcast sky for next 5 days.**

Block	Cuttack sadar (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	19/07/2022	20/07/2022	21/07/2022	22/07/2022	23/07/2022
Rainfall(mm)	5	4	7	28	32
T-Max(°C)	33	32	33	32	33
T-Min(°C)	26	26	26	26	26
Cloud Cover(octa)	6	6	8	8	8
RH Max (%)	78	80	81	82	83
RH Min (%)	48	47	50	55	57
Wind Speed (Kmph)	17	14	12	10	10
Wind Direction (deg)	225	212	205	200	199

Agromet Advisory

General advisory: Drain out excess water from rice nurseries, orchards, maize and jute fields. Postpone irrigation, intercultural operations, application of plant protection measures and fertilizers to the standing crops from Thursday to Saturday and they may carry out weeding, pesticide and fertilizer application in upcoming 2 days. Keep the harvested produce at safer places. Cover the vegetable seedling with polythene to protect them from heavy shower.

Kharifrice:(DIRECT SEEDED RICE):

Direct seeded rice are at three to four week stage. As light to moderate amount of rainfall is forecasted in the upcoming days, farmers should go for beushanning. Apply 35 kg urea/ac at beushanning followed by weeding and gap filling. Drain out the excess water at the time of fertilizer application and pond it after one day. If basal fertilizer not applied, apply 18 kg urea, 100kg ssp and 26 kg MOP while beushanning followed by top dressing of 35 kg urea after 7-10 days.

TRANSPLANTED RICE:

Considering the anticipated rainfall, paddy transplanting may be completed before 30th July. Apply 400-gram Chlorantraniliprole 0.4 % G or 800-gram Cartap Hydrochloride 4% GR in the 10 decimal nursery area 7 days before transplanting to manage gall midge, stem borer, caseworm, leaf folder and root knot nematode up to 3 weeks after transplanting. Intermittent dry and wet weather condition, is favourable for incidence of **BLB** and **Sheath blight** in early sown nursery seedlings. Go for foliar spray of Hexaconazole with Plantomycin@1.0 g/l to prevent the spread of the disease. Prevailing high temperature condition may lead to infestation of **rice thrips** in rice nursery. Apply Fipronil 5SC@2ml/l of water or Thiomethoxam@0.4g/ litre of water.

Sugarcane: For **Red rot and wilt disease**, spray the crop with Mancozeb 0.3% or carbendazim 0.1% or Azoxystrobin 18.2%SC/ Difenconazole 11.45SC @ 1ml per liter water in dry days. weather is conducive for temperatures between 29-35 degree Celsius, humidity 60-70 %; this may favour the growth of Whiteflies in sugarcane;; spray Thiamethoxam 25 WG @ 125g/ha or apply Chloropyrifos @ 2.5ml/litre water. Present rainy weather favors infestation Wilt disease in Sugarcane, to control this drain out excess water and apply Cupper oxychloride 50.0% 3 gram per liter of water.

ARHAR :

In arhar crop, for early variety apply 35 kg DAP, 4 kg Urea and 14 kg MOP as basal fertilizer per acre and for late variety apply 52 kg DAP, 6 kg Urea and 14 kg MOP as basal fertilizer per acre in dry days.

VEGETABLES:

June-july is the best time for sowing of tomato in raised bed for nursery seedlings. Do seed treatment with Trichoderma viride @ 4g/kg seeds. The seed treatment must be done 24 hours before sowing of seeds.

- Due to forecast of rain, Chances of **fruit rot** infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever **sucking pests** incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. **To prevent of soil borne diseases like seedling rot, stem rot and wilt diseases apply 3g of Copper oxychloride per litre of water around the main stem.**
- Incidence of **thrips** may observe in chilli, if observe spray Spinosad 45% SC @ 0.4 ml/litre of water or Imidacloprid 17.8% SL @ 0.3 ml/litre of water in clean and dry weather. The present weather conditions are favourable for Upward and downward leafcurl in chilli. To control spray phosalon (Zolone) 3 ml. or diphenthiuron @ 1g./L of water.
- To control **fruit and shoot borer** in okra, Spray Bacillus thuringiensis 5% W.P @1.5 gram/lit of water. Present weather condition favours incidence of white fly and jassid. To control this apply Thiomethoxam@0.4g/ litre of water.

Fruits: Spray mancozeb 2g/litre, to control sigatoka leaf spot in **banana**. To control **papaya yellow mosaic virus**, spray Imidachloprid @ 0.5 ml / litre.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-314
Date- 19/07/2022

AGROMET ADVISORY BULLETIN FOR KANTAPADA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 28mm and district received average 21mm rainfall during the last week. The mean daily maximum and minimum temperature was 32°C (-0.7°C from normal) and 25°C (+1.1°C from normal), respectively. Mean relative humidity was around 88-95%.

Forecasted period: Today(8.30IST) to 24th July(8.30IST)2022

Weather advisory: Moderate with Isolated heavy rainfall is forecasted for Friday and Saturday and very light to light rainfall is forecasted for other 3 days in the block. The day temperature will be 33-34°C and night temperature will be 26-27°C for upcoming 5 days. Maximum and minimum relative humidity will be 78-80% and 46-53% respectively. **There will be partly cloudy to overcast sky for next 5 days.**

Block	Kantapada (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	19/07/2022	20/07/2022	21/07/2022	22/07/2022	23/07/2022
Rainfall(mm)	2	4	6	25	31
T-Max(°C)	34	33	33	33	33
T-Min(°C)	26	26	27	27	27
Cloud Cover(octa)	7	6	8	8	8
RH Max (%)	78	78	78	80	80
RH Min (%)	48	46	47	52	53
Wind Speed (Kmph)	18	16	13	11	12
Wind Direction (deg)	225	204	207	203	205

Agromet Advisory

General advisory: Drain out excess water from rice nurseries, orchards, maize and jute fields. Postpone irrigation, intercultural operations, application of plant protection measures and fertilizers to the standing crops from Thursday to Saturday and they may carry out weeding, pesticide and fertilizer application in upcoming 2 days. Keep the harvested produce at safer places. Cover the vegetable seedling with polythene to protect them from heavy shower.

Khari rice: (DIRECT SEEDED RICE):

Direct seeded rice are at three to four week stage. As light to moderate amount of rainfall is forecasted in the upcoming days, farmers should go for beushanning. Apply 35 kg urea/ac at beushanning followed by weeding and gap filling. Drain out the excess water at the time of fertilizer application and pond it after one day. If basal fertilizer not applied, apply 18 kg urea, 100kg ssp and 26 kg MOP while beushanning followed by top dressing of 35 kg urea after 7-10 days.

TRANSPLANTED RICE:

Considering the anticipated rainfall, paddy transplanting may be completed before 30th July. Apply 400-gram Chlorantraniliprole 0.4 % G or 800-gram Cartap Hydrochloride 4% GR in the 10 decimal nursery area 7 days before transplanting to manage gall midge, stem borer, caseworm, leaf folder and root knot nematode up to 3 weeks after transplanting. Intermittent dry and wet weather condition, is favourable for incidence of **BLB** and **Sheath blight** in early sown nursery seedlings. Go for foliar spray of Hexaconazole with Plantomycin @ 1.0 g/l to prevent the spread of the disease. Prevailing high temperature condition may lead to infestation of **rice thrips** in rice nursery. Apply Fipronil 5SC @ 2ml/l of water or Thiomethoxam @ 0.4g/ litre of water.

Sugarcane:

For **Red rot and wilt disease**, spray the crop with Mancozeb 0.3% or carbendazim 0.1% or Azoxystrobin 18.2% SC/ Difenconazole 11.45SC @ 1ml per liter water in dry days. weather is conducive for temperatures between 29-35 degree Celsius, humidity 60-70 %; this may favour the growth of Whiteflies in sugarcane;; spray Thiamethoxam 25 WG @ 125g/ha or apply Chloropyriphos @ 2.5ml/litre water. Present rainy weather favors infestation Wilt disease in Sugarcane, to control this drain out excess water and apply Copper oxychloride 50.0% 3 gram per liter of water.

ARHAR :

In arhar crop, for early variety apply 35 kg DAP, 4 kg Urea and 14 kg MOP as basal fertilizer per acre and for late variety apply 52 kg DAP, 6 kg Urea and 14 kg MOP as basal fertilizer per acre in dry days.

VEGETABLES:

June-july is the best time for sowing of tomato in raised bed for nursery seedlings. Do seed treatment with Trichoderma viride @ 4g/kg seeds. The seed treatment must be done 24 hours before sowing of seeds.

- Due to forecast of rain, Chances of **fruit rot** infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever **sucking pests** incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. **To prevent of soil borne diseases like seedling rot, stem rot and wilt diseases apply 3g of Copper oxychloride per litre of water around the main stem.**
- Incidence of **thrips** may observe in chilli, if observe spray Spinosad 45% SC @ 0.4 ml/litre of water or Imidacloprid 17.8% SL @ 0.3 ml/litre of water in clean and dry weather. The present weather conditions are favourable for Upward and downward leafcurl in chilli. To control spray phosalon (Zolone) 3 ml. or diphenthiuron @ 1g./L of water.
- To control **fruit and shoot borer** in okra, Spray Bacillus thuringiensis 5% W.P @ 1.5 gram/lit of water. Present weather condition favours incidence of white fly and jassid. To control this apply Thiomethoxam @ 0.4g/ litre of water.

Fruits: Spray mancozeb 2g/litre, to control sigatoka leaf spot in **banana**. To control **papaya yellow mosaic virus**, spray Imidachloprid @ 0.5 ml / litre.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-314
Date- 19/07/2022

AGROMET ADVISORY BULLETIN FOR MAHANGA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 31mm and district received average 21mm rainfall during the last week. The mean daily maximum and minimum temperature was 32°C (-0.7°C from normal) and 25°C (+1.1°C from normal), respectively. Mean relative humidity was around 88-95%.

Forecasted period: Today(8.30IST) to 24th July(8.30IST)2022

Weather advisory: Moderate with Isolated heavy rainfall is forecasted for Friday and Saturday and light rainfall is forecasted for other 3 days in the block. The day temperature will be and it will be 32-34°C and night temperature will be 26°C for upcoming 5 days. Maximum and minimum relative humidity will be 79-83% and 45-56% respectively. **There will be partly cloudy to overcast sky for next 5 days.**

Block	Mahanga (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	19/07/2022	20/07/2022	21/07/2022	22/07/2022	23/07/2022
Rainfall(mm)	5	7	9	35	31
T-Max(°C)	34	33	33	33	32
T-Min(°C)	26	26	26	26	26
Cloud Cover(octal)	7	6	8	8	8
RH Max (%)	79	80	82	82	83
RH Min (%)	48	45	49	51	56
Wind Speed (Kmph)	18	15	13	10	11
Wind Direction (deg)	225	207	200	203	210

Agromet Advisory

General advisory: Drain out excess water from rice nurseries, orchards, maize and jute fields. Postpone irrigation, intercultural operations, application of plant protection measures and fertilizers to the standing crops from Thursday to Saturday and they may carry out weeding, pesticide and fertilizer application in upcoming 2 days. Keep the harvested produce at safer places. Cover the vegetable seedling with polythene to protect them from heavy shower.

Khari rice: (DIRECT SEEDED RICE):

Direct seeded rice are at three to four week stage. As light to moderate amount of rainfall is forecasted in the upcoming days, farmers should go for beushanning. Apply 35 kg urea/ac at beushanning followed by weeding and gap filling. Drain out the excess water at the time of fertilizer application and pond it after one day. If basal fertilizer not applied, apply 18 kg urea, 100kg ssp and 26 kg MOP while beushanning followed by top dressing of 35 kg urea after 7-10 days.

TRANSPLANTED RICE:

Considering the anticipated rainfall, paddy transplanting may be completed before 30th July. Apply 400-gram Chlorantraniliprole 0.4 % G or 800-gram Cartap Hydrochloride 4% GR in the 10 decimal nursery area 7 days before transplanting to manage gall midge, stem borer, caseworm, leaf folder and root knot nematode up to 3 weeks after transplanting. Intermittent dry and wet weather condition, is favourable for incidence of **BLB** and **Sheath blight** in early sown nursery seedlings. Go for foliar spray of Hexaconazole with Plantomycin @ 1.0 g/l to prevent the spread of the disease. Prevailing high temperature condition may lead to infestation of **rice thrips** in rice nursery. Apply Fipronil 5SC @ 2ml/l of water or Thiomethoxam @ 0.4g/ litre of water.

Sugarcane:

For **Red rot and wilt disease**, spray the crop with Mancozeb 0.3% or carbendazim 0.1% or Azoxystrobin 18.2% SC/ Difenconazole 11.45SC @ 1ml per liter water in dry days. weather is conducive for temperatures between 29-35 degree Celsius, humidity 60-70 %; this may favour the growth of Whiteflies in sugarcane; spray Thiamethoxam 25 WG @ 125g/ha or apply Chloropyrifos @ 2.5ml/litre water. Present rainy weather favors infestation Wilt disease in Sugarcane, to control this drain out excess water and apply Copper oxychloride 50.0% 3 gram per liter of water.

Jute: Recent rain has aggravated **weed problem** in early sown crop. To control grass weeds, apply Quizalofop-pethyl @ 2ml/litre of water at 2-3 leaf stage of weeds. After rain in vegetative stage to control **stem rot of jute**, spray copper oxychloride @ 3-4 gram/litre of water. To control **indigo caterpillar** apply Spinosad (45% S.C) 0.3 ml/litre of water or Emamectin benzoate (5% SC) 0.5 gram per liter of water.

VEGETABLES:

June-july is the best time for sowing of tomato in raised bed for nursery seedlings. Do seed treatment with Trichoderma viride @ 4g/kg seeds. The seed treatment must be done 24 hours before sowing of seeds.

- Due to forecast of rain, Chances of **fruit rot** infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever **sucking pests** incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. To prevent of soil borne diseases like seedling rot, stem rot and wilt diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Incidence of **thrips** may observe in chilli, if observe spray Spinosad 45% SC @ 0.4 ml/litre of water or Imidacloprid 17.8% SL @ 0.3 ml/litre of water in clean and dry weather. The present weather conditions are favourable for Upward and downward leafcurl in chilli. To control spray phosalon (Zolone) 3 ml. or diphenhiuron @ 1g./L of water.
- To control **fruit and shoot borer** in okra, Spray Bacillus thuringiensis 5% W.P @ 1.5 gram/lit of water. Present weather condition favours incidence of white fly and jassid. To control this apply Thiomethoxam @ 0.4g/ litre of water.

Fruits: Spray mancozeb 2g/litre, to control sigatoka leaf spot in **banana**. To control **papaya yellow mosaic virus**, spray Imidachloprid @ 0.5 ml / litre.

Dr Sujata Sethy
 (Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-314
Date- 19/07/2022

AGROMET ADVISORY BULLETIN FOR NARASINGHPUR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 26mm and district received average 21mm rainfall during the last week. The mean daily maximum and minimum temperature was 32°C (-0.7°C from normal) and 25°C (+1.1°C from normal), respectively. Mean relative humidity was around 88-95%.

Forecasted period: Today(8.30IST) to 24th July(8.30IST)2022

Weather advisory: Moderate with isolated heavy rainfall is forecasted for Friday and Saturday and light to moderate rainfall is forecasted for other 3 days in the block. The day temperature will be 31-32°C and night temperature will be 24-25°C for upcoming 5 days. Maximum and minimum relative humidity will be 82-89% and 53-65% respectively. **There will be partly cloudy to overcast sky for next 5 days.**

Block	Narasinghpur (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	19/07/2022	20/07/2022	21/07/2022	22/07/2022	23/07/2022
Rainfall(mm)	2	9	19	33	35
T-Max(°C)	32	31	31	31	32
T-Min(°C)	24	24	25	25	24
Cloud Cover(/octa)	6	6	8	8	8
RH Max (%)	82	85	88	86	89
RH Min (%)	55	53	62	65	57
Wind Speed (Kmph)	9	7	7	5	6
Wind Direction (deg)	242	234	205	240	270

Agromet Advisory

General advisory: Drain out excess water from rice nurseries, orchards, maize and jute fields. Postpone irrigation, intercultural operations, application of plant protection measures and fertilizers to the standing crops from Thursday to Saturday and they may carry out weeding, pesticide and fertilizer application in upcoming 2 days. Keep the harvested produce at safer places. Cover the vegetable seedling with polythene to protect them from heavy shower.

Kharrifrice:(DIRECT SEEDED RICE):

Direct seeded rice are at three to four week stage. As light to moderate amount of rainfall is forecasted in the upcoming days, farmers should go for beushanning. Apply 35 kg urea/ac at beushanning followed by weeding and gap filling. Drain out the excess water at the time of fertilizer application and pond it after one day. If basal fertilizer not applied, apply 18 kg urea, 100kg ssp and 26 kg MOP while beushanning followed by top dressing of 35 kg urea after 7-10 days.

TRANSPLANTED RICE:

Considering the anticipated rainfall, paddy transplanting may be completed before 30th July. Apply 400-gram Chlorantraniliprole 0.4 % G or 800-gram Cartap Hydrochloride 4% GR in the 10 decimal nursery area 7 days before transplanting to manage gall midge, stem borer, caseworm, leaf folder and root knot nematode up to 3 weeks after transplanting. Intermittent dry and wet weather condition, is favourable for incidence of **BLB** and **Sheath blight** in early sown nursery seedlings. Go for foliar spray of Hexaconazole with Plantomycin@1.0 g/l to prevent the spread of the disease. Prevailing high temperature condition may lead to infestation of **rice thrips** in rice nursery. Apply Fipronil 5SC@2ml/l of water or Thiomethoxam@0.4g/ litre of water.

Maize:

Stem fly in Maize infects the plants from sowing to 5 weeks after sowing, temperature between 25-30 deg. celsius and humidity above 60% favour pest incidence. In vegetative stage to control stem fly spray Methyl demeton @1 ml/l or Dimethoate @ 2ml/l or apply Carbofuran 3G granuals @ 14 kg/acre. High humid conditions, temperature between 18-27 Deg. Celsius favours the occurrence of **leaf blight** in Maize. To control spray Azoxystrobin + Mancozeb (Pluton) @ 300 g /ac.

ARHAR: In arhar crop, for early variety apply 35 kg DAP, 4 kg Urea and 14 kg MOP as basal fertilizer per acre and for late variety apply 52 kg DAP, 6 kg Urea and 14 kg MOP as basal fertilizer per acre in dry days.

VEGETABLES:

June-july is the best time for sowing of tomato in raised bed for nursery seedlings. Do seed treatment with Trichoderma viride @ 4g/kg seeds. The seed treatment must be done 24 hours before sowing of seeds.

- Due to forecast of rain, Chances of **fruit rot** infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever **sucking pests** incidence was observed spray 5 ml Neem oil or Thiomethoxam 0.2 g. per litre of water. **To prevent of soil borne diseases like seedling rot, stem rot and wilt diseases apply 3g of Copper oxychloride per litre of water around the main stem.**
- Incidence of **thrips** may observe in chilli, if observe spray Spinosad 45% SC @ 0.4 ml/litre of water or Imidacloprid 17.8% SL @ 0.3 ml/litre of water in clean and dry weather. The present weather conditions are favourable for Upward and downward leafcurl in chilli. To control spray phosalon (Zolone) 3 ml. or diphenhiuron @ 1g./L of water.
- To control **fruit and shoot borer** in okra, Spray Bacillus thuringiensis 5% W.P @1.5 gram/lit of water. Present weather condition favours incidence of white fly and jassid. To control this apply Thiomethoxam@0.4g/ litre of water.

Fruits:

Spray mancozeb 2g/litre, to control sigatoka leaf spot in **banana**. To control **papaya yellow mosaic virus**, spray Imidachloprid @ 0.5 ml / litre.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-314
Date- 19/07/2022

AGROMET ADVISORY BULLETIN FOR NIALI

District Past week average weather condition and Block past week average Rainfall condition

The block received average 11mm and district received average 21mm rainfall during the last week. The mean daily maximum and minimum temperature was 32°C (-0.7°C from normal) and 25°C (+1.1°C from normal), respectively. Mean relative humidity was around 88-95%.

Forecasted period: Today(8.30IST) to 24th July(8.30IST)2022

Weather advisory: Moderate with isolated heavy rainfall is forecasted for Friday and Saturday and very light to light rainfall is forecasted for other 3 days in the block. The day temperature will be 33-34°C and night temperature will be 26-27°C for upcoming 5 days. Maximum and minimum relative humidity will be 77-79% and 47-52% respectively. **There will be partly cloudy to overcast sky for next 5 days.**

Block	Niali (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	19/07/2022	20/07/2022	21/07/2022	22/07/2022	23/07/2022
Rainfall(mm)	1	3	6	16	31
T-Max(°C)	34	33	33	34	33
T-Min(°C)	26	26	27	27	27
Cloud Cover(octa)	7	5	8	8	8
RH Max (%)	78	77	78	79	79
RH Min (%)	49	47	48	52	51
Wind Speed(Kmph)	19	18	15	13	14
Wind Direction(deg)	212	204	204	203	208

Agromet Advisory

General advisory: Drain out excess water from rice nurseries, orchards, maize and jute fields. Postpone irrigation, intercultural operations, application of plant protection measures and fertilizers to the standing crops from Thursday to Saturday and they may carry out weeding, pesticide and fertilizer application in upcoming 2 days. Keep the harvested produce at safer places. Cover the vegetable seedling with polythene to protect them from heavy shower.

Kharifrice:(DIRECT SEEDED RICE):

Direct seeded rice are at three to four week stage. As light to moderate amount of rainfall is forecasted in the upcoming days, farmers should go for beushanning. Apply 35 kg urea/ac at beushanning followed by weeding and gap filling. Drain out the excess water at the time of fertilizer application and pond it after one day. If basal fertilizer not applied, apply 18 kg urea, 100kg ssp and 26 kg MOP while beushanning followed by top dressing of 35 kg urea after 7-10 days.

TRANSPLANTED RICE:

Considering the anticipated rainfall, paddy transplanting may be completed before 30th July. Apply 400-gram Chlorantraniliprole 0.4 % G or 800-gram Cartap Hydrochloride 4% GR in the 10 decimal nursery area 7 days before transplanting to manage gall midge, stem borer, caseworm, leaf folder and root knot nematode up to 3 weeks after transplanting. Intermittent dry and wet weather condition, is favourable for incidence of **BLB** and **Sheath blight** in early sown nursery seedlings. Go for foliar spray of Hexaconazole with Plantomycin@1.0 g/l to prevent the spread of the disease. Prevailing high temperature condition may lead to infestation of **rice thrips** in rice nursery. Apply Fipronil 5SC@2ml/l of water or Thiomethoxam@0.4g/ litre of water.

Sugarcane:

For **Red rot and wilt disease**, spray the crop with Mancozeb 0.3% or carbendazim 0.1% or Azoxystrobin 18.2%SC/ Difenconazole 11.45SC @ 1ml per liter water in dry days. weather is conducive for temperatures between 29-35 degree Celsius, humidity 60-70 %; this may favour the growth of Whiteflies in sugarcane;; spray Thiamethoxam 25 WG @ 125g/ha or apply Chloropyriphos @ 2.5ml/litre water. Present rainy weather favors infestation Wilt disease in Sugarcane, to control this drain out excess water and apply Cupper oxychloride 50.0% 3 gram per liter of water.

ARHAR :

In arhar crop, for early variety apply 35 kg DAP, 4 kg Urea and 14 kg MOP as basal fertilizer per acre and for late variety apply 52 kg DAP, 6 kg Urea and 14 kg MOP as basal fertilizer per acre in dry days.

VEGETABLES:

June-july is the best time for sowing of tomato in raised bed for nursery seedlings. Do seed treatment with Trichoderma viride @ 4g/kg seeds. The seed treatment must be done 24 hours before sowing of seeds.

- Due to forecast of rain, Chances of **fruit rot** infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever **sucking pests** incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. **To prevent of soil borne diseases like seedling rot, stem rot and wilt diseases apply 3g of Copper oxychloride per litre of water around the main stem.**
- Incidence of **thrips** may observe in chilli, if observe spray Spinosad 45% SC @ 0.4 ml/litre of water or Imidacloprid 17.8% SL @ 0.3 ml/litre of water in clean and dry weather. The present weather conditions are favourable for Upward and downward leafcurl in chilli. To control spray phosalon (Zolone) 3 ml. or diphenhiuron @ 1g./L of water.
- To control **fruit and shoot borer** in okra, Spray Bacillus thuringiensis 5% W.P @1.5 gram/lit of water. Present weather condition favours incidence of white fly and jassid. To control this apply Thiomethoxam@0.4g/ litre of water.

Fruits: Spray mancozeb 2g/litre, to control sigatoka leaf spot in **banana**. To control **papaya yellow mosaic virus**, spray Imidachloprid @ 0.5 ml / litre.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightning alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-314
Date- 19/07/2022

AGROMET ADVISORY BULLETIN FOR NISCHINTAKOILI
District Past week average weather condition and Block past week average Rainfall condition

The block received average 25mm and district received average 21mm rainfall during the last week. The mean daily maximum and minimum temperature was 32°C (-0.7°C from normal) and 25°C (+1.1°C from normal), respectively. Mean relative humidity was around 88-95%.

Forecasted period: Today(8.30IST) to 24th July(8.30IST)2022

Weather advisory: Moderate with Isolated heavy rainfall is forecasted for Friday and Saturday and light rainfall is forecasted for other 3 days in the block. The day temperature will be 32-34°C and night temperature will be 26°C for upcoming 5 days. Maximum and minimum relative humidity will be 79-83% and 45-57% respectively. **There will be partly cloudy to overcast sky for next 5 days.**

Block	Nischintakoili (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	19/07/2022	20/07/2022	21/07/2022	22/07/2022	23/07/2022
Rainfall(mm)	3	6	7	22	33
T-Max(°C)	34	33	33	33	32
T-Min(°C)	26	26	26	26	26
Cloud Cover(octa)	7	6	8	8	8
RH Max (%)	79	80	80	82	83
RH Min (%)	47	45	48	51	57
Wind Speed (Kmph)	18	16	13	11	11
Wind Direction (deg)	225	207	200	203	205

Agromet Advisory

General advisory: Drain out excess water from rice nurseries, orchards, maize and jute fields. Postpone irrigation, intercultural operations, application of plant protection measures and fertilizers to the standing crops from Thursday to Saturday and they may carry out weeding, pesticide and fertilizer application in upcoming 2 days. Keep the harvested produce at safer places. Cover the vegetable seedling with polythene to protect them from heavy shower.

Kharifrice:(DIRECT SEEDED RICE):

Direct seeded rice are at three to four week stage. As light to moderate amount of rainfall is forecasted in the upcoming days, farmers should go for beushanning. Apply 35 kg urea/ac at beushanning followed by weeding and gap filling. Drain out the excess water at the time of fertilizer application and pond it after one day. If basal fertilizer not applied, apply 18 kg urea, 100kg ssp and 26 kg MOP while beushanning followed by top dressing of 35 kg urea after 7-10 days.

TRANSPLANTED RICE:

Considering the anticipated rainfall, paddy transplanting may be completed before 30th July. Apply 400-gram Chlorantraniliprole 0.4 % G or 800-gram Cartap Hydrochloride 4% GR in the 10 decimal nursery area 7 days before transplanting to manage gall midge, stem borer, caseworm, leaf folder and root knot nematode up to 3 weeks after transplanting. Intermittent dry and wet weather condition, is favourable for incidence of **BLB** and **Sheath blight** in early sown nursery seedlings. Go for foliar spray of Hexaconazole with Plantomycin@1.0 g/l to prevent the spread of the disease. Prevailing high temperature condition may lead to infestation of **rice thrips** in rice nursery. Apply Fipronil 5SC@2ml/l of water or Thiomethoxam@0.4g/ litre of water.

Sugarcane:

For **Red rot and wilt disease**, spray the crop with Mancozeb 0.3% or carbendazim 0.1% or Azoxystrobin 18.2%SC/ Difenconazole 11.45SC @ 1ml per liter water in dry days. weather is conducive for temperatures between 29-35 degree Celsius, humidity 60-70 %; this may favour the growth of Whiteflies in sugarcane; spray Thiamethoxam 25 WG @ 125g/ha or apply Chloropyriphos @ 2.5ml/litre water. Present rainy weather favors infestation Wilt disease in Sugarcane, to control this drain out excess water and apply Cupper oxychloride 50.0% 3 g per liter of water.

ARHAR :

In arhar crop, for early variety apply 35 kg DAP, 4 kg Urea and 14 kg MOP as basal fertilizer per acre and for late variety apply 52 kg DAP, 6 kg Urea and 14 kg MOP as basal fertilizer per acre in dry days.

VEGETABLES:

June-july is the best time for sowing of tomato in raised bed for nursery seedlings. Do seed treatment with Trichoderma viride @ 4g/kg seeds. The seed treatment must be done 24 hours before sowing of seeds.

- Due to forecast of rain, Chances of **fruit rot** infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever **sucking pests** incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. **To prevent of soil borne diseases like seedling rot, stem rot and wilt diseases apply 3g of Copper oxychloride per litre of water around the main stem.**
- Incidence of **thrips** may observe in chilli, if observe spray Spinosad 45% SC @ 0.4 ml/litre of water or Imidacloprid 17.8% SL @ 0.3 ml/litre of water in clean and dry weather. The present weather conditions are favourable for Upward and downward leafcurl in chilli. To control spray phosalon (Zolone) 3 ml. or diphenthiuron @ 1g./L of water.
- To control **fruit and shoot borer** in okra, Spray Bacillus thuringiensis 5% W.P @1.5 gram/lit of water. Present weather condition favours incidence of white fly and jassid. To control this apply Thiomethoxam@0.4g/ litre of water.

Fruits: Spray mancozeb 2g/litre, to control sigatoka leaf spot in **banana**. To control **papaya yellow mosaic virus**, spray Imidachloprid @ 0.5 ml / litre.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-314
Date- 19/07/2022

AGROMET ADVISORY BULLETIN FOR SALEPUR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 17mm and district received average 21mm rainfall during the last week. The mean daily maximum and minimum temperature was 32°C (-0.7°C from normal) and 25°C (+1.1°C from normal), respectively. Mean relative humidity was around 88-95%.

Forecasted period: Today(8.30IST) to 24th July(8.30IST)2022

Weather advisory: *Moderate with Isolated heavy rainfall is forecasted for Friday and Saturday and light rainfall is forecasted for other 3 days in the block.* The day temperature will be 32-33°C and night temperature will be 26°C for upcoming 5 days. Maximum and minimum relative humidity will be 79-83% and 45-58% respectively. **There will be partly cloudy to overcast sky for next 5 days.**

Block	Salepur (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	19/07/2022	20/07/2022	21/07/2022	22/07/2022	23/07/2022
Rainfall(mm)	4	6	8	35	32
T-Max(°C)	33	33	33	33	32
T-Min(°C)	26	26	26	26	26
Cloud Cover(octa)	7	6	8	8	8
RH Max (%)	79	80	81	82	83
RH Min (%)	48	45	48	51	58
Wind Speed (Kmph)	18	15	13	10	11
Wind Direction (deg)	225	207	200	203	205

Agromet Advisory

General advisory: *Drain out excess water from rice nurseries, orchards, maize and jute fields. Postpone irrigation, intercultural operations, application of plant protection measures and fertilizers to the standing crops from Thursday to Saturday and they may carry out weeding, pesticide and fertilizer application in upcoming 2 days. Keep the harvested produce at safer places. Cover the vegetable seedling with polythene to protect them from heavy shower.*

Kharifrice:(DIRECT SEEDED RICE):

Direct seeded rice are at three to four week stage. As light to moderate amount of rainfall is forecasted in the upcoming days, farmers should go for beushanning. Apply 35 kg urea/ac at beushanning followed by weeding and gap filling. Drain out the excess water at the time of fertilizer application and pond it after one day. If basal fertilizer not applied, apply 18 kg urea, 100kg ssp and 26 kg MOP while beushanning followed by top dressing of 35 kg urea after 7-10 days.

TRANSPLANTED RICE:

Considering the anticipated rainfall, paddy transplanting may be completed before 30th July. Apply 400-gram Chlorantraniliprole 0.4 % G or 800-gram Cartap Hydrochloride 4% GR in the 10 decimal nursery area 7 days before transplanting to manage gall midge, stem borer, caseworm, leaf folder and root knot nematode up to 3 weeks after transplanting. Intermittent dry and wet weather condition, is favourable for incidence of **BLB** and **Sheath blight** in early sown nursery seedlings. Go for foliar spray of Hexaconazole with Plantomycin@1.0 g/l to prevent the spread of the disease. Prevailing high temperature condition may lead to infestation of **rice thrips** in rice nursery. Apply Fipronil 5SC@2ml/l of water or Thiomethoxam@0.4g/ litre of water.

Suagrcane:

For **Red rot and wilt disease**, spray the crop with Mancozeb 0.3% or carbendazim 0.1% or Azoxystrobin 18.2%SC/ Difenconazole 11.45SC @ 1ml per liter water in dry days. weather is conducive for temperatures between 29-35 degree Celsius, humidity 60-70 %; this may favour the growth of Whiteflies in sugarcane;; spray Thiamethoxam 25 WG @ 125g/ha or apply Chloropyrifos @ 2.5ml/litre water. Present rainy weather favors infestation Wilt disease in Sugarcane, to control this drain out excess water and apply Cupper oxychloride 50.0% 3 gram per liter of water.

Jute: Recent rain has aggravated **weed problem** in early sown crop. To control grass weeds, apply Quizalofop-pethyl@2ml/litre of water at 2-3 leaf stage of weeds. After rain in vegetative stage to control **stem rot of jute**, spray copper oxychloride @ 3-4 gram/litre of water. To control **indigo caterpillar** apply Spinosad (45% S.C) 0.3 ml/litre of water or Emamectin benzoate (5% SC) 0.5 gram per liter of water.

VEGETABLES:

June-july is the best time for sowing of tomato in raised bed for nursery seedlings. Do seed treatment with Trichoderma viride @ 4g/kg seeds. The seed treatment must be done 24 hours before sowing of seeds.

- Due to forecast of rain, Chances of **fruit rot** infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever **sucking pests** incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. **To prevent of soil borne diseases like seedling rot, stem rot and wilt diseases apply 3g of Copper oxychloride per litre of water around the main stem.**
- Incidence of **thrips** may observe in chilli, if observe spray Spinosad 45% SC @ 0.4 ml/litre of water or Imidacloprid 17.8% SL @ 0.3 ml/litre of water in clean and dry weather. The present weather conditions are favourable for Upward and downward leafcurl in chilli. To control spray phosalon (Zolone) 3 ml. or diphenthiuron @ 1g./L of water.
- To control **fruit and shoot borer** in okra, Spray Bacillus thuringiensis 5% W.P @1.5 gram/lit of water. Present weather condition favours incidence of white fly and jassid. To control this apply Thiomethoxam@0.4g/ litre of water.

Fruits: Spray mancozeb 2g/litre, to control sigatoka leaf spot in **banana**. To control **papaya yellow mosaic virus**, spray Imidachloprid @ 0.5 ml / litre.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-314
Date- 19/07/2022

AGROMET ADVISORY BULLETIN FOR TANGI-CHOUDWAR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 26mm and district received average 21mm rainfall during the last week. The mean daily maximum and minimum temperature was 32°C (-0.7°C from normal) and 25°C (+1.1°C from normal), respectively. Mean relative humidity was around 88-95%.

Forecasted period: Today(8.30IST) to 24th July(8.30IST)2022

Weather advisory: Moderate with Isolated heavy rainfall is forecasted for Friday and Saturday and light rainfall is forecasted for other 3 days in the block. The day temperature will be 32-33°C and night temperature will be 26°C for upcoming 5 days. Maximum and minimum relative humidity will be 77-80% and 49-55% respectively. **There will be partly cloudy to overcast sky for next 5 days.**

Block	Tangi-Choudwar (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	19/07/2022	20/07/2022	21/07/2022	22/07/2022	23/07/2022
Rainfall(mm)	5	7	10	35	35
T-Max(°C)	33	33	33	33	32
T-Min(°C)	26	26	26	26	26
Cloud Cover(octa)	7	5	8	7	8
RH Max (%)	77	79	80	80	78
RH Min (%)	52	49	52	55	55
Wind Speed (Kmph)	22	21	19	18	16
Wind Direction (deg)	207	203	201	198	205

Agromet Advisory

General advisory: Drain out excess water from rice nurseries, orchards, maize and jute fields. Postpone irrigation, intercultural operations, application of plant protection measures and fertilizers to the standing crops from Thursday to Saturday and they may carry out weeding, pesticide and fertilizer application in upcoming 2 days. Keep the harvested produce at safer places. Cover the vegetable seedling with polythene to protect them from heavy shower.

Kharrifrice:(DIRECT SEEDED RICE):

Direct seeded rice are at three to four week stage. As light to moderate amount of rainfall is forecasted in the upcoming days, farmers should go for beushanning. Apply 35 kg urea/ac at beushanning followed by weeding and gap filling. Drain out the excess water at the time of fertilizer application and pond it after one day. If basal fertilizer not applied, apply 18 kg urea, 100kg ssp and 26 kg MOP while beushanning followed by top dressing of 35 kg urea after 7-10 days.

TRANSPLANTED RICE:

Considering the anticipated rainfall, paddy transplanting may be completed before 30th July. Apply 400-gram Chlorantraniliprole 0.4 % G or 800-gram Cartap Hydrochloride 4% GR in the 10 decimal nursery area 7 days before transplanting to manage gall midge, stem borer, caseworm, leaf folder and root knot nematode up to 3 weeks after transplanting. Intermittent dry and wet weather condition, is favourable for incidence of **BLB** and **Sheath blight** in early sown nursery seedlings. Go for foliar spray of Hexaconazole with Plantomycin@1.0 g/l to prevent the spread of the disease. Prevailing high temperature condition may lead to infestation of **rice thrips** in rice nursery. Apply Fipronil 5SC@2ml/l of water or Thiomethoxam@0.4g/ litre of water.

Maize:

Stem fly in Maize infects the plants from sowing to 5 weeks after sowing, temperature between 25-30 deg. celsius and humidity above 60% favour pest incidence. In vegetative stage to control stem fly spray Methyl demeton @1 ml/lit or Dimethoate @ 2ml/lit or apply Carbofuran 3G granuals @ 14 kg/acre. High humid conditions, temperature between 18-27 Deg. Celsius favours the occurrence of **leaf blight** in Maize. To control spray Azoxystrobin + Mancozeb (Pluton) @ 300 g /ac.

ARHAR: In arhar crop, for early variety apply 35 kg DAP, 4 kg Urea and 14 kg MOP as basal fertilizer per acre and for late variety apply 52 kg DAP, 6 kg Urea and 14 kg MOP as basal fertilizer per acre in dry days.

VEGETABLES:

June-july is the best time for sowing of tomato in raised bed for nursery seedlings. Do seed treatment with Trichoderma viride @ 4g/kg seeds. The seed treatment must be done 24 hours before sowing of seeds.

- Due to forecast of rain, Chances of **fruit rot** infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever **sucking pests** incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. **To prevent of soil borne diseases like seedling rot, stem rot and wilt diseases apply 3g of Copper oxychloride per litre of water around the main stem.**
- Incidence of **thrips** may observe in chilli, if observe spray Spinosad 45% SC @ 0.4 ml/litre of water or Imidacloprid 17.8% SL @ 0.3 ml/litre of water in clean and dry weather. The present weather conditions are favourable for Upward and downward leafcurl in chilli. To control spray phosalon (Zolone) 3 ml. or diphenhiuron @ 1g./L of water.
- To control **fruit and shoot borer** in okra, Spray Bacillus thuringiensis 5% W.P @1.5 gram/lit of water. Present weather condition favours incidence of white fly and jassid. To control this apply Thiomethoxam@0.4g/ litre of water.

Fruits:

Spray mancozeb 2g/litre, to control sigatoka leaf spot in **banana**. To control **papaya yellow mosaic virus**, spray Imidachloprid @ 0.5 ml / litre.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-314
Date- 19/07/2022

AGROMET ADVISORY BULLETIN FOR TIGIRIA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 18mm and district received average 21mm rainfall during the last week. The mean daily maximum and minimum temperature was 32°C (-0.7°C from normal) and 25°C (+1.1°C from normal), respectively. Mean relative humidity was around 88-95%.

Forecasted period: Today(8.30IST) to 24th July(8.30IST)2022

Weather advisory: *Moderate with isolated heavy rainfall is forecasted for Friday and Saturday and light to moderate rainfall is forecasted for other 3 days in the block.* The day temperature will be 31-33°C and night temperature will be 25-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 83-86% and 51-57% respectively. **There will be partly cloudy to overcast sky for next 5 days.**

Block	Tigiria (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	19/07/2022	20/07/2022	21/07/2022	22/07/2022	23/07/2022
Rainfall(mm)	2	8	17	36	35
T-Max(°C)	32	31	32	33	32
T-Min(°C)	25	25	25	26	25
Cloud Cover(octa)	7	6	8	8	8
RH Max (%)	83	84	86	84	86
RH Min (%)	53	51	57	54	54
Wind Speed (Kmph)	14	11	10	8	9
Wind Direction (deg)	225	212	203	207	210

Agro met Advisory

General General advisory: *Drain out excess water from rice nurseries, orchards, maize and jute fields. Postpone irrigation, intercultural operations, application of plant protection measures and fertilizers to the standing crops from Thursday to Saturday and they may carry out weeding, pesticide and fertilizer application in upcoming 2 days. Keep the harvested produce at safer places. Cover the vegetable seedling with polythene to protect them from heavy shower.*

Kharifrice:(DIRECT SEEDED RICE):

Direct seeded rice are at three to four week stage. As light to moderate amount of rainfall is forecasted in the upcoming days, farmers should go for beushanning. Apply 35 kg urea/ac at beushanning followed by weeding and gap filling. Drain out the excess water at the time of fertilizer application and pond it after one day. If basal fertilizer not applied, apply 18 kg urea, 100kg ssp and 26 kg MOP while beushanning followed by top dressing of 35 kg urea after 7-10 days.

TRANSPLANTED RICE:

Considering the anticipated rainfall, paddy transplanting may be completed before 30th July. Apply 400-gram Chlorantraniliprole 0.4 % G or 800-gram Cartap Hydrochloride 4% GR in the 10 decimal nursery area 7 days before transplanting to manage gall midge, stem borer, caseworm, leaf folder and root knot nematode up to 3 weeks after transplanting. Intermittent dry and wet weather condition, is favourable for incidence of **BLB** and **Sheath blight** in early sown nursery seedlings. Go for foliar spray of Hexaconazole with Plantomycin@1.0 g/l to prevent the spread of the disease. Prevailing high temperature condition may lead to infestation of **rice thrips** in rice nursery. Apply Fipronil 5SC@2ml/l of water or Thiomethoxam@0.4g/ litre of water.

Maize:

Stem fly in Maize infects the plants from sowing to 5 weeks after sowing, temperature between 25-30 deg. celsius and humidity above 60% favour pest incidence. In vegetative stage to control stem fly spray Methyl demeton @1 ml/lit or Dimethoate @ 2ml/lit or apply Carbofuran 3G granuals @ 14 kg/acre. High humid conditions, temperature between 18-27 Deg. Celsius favours the occurrence of **leaf blight** in Maize. To control spray Azoxystrobin + Mancozeb (Pluton) @ 300 g /ac.

ARHAR: In arhar crop, for early variety apply 35 kg DAP, 4 kg Urea and 14 kg MOP as basal fertilizer per acre and for late variety apply 52 kg DAP, 6 kg Urea and 14 kg MOP as basal fertilizer per acre in dry days.

VEGETABLES: June-july is the best time for sowing of tomato in raised bed for nursery seedlings. Do seed treatment with Trichoderma viride @ 4g/kg seeds. The seed treatment must be done 24 hours before sowing of seeds.

- Due to forecast of rain, Chances of **fruit rot** infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever **sucking pests** incidence was observed spray 5 ml Neem oil or Thiomethoxam 0.2 g. per litre of water. **To prevent of soil borne diseases like seedling rot, stem rot and wilt diseases apply 3g of Copper oxychloride per litre of water around the main stem.**
- Incidence of **thrips** may observe in chilli, if observe spray Spinosad 45% SC @ 0.4 ml/litre of water or Imidacloprid 17.8% SL @ 0.3 ml/litre of water in clean and dry weather. The present weather conditions are favourable for Upward and downward leafcurl in chilli. To control spray phosalon (Zolone) 3 ml. or diphenthiuron @ 1g./L of water.
- To control **fruit and shoot borer** in okra, Spray Bacillus thuringiensis 5% W.P @1.5 gram/lit of water. Present weather condition favours incidence of white fly and jassid. To control this apply Thiomethoxam@0.4g/ litre of water.

Fruits: Spray mancozeb 2g/litre, to control sigatoka leaf spot in **banana**. To control **papaya yellow mosaic virus**, spray Imidachloprid @ 0.5 ml / litre.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-314
Date- 19/07/2022

				
MEGHDOOT APP	DAMINI APP	FACEBOOK (Damu Cuttack)	BLOCK SPECIFIC WHAT'S APP GROUPS	@DAMUCUTTACK



- Kindly install and use **“Meghdoot mobile App”** for Cuttack district weather forecast, warnings and weather based agromet advisory services.

<https://play.google.com/store/apps/details?id=com.aas.meghdoot>

- Kindly install and use **“Damini mobile App”** for location specific for lightening forecast and warnings.

<https://play.google.com/store/apps/details?id=com.lightening.live.damini>

- Follow the **block specific whats app groups** and join in your block wise groups for getting weather based agromet advisory services twice in a week (Tuesday and Friday) and other weather warning based farm management advisories.
- Follow the **Official Website links** to get the weather based agromet advisory service bulletins (**English & Odia**) twice in a week (Tuesday and Friday).

1	ICAR-NRRI, Cuttack, Odisha	https://icar-nrri.in/agromate/	
2	Agromet division, IMD, Pune	https://imdagrmet.gov.in/imdproject/AGIndex.php	

Dr Sujata Sethy
 (Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
 Contact No-9861445080/7008453427