




Cuttack District Block Wise Agro-met Advisory Service Bulletin

*Friday, 02nd December, 2022
(For the period 02nd to 06th December, 2022)*



*Issued by
District Agromet Unit,
KrishiVigyan Kendra Cuttack*







GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-353
Date- 02/12/2022

STANDARD PRECIPITATION INDEX AND NDVI MAP

Standard Precipitation Index (SPI)	Normalized difference vegetation Index (NDVI)
<p style="text-align: center;">Standardised Precipitation Index for the period 3rd November 2022 to 30th November 2022</p> <p style="text-align: center;">MINISTRY OF EARTH SCIENCES INDIA METEOROLOGICAL DEPARTMENT HYDROMET SECTION, CRS PUNE</p> 	<p style="text-align: center;">NOAA/VIIIRS/BLENDED NDVI Composite ending on Week no 47 (19.11.2022 to 25.11.2022) over Agricultural region</p> <p style="text-align: center;">Odisha</p>  <p>Agriculture vigour is good over few places of the state and moderate over rest of the state.</p>
<p><i>As per SPI value (cumulative 4weeks - 03November2022 – 30 November2022), milddrycondition was there in Cuttack district.</i></p>	<p><i>According to NDVI map (19.11.2022 to 25.11.2022) the agriculture vigor was moderate overmost places of Cuttack and light to moderate over BankiDampada, Athagad and Baranga,Cuttacksadar, Nialiblock of Cuttack district.</i></p>

LEGEND (Medium range block level rainfall forecast)		
	No Rainfall	00 mm
	Very light rainfall	01-05 mm
	Light rainfall	06-10 mm
	Light to moderate rainfall	11-30 mm
	Moderate to heavy rainfall	31-65 mm
	Heavy to very heavy rainfall	66-115 mm
	Extremely heavy rainfall	>116 mm

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com




Bulletin No.-353
Date- 02/12/2022

Past week (25 Nov-01 Dec, 2022) cumulative rainfall departure% from normal and (26 Nov-02 Dec, 2022) average minimum temperature and its departure from normal for Cuttack district



Medium range (Today (8.30IST) to Wednesday (8.30IST)) Rainfall (mm) forecast for Cuttack district



Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427
















GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT

KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-353
Date- 02/12/2022

Weather (Past and Forecasted) based Disease and pest in Major Crops and vegetables

Sl No	Crop	Stage and condition of the crop	Anticipated Disease and pest	
1	Rice	Milk dough to maturity		False smut
2	Sugarcane	Grand growth stage to cane development stag		Woolly Aphid
				Scale insect
				White fly
				Mealy bug
3	Mustard	Vegetative		Painted Bug
				Saw fly
4	Cole crops	Vegetative to fruiting		DBM
				Leaf eating caterpillar
				Aphid
				Fruit Borer
5	Vegetables	Vegetative to fruiting		sucking pest(Aphid, Mites, Thrips)
6	Tomato	Fruiting		Late Blight

Dr Sujata Sethy
 (Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-353
Date- 02/12/2022

AGROMET ADVISORY BULLETIN FOR ATHAGAD

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0mm rainfall during the last week. The mean daily maximum and minimum temperature was 29.2°C (-0.6) and 17°C (1.33), respectively. The mean relative humidity was 66-74%.

Forecasted period: Today(8.30IST) to 07th December(8.30IST)2022

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 29-30°C and night temperature will be 17-18°C for upcoming 5 days. Maximum and minimum relative humidity will be 54-77% and 24-35% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Athagad (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	02/12/2022	03/12/2022	04/12/2022	05/12/2022	06/12/2022
Rainfall (mm)	0	0	0	0	0
T-Max (°C)	30	30	30	30	29
T-Min (°C)	18	18	18	18	17
Cloud Cover (octa)	0	0	1	2	7
RH Max (%)	77	59	64	61	54
RH Min (%)	35	31	30	31	24
Wind Speed (kmph)	5	6	6	7	7
Wind Direction (deg)	15	72	23	25	11

Agromet Advisory

General Advisory: Due to forecasted low night temperature coupled with fog during upcoming days, there will be chance of infestation of sucking pests, leaf blight, powdery mildew in the rabi crops and vegetables, monitor the crops and vegetables, if symptoms are noticed, undertake plant protection measures. Provide light irrigation to standing vegetables and crops during morning hours to prevent damage from fog, low temperature and moisture stress. Carry out weeding, hoeing and other intercultural operations in rabi vegetables.

Kharif Rice:

Paddy is at milk dough stage to maturity stage. Drainout filed before 10-15 days of harvesting. **As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop.** For prevention of false smut spray the field before panicle initiation or flowering with propiconazole 25%EC @ 2ml/liter of water.

PULSES:

21-26°C soil temperature is required for proper germination of pulses. Due to forecast of dry weather for optimum germination irrigate the field after sowing. This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture.

Mustard:

Taking the advantage of available soil moisture, land preparation and sowing of the crop should be started to get optimum yield. **Apply irrigation in field if mustard was sown before 15-20 days. Due to increase in day temperature there may be infestation of saw fly and painted bug in Mustard, to control bug spray Dimethote 30 EC @ 1ml in liter of water and to control Saw fly spray imidachloprid 0.3 ml /liter of water.**

VEGETABLES:

As dry weather was prevailed in last 4 weeks and also forecasted for upcoming 5 days, farmers should irrigate their vegetable field in morning and evening hrs.

- Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. There is a chance of occurrence of **thrip, mites and Aphids** in Brinjal, Tomato, Chilli, cole crops, for control spray imidachloprid 0.3 ml /liter or fipronil 2 ml /liter of water.
- As Onion is shallow rooted crop, so at bulb initiation stage (after 60 days) due to forecasted dry weather farmers are advised to go for irrigation in the moisture sensitive stage.**
- Cabbage and Cauliflower:**
- The transplanted cabbage is at seedling establishment to three week stage. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by **diamond back moth**. To control this apply Spinosad 45SC @ 0.5ml/l.
- Brinjal:**
- Shoot and fruit borer and leaf eating Caterpillar may observe in brinjal field, to control it spray Emamectin benzoate 5% SG @ 0.4 g/litre of water or Chlorantraniliprole 18.5% SC @ 0.4 ml/litre of water. Spraying should be done during the early morning or evening hours.
- Tomato:**
- Due to decrease in minimum temperature and foggy condition there may be chances of late blight in tomato,** to avoid late blight precaution must be taken during the initial infection stage. Spray 2.5-gram Metalaxyl + Mancozeb/lit of water. Mix 3 gram Trichoderma with per kg of seed during seed treatment. Avoid excess nitrogen fertilizer and close spacing of plants.

Animal husbandry:

In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-353
Date- 02/12/2022

AGROMET ADVISORY BULLETIN FOR BADAMBA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0mm rainfall during the last week. The mean daily maximum and minimum temperature was 29.2°C (-0.6) and 17°C (1.33), respectively. The mean relative humidity was 66-74%.

Forecasted period: Today(8.30IST) to 07th December(8.30IST)2022

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 29-31°C and night temperature will be 16-18°C for upcoming 5 days. Maximum and minimum relative humidity will be 56-85% and 24-31% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Badamba (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	02/12/2022	03/12/2022	04/12/2022	05/12/2022	06/12/2022
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	31	30	31	30	29
T-Min(°C)	18	18	17	17	16
Cloud Cover(octa)	0	0	1	2	7
RH Max (%)	85	68	70	65	56
RH Min (%)	31	31	30	30	24
Wind Speed (Kmph)	5	5	5	6	6
Wind Direction (deg)	11	72	19	23	11

Agromet Advisory

General Advisory: Due to forecasted low night temperature coupled with fog during upcoming days, there will be chance of infestation of sucking pests, leaf blight, powdery mildew in the rabi crops and vegetables, monitor the crops and vegetables, if symptoms are noticed, undertake plant protection measures. Provide light irrigation to standing vegetables and crops during morning hours to prevent damage from fog, low temperature and moisture stress. Carry out weeding, hoeing and other intercultural operations in rabi vegetables.

Kharif Rice:

Paddy is at milk dough stage to maturity stage. Drainout filed before 10-15 days of harvesting. As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop. For prevention of false smut spray the field before panicle initiation or flowering with propiconazole 25%EC @ 2ml/liter of water.

PULSES:

21-26°C soil temperature is required for proper germination of pulses. Due to forecast of dry weather for optimum germination irrigate the field after sowing. This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture.

Mustard:

Taking the advantage of available soil moisture, land preparation and sowing of the crop should be started to get optimum yield. Apply irrigation in field if mustard was sown before 15-20 days. Due to increase in day temperature there may be infestation of saw fly and painted bug in Mustard, to control bug spray Dimethote 30 EC @ 1ml in liter of water and to control Saw fly spray imidachloprid 0.3 ml /liter of water.

VEGETABLES:

As dry weather was prevailed in last 4 weeks and also forecasted for upcoming 5 days, farmers should irrigate their vegetable field in morning and evening hrs.

- Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. There is a chance of occurrence of thrip, mites and Aphids in Brinjal, Tomato, Chilli, cole crops, for control spray imidachloprid 0.3 ml /liter or fipronil 2 ml /liter of water.
- As Onion is shallow rooted crop, so at bulb initiation stage (after 60 days) due to forecasted dry weather farmers are advised to go for irrigation in the moisture sensitive stage.
- Cabbage and Cauliflower:**
- The transplanted cabbage is at seedling establishment to three week stage. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by diamond back moth. To control this apply Spinosad 45SC @ 0.5ml/l.
- Brinjal:**
- Shoot and fruit borer and leaf eating Caterpillar may observe in brinjal field, to control it spray Emamectin benzoate 5% SG @ 0.4 g/litre of water or Chlorantraniliprole 18.5% SC @ 0.4 ml/litre of water. Spraying should be done during the early morning or evening hours.
- Tomato:**
- Due to decrease in minimum temperature and foggy condition there may be chances of late blight in tomato, to avoid late blight precaution must be taken during the initial infection stage. Spray 2.5-gram Metalaxyl + Mancozeb/lit of water. Mix 3 gram Trichoderma with per kg of seed during seed treatment. Avoid excess nitrogen fertilizer and close spacing of plants.

Animal husbandry:

In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-353
Date- 02/12/2022

AGROMET ADVISORY BULLETIN FOR BANKI

District Past week average weather condition

The block received average 0mm and district received average 0mm rainfall during the last week. The mean daily maximum and minimum temperature was 29.2°C (-0.6) and 17°C (1.33), respectively. The mean relative humidity was 66-74%.

Forecasted period: Today(8.30IST) to 07th December(8.30IST)2022

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 29-31°C and night temperature will be 17-18°C for upcoming 5 days. Maximum and minimum relative humidity will be 55-82% and 24-32% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Banki (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	02/12/2022	03/12/2022	04/12/2022	05/12/2022	06/12/2022
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	31	30	31	30	29
T-Min(°C)	18	18	18	18	17
Cloud Cover(octa)	0	0	1	2	8
RH Max (%)	82	64	67	62	55
RH Min (%)	32	31	30	30	24
Wind Speed (Kmph)	5	6	6	7	7
Wind Direction (deg)	15	72	23	25	11

Agromet Advisory

General Advisory: Due to forecasted low night temperature coupled with fog during upcoming days, there will be chance of infestation of sucking pests, leaf blight, powdery mildew in the rabi crops and vegetables, monitor the crops and vegetables, if symptoms are noticed, undertake plant protection measures. Provide light irrigation to standing vegetables and crops during morning hours to prevent damage from fog, low temperature and moisture stress. Carry out weeding, hoeing and other intercultural operations in rabi vegetables.

Kharif Rice:

Paddy is at milk dough stage to maturity stage. Drainout filed before 10-15 days of harvesting. **As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop.** For prevention of false smut spray the field before panicle initiation or flowering with propiconazole 25%EC @ 2ml/liter of water.

PULSES:

21-26°C soil temperature is required for proper germination of pulses. Due to forecast of dry weather for optimum germination irrigate the field after sowing. This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture.

Mustard:

Taking the advantage of available soil moisture, land preparation and sowing of the crop should be started to get optimum yield. **Apply irrigation in field if mustard was sown before 15-20 days. Due to increase in day temperature there may be infestation of saw fly and painted bug in Mustard, to control bug spray Dimethote 30 EC @ 1ml in liter of water and to control Saw fly spray imidachloprid 0.3 ml /liter of water.**

VEGETABLES:

As dry weather was prevailed in last 4 weeks and also forecasted for upcoming 5 days, farmers should irrigate their vegetable field in morning and evening hrs.

- Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. There is a chance of occurrence of **thrip, mites and Aphids** in Brinjal, Tomato, Chilli, cole crops, for control spray imidachloprid 0.3 ml /liter or fipronil 2 ml /liter of water.
- As Onion is shallow rooted crop, so at bulb initiation stage (after 60 days) due to forecasted dry weather farmers are advised to go for irrigation in the moisture sensitive stage.**
- Cabbage and Cauliflower:**
- The transplanted cabbage is at seedling establishment to three week stage. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by **diamond back moth**. To control this apply Spinosad 45SC @ 0.5ml/l.
- Brinjal:**
- Shoot and fruit borer and leaf eating Caterpillar** may observe in brinjal field, to control it spray Emamectin benzoate 5% SG @ 0.4 g/litre of water or Chlorantraniliprole 18.5% SC @ 0.4 ml/litre of water. Spraying should be done during the early morning or evening hours.
- Tomato:**
- Due to decrease in minimum temperature and foggy condition there may be chances of late blight in tomato,** to avoid late blight precaution must be taken during the initial infection stage. Spray 2.5-gram Metalaxyl + Mancozeb/lit of water. Mix 3 gram Trichoderma with per kg of seed during seed treatment. Avoid excess nitrogen fertilizer and close spacing of plants.

Animal husbandry:

- In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-353
Date- 02/12/2022

AGROMET ADVISORY BULLETIN FOR BANKI-DAMPADA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0mm rainfall during the last week. The mean daily maximum and minimum temperature was 29.2°C (-0.6) and 17°C (1.33), respectively. The mean relative humidity was 66-74%.

Forecasted period: Today(8.30IST) to 07th December(8.30IST)2022

Weather Advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 29-31°C and night temperature will be 17-18°C for upcoming 5 days. Maximum and minimum relative humidity will be 55-80% and 24-33% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Banki-Dampada (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	02/12/2022	03/12/2022	04/12/2022	05/12/2022	06/12/2022
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	31	30	30	30	29
T-Min(°C)	18	18	18	18	17
Cloud Cover(octa)	0	0	1	2	8
RH Max (%)	80	63	66	61	55
RH Min (%)	33	30	29	30	24
Wind Speed (Kmph)	5	6	6	7	7
Wind Direction (deg)	15	72	23	25	11

Agromet Advisory

General Advisory: Due to forecasted low night temperature coupled with fog during upcoming days, there will be chance of infestation of sucking pests, leaf blight, powdery mildew in the rabi crops and vegetables, monitor the crops and vegetables, if symptoms are noticed, undertake plant protection measures. Provide light irrigation to standing vegetables and crops during morning hours to prevent damage from fog, low temperature and moisture stress. Carry out weeding, hoeing and other intercultural operations in rabi vegetables.

Kharif Rice:

Paddy is at milk dough stage to maturity stage. Drainout filed before 10-15 days of harvesting. **As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop.** For prevention of false smut spray the field before panicle initiation or flowering with propiconazole 25%EC@ 2ml/liter of water.

PULSES:

21-26°C soil temperature is required for proper germination of pulses. Due to forecast of dry weather for optimum germination irrigate the field after sowing. This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture.

Mustard:

Taking the advantage of available soil moisture, land preparation and sowing of the crop should be started to get optimum yield. **Apply irrigation in field if mustard was sown before 15-20 days.** Due to increase in day temperature there may be infestation of saw fly and painted bug in Mustard, to control bug spray Dimethote 30 EC @ 1ml in liter of water and to control Saw fly spray imidachloprid 0.3 ml /liter of water.

VEGETABLES:

As dry weather was prevailed in last 4 weeks and also forecasted for upcoming 5 days, farmers should irrigate their vegetable field in morning and evening hrs.

- Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. There is a chance of occurrence of **thrip, mites and Aphids** in Brinjal, Tomato, Chilli, cole crops, for control spray imidachloprid 0.3 ml /liter or fipronil 2 ml /liter of water.
- As Onion is shallow rooted crop, so at bulb initiation stage (after 60 days) due to forecasted dry weather farmers are advised to go for irrigation in the moisture sensitive stage.**
- Cabbage and Cauliflower:**
- The transplanted cabbage is at seedling establishment to three week stage. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by **diamond back moth**. To control this apply Spinosad 45SC @ 0.5ml/l.
- Brinjal:**
- Shoot and fruit borer and leaf eating Caterpillar** may observe in brinjal field, to control it spray Emamectin benzoate 5% SG @ 0.4 g/litre of water or Chlorantraniliprole 18.5% SC @ 0.4 ml/litre of water. Spraying should be done during the early morning or evening hours.
- Tomato:**
- Due to decrease in minimum temperature and foggy condition there may be chances of late blight in tomato,** to avoid late blight precaution must be taken during the initial infection stage. Spray 2.5-gram Metalaxyl + Mancozeb/lit of water. Mix 3 gram Trichoderma with per kg of seed during seed treatment. Avoid excess nitrogen fertilizer and close spacing of plants.

Animal husbandry:

- In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-353
Date- 02/12/2022

AGROMET ADVISORY BULLETIN FOR BARANGA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0mm rainfall during the last week. The mean daily maximum and minimum temperature was 29.2°C (-0.6) and 17°C (1.33), respectively. The mean relative humidity was 66-74%.

Forecasted period: Today(8.30IST) to 07th December(8.30IST)2022

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 29-30°C and night temperature will be 17-18°C for upcoming 5 days. Maximum and minimum relative humidity will be 51-69% and 25-37% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Baranga (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	02/12/2022	03/12/2022	04/12/2022	05/12/2022	06/12/2022
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	30	30	30	30	29
T-Min(°C)	18	18	18	18	17
Cloud Cover	0	0	1	1	7
RH Max (%)	69	55	57	56	51
RH Min (%)	37	29	30	29	25
Wind Speed (Kmph)	7	7	7	8	9
Wind Direction (deg)	23	23	15	25	27

Agromet Advisory

General Advisory: Due to forecasted low night temperature coupled with fog during upcoming days, there will be chance of infestation of sucking pests, leaf blight, powdery mildew in the rabi crops and vegetables, monitor the crops and vegetables, if symptoms are noticed, undertake plant protection measures. Provide light irrigation to standing vegetables and crops during morning hours to prevent damage from fog, low temperature and moisture stress. Carry out weeding, hoeing and other intercultural operations in rabi vegetables.

Kharif Rice:

Paddy is at milk dough stage to maturity stage. Drainout filed before 10-15 days of harvesting. **As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop.** For prevention of false smut spray the field before panicle initiation or flowering with propiconazole 25%EC @ 2ml/liter of water.

PULSES:

21-26°C soil temperature is required for proper germination of pulses. Due to forecast of dry weather for optimum germination irrigate the field after sowing. This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture.

Mustard:

Taking the advantage of available soil moisture, land preparation and sowing of the crop should be started to get optimum yield. **Apply irrigation in field if mustard was sown before 15-20 days.** Due to increase in day temperature there may be infestation of saw fly and painted bug in Mustard, to control bug spray Dimethote 30 EC @ 1ml in liter of water and to control Saw fly spray imidachloprid 0.3 ml /liter of water.

VEGETABLES:

As dry weather was prevailed in last 4 weeks and also forecasted for upcoming 5 days, farmers should irrigate their vegetable field in morning and evening hrs.

- Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. There is a chance of occurrence of **thrip, mites and Aphids** in Brinjal, Tomato, Chilli, cole crops, for control spray imidachloprid 0.3 ml /liter or fipronil 2 ml /liter of water.
- As Onion is shallow rooted crop, so at bulb initiation stage (after 60 days) due to forecasted dry weather farmers are advised to go for irrigation in the moisture sensitive stage.**
- Cabbage and Cauliflower:**
- The transplanted cabbage is at seedling establishment to three week stage. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by **diamond back moth**. To control this apply Spinosad 45SC @ 0.5ml/l.
- Brinjal:**
- Shoot and fruit borer and leaf eating Caterpillar** may observe in brinjal field, to control it spray Emamectin benzoate 5% SG @ 0.4 g/litre of water or Chlorantraniliprole 18.5% SC @ 0.4 ml/litre of water. Spraying should be done during the early morning or evening hours.
- Tomato:**
- Due to decrease in minimum temperature and foggy condition there may be chances of late blight in tomato,** to avoid late blight precaution must be taken during the initial infection stage. Spray 2.5-gram Metalaxyl + Mancozeb/lit of water. Mix 3 gram Trichoderma with per kg of seed during seed treatment. Avoid excess nitrogen fertilizer and close spacing of plants.

Animal husbandry:

- In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-353
Date- 02/12/2022

AGROMET ADVISORY BULLETIN FOR CUTTACK SADAR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0mm rainfall during the last week. The mean daily maximum and minimum temperature was 29.2°C (-0.6) and 17°C (1.33), respectively. The mean relative humidity was 66-74%.

Forecasted period: Today(8.30IST) to 07th December(8.30IST)2022

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 29-30°C and night temperature will be 18-19°C for upcoming 5 days. Maximum and minimum relative humidity will be 51-67% and 25-37% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Cuttack sadar (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	02/12/2022	03/12/2022	04/12/2022	05/12/2022	06/12/2022
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	30	30	30	30	29
T-Min(°C)	18	19	18	18	18
Cloud Cover(octa)	0	0	1	1	7
RH Max (%)	67	53	59	57	51
RH Min (%)	37	29	30	30	25
Wind Speed (Kmph)	6	7	7	8	8
Wind Direction (deg)	23	68	23	28	27

Agromet Advisory

General Advisory: Due to forecasted low night temperature coupled with fog during upcoming days, there will be chance of infestation of sucking pests, leaf blight, powdery mildew in the rabi crops and vegetables, monitor the crops and vegetables, if symptoms are noticed, undertake plant protection measures. Provide light irrigation to standing vegetables and crops during morning hours to prevent damage from fog, low temperature and moisture stress. Carry out weeding, hoeing and other intercultural operations in rabi vegetables.

Kharif Rice:

Paddy is at milk dough stage to maturity stage. Drainout filed before 10-15 days of harvesting. **As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop.** For prevention of false smut spray the field before panicle initiation or flowering with propiconazole 25%EC @ 2ml/liter of water.

Sugarcane:

The crop is at maturity (40 - 41 week) stage. Postpone irrigation to crops and spraying insecticides and pesticides. For control of Sucking pest in sugarcane like **Woolly aphid, Whitefly, Scale insects, mealybug** in present weather conditions go for alternate spraying of thiomethoxam 25WG @ 0.25 gm/lit or imidacloprid 0.45 ml/lit of water or Dinotefuron 20% SG 3gm/lit of Water.

Mustard:

Taking the advantage of available soil moisture, land preparation and sowing of the crop should be started to get optimum yield. **Apply irrigation in field if mustard was sown before 15-20 days. Due to increase in day temperature there may be infestation of saw fly and painted bug in Mustard, to control bug spray Dimethote 30 EC @ 1ml in liter of water and to control Saw fly spray imidachloprid 0.3 ml /liter of water.**

VEGETABLES:

As dry weather was prevailed in last 4 weeks and also forecasted for upcoming 5 days, farmers should irrigate their vegetable field in morning and evening hrs.

- Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. There is a chance of occurrence of **thrip, mites and Aphids** in Brinjal, Tomato, Chilli, cole crops, for control spray imidachloprid 0.3 ml /liter or fipronil 2 ml /liter of water.
- As Onion is shallow rooted crop, so at bulb initiation stage (after 60 days) due to forecasted dry weather farmers are advised to go for irrigation in the moisture sensitive stage.**
- Cabbage and Cauliflower:**
- The transplanted cabbage is at seedling establishment to three week stage. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by **diamond back moth**. To control this apply Spinosad 45SC @ 0.5ml/l.
- Brinjal:**
- Shoot and fruit borer and leaf eating Caterpillar** may observe in brinjal field, to control it spray Emamectin benzoate 5% SG @ 0.4 g/litre of water or Chlorantraniliprole 18.5% SC @ 0.4 ml/litre of water. Spraying should be done during the early morning or evening hours.
- Tomato:**
- Due to decrease in minimum temperature and foggy condition there may be chances of late blight in tomato,** to avoid late blight precaution must be taken during the initial infection stage. Spray 2.5-gram Metalaxyl + Mancozeb/lit of water. Mix 3 gram Trichoderma with per kg of seed during seed treatment. Avoid excess nitrogen fertilizer and close spacing of plants.

Animal husbandry:

In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-353
Date- 02/12/2022

AGROMET ADVISORY BULLETIN FOR KANTAPADA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0mm rainfall during the last week. The mean daily maximum and minimum temperature was 29.2°C (-0.6) and 17°C (1.33), respectively. The mean relative humidity was 66-74%.

Forecasted period: Today(8.30IST) to 07th December(8.30IST)2022

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 29-31°C and night temperature will be 17-19°C for upcoming 5 days. Maximum and minimum relative humidity will be 50-66% and 26-38% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Kantapada (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	02/12/2022	03/12/2022	04/12/2022	05/12/2022	06/12/2022
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	31	30	30	30	29
T-Min(°C)	18	19	18	18	17
Cloud Cover(octa)	0	0	3	1	8
RH Max (%)	66	56	56	55	50
RH Min (%)	38	29	30	28	26
Wind Speed (Kmph)	8	8	9	10	11
Wind Direction (deg)	15	68	23	26	25

Agromet Advisory

General Advisory: Due to forecasted low night temperature coupled with fog during upcoming days, there will be chance of infestation of sucking pests, leaf blight, powdery mildew in the rabi crops and vegetables, monitor the crops and vegetables, if symptoms are noticed, undertake plant protection measures. Provide light irrigation to standing vegetables and crops during morning hours to prevent damage from fog, low temperature and moisture stress. Carry out weeding, hoeing and other intercultural operations in rabi vegetables.

Kharif Rice:

Paddy is at milk dough stage to maturity stage. Drainout filed before 10-15 days of harvesting. **As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop.** For prevention of false smut spray the field before panicle initiation or flowering with propiconazole 25%EC @ 2ml/liter of water.

Sugarcane:

The crop is at maturity (40 - 41 week) stage. Postpone irrigation to crops and spraying insecticides and pesticides. For control of Sucking pest in sugarcane like Woolly aphid, Whitefly, Scale insects, mealybug in present weather conditions go for alternate spraying of thiomethoxam 25WG @ 0.25 gm/lit or imidacloprid 0.45 ml/lit of water or Dinotefuron 20% SG 3gm/lit of Water.

Mustard:

Taking the advantage of available soil moisture, land preparation and sowing of the crop should be started to get optimum yield. **Apply irrigation in field if mustard was sown before 15-20 days. Due to increase in day temperature there may be infestation of saw fly and painted bug in Mustard, to control bug spray Dimethote 30 EC @ 1ml in liter of water and to control Saw fly spray imidachloprid 0.3 ml /liter of water.**

VEGETABLES:

As dry weather was prevailed in last 4 weeks and also forecasted for upcoming 5 days, farmers should irrigate their vegetable field in morning and evening hrs.

- Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. There is a chance of occurrence of thrip, mites and Aphids in Brinjal, Tomato, Chilli, cole crops, for control spray imidachloprid 0.3 ml /liter or fipronil 2 ml /liter of water.
- As Onion is shallow rooted crop, so at bulb initiation stage (after 60 days) due to forecasted dry weather farmers are advised to go for irrigation in the moisture sensitive stage.**
- Cabbage and Cauliflower:**
- The transplanted cabbage is at seedling establishment to three week stage. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by diamond back moth. To control this apply Spinosad 45SC @ 0.5ml/l.
- Brinjal:**
- Shoot and fruit borer and leaf eating Caterpillar may observe in brinjal field, to control it spray Emamectin benzoate 5% SG @ 0.4 g/litre of water or Chlorantraniliprole 18.5% SC @ 0.4 ml/litre of water. Spraying should be done during the early morning or evening hours.
- Tomato:**
- Due to decrease in minimum temperature and foggy condition there may be chances of late blight in tomato,** to avoid late blight precaution must be taken during the initial infection stage. Spray 2.5-gram Metalaxyl + Mancozeb/lit of water. Mix 3 gram Trichoderma with per kg of seed during seed treatment. Avoid excess nitrogen fertilizer and close spacing of plants.

Animal husbandry:

In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-353
Date- 02/12/2022

AGROMET ADVISORY BULLETIN FOR MAHANGA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0mm rainfall during the last week. The mean daily maximum and minimum temperature was 29.2°C (-0.6) and 17°C (1.33), respectively. The mean relative humidity was 66-74%.

Forecasted period: Today(8.30IST) to 07th December(8.30IST)2022

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be and it will be 29-30°C and night temperature will be 17-18°C for upcoming 5 days. Maximum and minimum relative humidity will be 50-59% and 26-39% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Mahanga (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	02/12/2022	03/12/2022	04/12/2022	05/12/2022	06/12/2022
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	30	30	30	30	29
T-Min(°C)	18	18	18	17	17
Cloud Cover(octal)	0	0	1	1	8
RH Max (%)	59	50	58	56	50
RH Min (%)	39	28	30	29	26
Wind Speed (Kmph)	8	8	8	9	10
Wind Direction (deg)	23	30	18	27	25

Agromet Advisory

General Advisory: Due to forecasted low night temperature coupled with fog during upcoming days, there will be chance of infestation of sucking pests, leaf blight, powdery mildew in the rabi crops and vegetables, monitor the crops and vegetables, if symptoms are noticed, undertake plant protection measures. Provide light irrigation to standing vegetables and crops during morning hours to prevent damage from fog, low temperature and moisture stress. Carry out weeding, hoeing and other intercultural operations in rabi vegetables.

Kharif Rice:

Paddy is at milk dough stage to maturity stage. Drainout filed before 10-15 days of harvesting. **As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop.** For prevention of false smut spray the field before panicle initiation or flowering with propiconazole 25%EC @ 2ml/liter of water.

Sugarcane:

The crop is at maturity (40 - 41 week) stage. Postpone irrigation to crops and spraying insecticides and pesticides. For control of Sucking pest in sugarcane like Woolly aphid, Whitefly, Scale insects, mealybug in present weather conditions go for alternate spraying of thiomethoxam 25WG @ 0.25 gm/lit or imidacloprid 0.45 ml/lit of water or Dinotefuron 20% SG 3gm/lit of Water.

Mustard:

Taking the advantage of available soil moisture, land preparation and sowing of the crop should be started to get optimum yield. **Apply irrigation in field if mustard was sown before 15-20 days. Due to increase in day temperature there may be infestation of saw fly and painted bug in Mustard, to control bug spray Dimethote 30 EC @ 1ml in liter of water and to control Saw fly spray imidachloprid 0.3 ml /liter of water.**

VEGETABLES:

As dry weather was prevailed in last 4 weeks and also forecasted for upcoming 5 days, farmers should irrigate their vegetable field in morning and evening hrs.

- Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. There is a chance of occurrence of **thrip, mites and Aphids** in Brinjal, Tomato, Chilli, cole crops, for control spray imidachloprid 0.3 ml /liter or fipronil 2 ml /liter of water.
- As Onion is shallow rooted crop, so at bulb initiation stage (after 60 days) due to forecasted dry weather farmers are advised to go for irrigation in the moisture sensitive stage.**
- Cabbage and Cauliflower:**
- The transplanted cabbage is at seedling establishment to three week stage. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by **diamond back moth**. To control this apply Spinosad 45SC @ 0.5ml/l.
- Brinjal:**
- Shoot and fruit borer and leaf eating Caterpillar** may observe in brinjal field, to control it spray Emamectin benzoate 5% SG @ 0.4 g/litre of water or Chlorantraniliprole 18.5% SC @ 0.4 ml/litre of water. Spraying should be done during the early morning or evening hours.
- Tomato:**
- Due to decrease in minimum temperature and foggy condition there may be chances of late blight in tomato,** to avoid late blight precaution must be taken during the initial infection stage. Spray 2.5-gram Metalaxyl + Mancozeb/lit of water. Mix 3 gram Trichoderma with per kg of seed during seed treatment. Avoid excess nitrogen fertilizer and close spacing of plants.

Animal husbandry:

- In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-353
Date- 02/12/2022

AGROMET ADVISORY BULLETIN FOR NARASINGHPUR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0mm rainfall during the last week. The mean daily maximum and minimum temperature was 29.2°C (-0.6) and 17°C (1.33), respectively. The mean relative humidity was 66-74%.

Forecasted period: Today(8.30IST) to 07th December(8.30IST)2022

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 28-31°C and night temperature will be 15-17°C for upcoming 5 days. Maximum and minimum relative humidity will be 56-87% and 24-32% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Narasinghpur (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	02/12/2022	03/12/2022	04/12/2022	05/12/2022	06/12/2022
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	30	31	31	30	28
T-Min(°C)	17	16	16	16	15
Cloud Cover(/octa)	0	0	1	1	7
RH Max (%)	87	75	75	68	56
RH Min (%)	28	32	29	29	24
Wind Speed (Kmph)	3	4	4	4	5
Wind Direction (deg)	30	90	23	15	11

Agromet Advisory

General Advisory: Due to forecasted low night temperature coupled with fog during upcoming days, there will be chance of infestation of sucking pests, leaf blight, powdery mildew in the rabi crops and vegetables, monitor the crops and vegetables, if symptoms are noticed, undertake plant protection measures. Provide light irrigation to standing vegetables and crops during morning hours to prevent damage from fog, low temperature and moisture stress. Carry out weeding, hoeing and other intercultural operations in rabi vegetables.

Kharif Rice:

Paddy is at milk dough stage to maturity stage. Drainout filed before 10-15 days of harvesting. **As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop.** For prevention of false smut spray the field before panicle initiation or flowering with propiconazole 25%EC @ 2ml/liter of water.

PULSES:

21-26°C soil temperature is required for proper germination of pulses. Due to forecast of dry weather for optimum germination irrigate the field after sowing. This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture.

Mustard:

Taking the advantage of available soil moisture, land preparation and sowing of the crop should be started to get optimum yield. **Apply irrigation in field if mustard was sown before 15-20 days.** Due to increase in day temperature there may be infestation of saw fly and painted bug in Mustard, to control bug spray Dimethote 30 EC @ 1ml in liter of water and to control Saw fly spray imidachloprid 0.3 ml /liter of water.

VEGETABLES:

As dry weather was prevailed in last 4 weeks and also forecasted for upcoming 5 days, farmers should irrigate their vegetable field in morning and evening hrs.

- Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. There is a chance of occurrence of **thrip, mites and Aphids** in Brinjal, Tomato, Chilli, cole crops, for control spray imidachloprid 0.3 ml /liter or fipronil 2 ml /liter of water.
- As Onion is shallow rooted crop, so at bulb initiation stage (after 60 days) due to forecasted dry weather farmers are advised to go for irrigation in the moisture sensitive stage.**
- Cabbage and Cauliflower:**
- The transplanted cabbage is at seedling establishment to three week stage. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by **diamond back moth**. To control this apply Spinosad 45SC @ 0.5ml/l.
- Brinjal:**
- Shoot and fruit borer and leaf eating Caterpillar** may observe in brinjal field, to control it spray Emamectin benzoate 5% SG @ 0.4 g/litre of water or Chlorantraniliprole 18.5% SC @ 0.4 ml/litre of water. Spraying should be done during the early morning or evening hours.
- Tomato:**
- Due to decrease in minimum temperature and foggy condition there may be chances of late blight in tomato,** to avoid late blight precaution must be taken during the initial infection stage. Spray 2.5-gram Metalaxyl + Mancozeb/lit of water. Mix 3 gram Trichoderma with per kg of seed during seed treatment. Avoid excess nitrogen fertilizer and close spacing of plants.

Animal husbandry:

- In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-353
Date- 02/12/2022

AGROMET ADVISORY BULLETIN FOR NIALI

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0mm rainfall during the last week. The mean daily maximum and minimum temperature was 29.2°C (-0.6) and 17°C (1.33), respectively. The mean relative humidity was 66-74%.

Forecasted period: Today(8.30IST) to 07th December(8.30IST)2022

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 29-30°C and night temperature will be 17-19°C for upcoming 5 days. Maximum and minimum relative humidity will be 50-66% and 27-39% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Niali (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	02/12/2022	03/12/2022	04/12/2022	05/12/2022	06/12/2022
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	30	30	30	30	29
T-Min(°C)	18	19	18	18	17
Cloud Cover(octa)	0	0	4	2	8
RH Max (%)	66	58	57	57	50
RH Min (%)	39	30	30	28	27
Wind Speed(Kmph)	9	9	10	11	12
Wind Direction(deg)	15	60	18	23	27

Agromet Advisory

General Advisory: Due to forecasted low night temperature coupled with fog during upcoming days, there will be chance of infestation of sucking pests, leaf blight, powdery mildew in the rabi crops and vegetables, monitor the crops and vegetables, if symptoms are noticed, undertake plant protection measures. Provide light irrigation to standing vegetables and crops during morning hours to prevent damage from fog, low temperature and moisture stress. Carry out weeding, hoeing and other intercultural operations in rabi vegetables.

Kharif Rice:

Paddy is at milk dough stage to maturity stage. Drainout filed before 10-15 days of harvesting. **As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop.** For prevention of false smut spray the field before panicle initiation or flowering with propiconazole 25%EC @ 2ml/liter of water.

Sugarcane:

The crop is at maturity (40 - 41 week) stage. Postpone irrigation to crops and spraying insecticides and pesticides. For control of Sucking pest in sugarcane like Woolly aphid, Whitefly, Scale insects, mealybug in present weather conditions go for alternate spraying of thiomethoxam 25WG @ 0.25 gm/lit or imidacloprid 0.45 ml/lit of water or Dinotefuron 20% SG 3gm/lit of Water.

Mustard:

Taking the advantage of available soil moisture, land preparation and sowing of the crop should be started to get optimum yield. **Apply irrigation in field if mustard was sown before 15-20 days. Due to increase in day temperature there may be infestation of saw fly and painted bug in Mustard, to control bug spray Dimethothate 30 EC @ 1ml in liter of water and to control Saw fly spray imidachloprid 0.3 ml /liter of water.**

VEGETABLES:

As dry weather was prevailed in last 4 weeks and also forecasted for upcoming 5 days, farmers should irrigate their vegetable field in morning and evening hrs.

- Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. There is a chance of occurrence of **thrip, mites and Aphids** in Brinjal, Tomato, Chilli, cole crops, for control spray imidachloprid 0.3 ml /liter or fipronil 2 ml /liter of water.
- As Onion is shallow rooted crop, so at bulb initiation stage (after 60 days) due to forecasted dry weather farmers are advised to go for irrigation in the moisture sensitive stage.**
- Cabbage and Cauliflower:**
- The transplanted cabbage is at seedling establishment to three week stage. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by **diamond back moth**. To control this apply Spinosad 45SC @ 0.5ml/l.
- Brinjal:**
- Shoot and fruit borer and leaf eating Caterpillar** may observe in brinjal field, to control it spray Emamectin benzoate 5% SG @ 0.4 g/litre of water or Chlorantraniliprole 18.5% SC @ 0.4 ml/litre of water. Spraying should be done during the early morning or evening hours.
- Tomato:**
- Due to decrease in minimum temperature and foggy condition there may be chances of late blight in tomato**, to avoid late blight precaution must be taken during the initial infection stage. Spray 2.5-gram Metalaxyl + Mancozeb/lit of water. Mix 3 gram Trichoderma with per kg of seed during seed treatment. Avoid excess nitrogen fertilizer and close spacing of plants.

Animal husbandry:

- In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-353
Date- 02/12/2022

AGROMET ADVISORY BULLETIN FOR NISCHINTAKOILI
District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0mm rainfall during the last week. The mean daily maximum and minimum temperature was 29.2°C (-0.6) and 17°C (1.33), respectively. The mean relative humidity was 66-74%.

Forecasted period: Today(8.30IST) to 07th December(8.30IST)2022

Weather advisory: Dry weather is forecasted for upcoming 5 Days in the block. The day temperature will be 29-30°C and night temperature will be 17-18°C for upcoming 5 days. Maximum and minimum relative humidity will be 50-59% and 27-39% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Nischintakoili (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	02/12/2022	03/12/2022	04/12/2022	05/12/2022	06/12/2022
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	30	30	30	30	29
T-Min(°C)	18	18	18	18	17
Cloud Cover(octa)	0	0	2	1	8
RH Max (%)	59	51	57	56	50
RH Min (%)	39	28	30	28	27
Wind Speed (Kmph)	8	8	8	10	11
Wind Direction (deg)	23	30	23	26	25

Agromet Advisory

General Advisory: Due to forecasted low night temperature coupled with fog during upcoming days, there will be chance of infestation of sucking pests, leaf blight, powdery mildew in the rabi crops and vegetables, monitor the crops and vegetables, if symptoms are noticed, undertake plant protection measures. Provide light irrigation to standing vegetables and crops during morning hours to prevent damage from fog, low temperature and moisture stress. Carry out weeding, hoeing and other intercultural operations in rabi vegetables.

Kharif Rice:

Paddy is at milk dough stage to maturity stage. Drainout filed before 10-15 days of harvesting. **As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop.** For prevention of false smut spray the field before panicle initiation or flowering with propiconazole 25%EC @ 2ml/liter of water.

Sugarcane:

The crop is at maturity (40 - 41 week) stage. Postpone irrigation to crops and spraying insecticides and pesticides. For control of Sucking pest in sugarcane like **Woolly aphid, Whitefly, Scale insects, mealy bug** in present weather conditions go for alternate spraying of thiomethoxam 25WG @ 0.25 gm/lit or imidacloprid 0.45 ml/lit of water or Dinotefuron 20% SG 3gm/lit of Water.

Mustard:

Taking the advantage of available soil moisture, land preparation and sowing of the crop should be started to get optimum yield. **Apply irrigation in field if mustard was sown before 15-20 days. Due to increase in day temperature there may be infestation of saw fly and painted bug in Mustard, to control bug spray Dimethote 30 EC @ 1ml in liter of water and to control Saw fly spray imidachloprid 0.3 ml /liter of water.**

VEGETABLES:

As dry weather was prevailed in last 4 weeks and also forecasted for upcoming 5 days, farmers should irrigate their vegetable field in morning and evening hrs.

- Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. There is a chance of occurrence of **thrip, mites and Aphids** in Brinjal, Tomato, Chilli, cole crops, for control spray imidachloprid 0.3 ml /liter or fipronil 2 ml /liter of water.
- As Onion is shallow rooted crop, so at bulb initiation stage (after 60 days) due to forecasted dry weather farmers are advised to go for irrigation in the moisture sensitive stage.**
- Cabbage and Cauliflower:**
- The transplanted cabbage is at seedling establishment to three week stage. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by **diamond back moth**. To control this apply Spinosad 45SC @ 0.5ml/l.
- Brinjal:**
- Shoot and fruit borer and leaf eating Caterpillar** may observe in brinjal field, to control it spray Emamectin benzoate 5% SG @ 0.4 g/litre of water or Chlorantraniliprole 18.5% SC @ 0.4 ml/litre of water. Spraying should be done during the early morning or evening hours.
- Tomato:**
- Due to decrease in minimum temperature and foggy condition there may be chances of late blight in tomato,** to avoid late blight precaution must be taken during the initial infection stage. Spray 2.5-gram Metalaxyl + Mancozeb/lit of water. Mix 3 gram Trichoderma with per kg of seed during seed treatment. Avoid excess nitrogen fertilizer and close spacing of plants.

Animal husbandry:

- In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-353
Date- 02/12/2022

AGROMET ADVISORY BULLETIN FOR SALEPUR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0mm rainfall during the last week. The mean daily maximum and minimum temperature was 29.2°C (-0.6) and 17°C (1.33), respectively. The mean relative humidity was 66-74%.

Forecasted period: Today(8.30IST) to 07th December(8.30IST)2022

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 29-30°C and night temperature will be 17-18°C for upcoming 5 days. Maximum and minimum relative humidity will be 50-60% and 26-39% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Salepur (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	02/12/2022	03/12/2022	04/12/2022	05/12/2022	06/12/2022
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	30	30	30	30	29
T-Min(°C)	18	18	18	18	17
Cloud Cover(octa)	0	0	1	1	8
RH Max (%)	60	50	57	56	50
RH Min (%)	39	28	30	29	26
Wind Speed (Kmph)	8	8	8	9	11
Wind Direction (deg)	23	30	18	26	25

Agromet Advisory

General Advisory: Due to forecasted low night temperature coupled with fog during upcoming days, there will be chance of infestation of sucking pests, leaf blight, powdery mildew in the rabi crops and vegetables, monitor the crops and vegetables, if symptoms are noticed, undertake plant protection measures. Provide light irrigation to standing vegetables and crops during morning hours to prevent damage from fog, low temperature and moisture stress. Carry out weeding, hoeing and other intercultural operations in rabi vegetables.

Kharif Rice:

Paddy is at milk dough stage to maturity stage. Drainout filed before 10-15 days of harvesting. **As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop.** For prevention of false smut spray the field before panicle initiation or flowering with propiconazole 25%EC @ 2ml/liter of water.

PULSES:

21-26°C soil temperature is required for proper germination of pulses. Due to forecast of dry weather for optimum germination irrigate the field after sowing. This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture.

Mustard:

Taking the advantage of available soil moisture, land preparation and sowing of the crop should be started to get optimum yield. **Apply irrigation in field if mustard was sown before 15-20 days.** Due to increase in day temperature there may be infestation of saw fly and painted bug in Mustard, to control bug spray Dimethote 30 EC @ 1ml in liter of water and to control Saw fly spray imidachloprid 0.3 ml /liter of water.

VEGETABLES:

As dry weather was prevailed in last 4 weeks and also forecasted for upcoming 5 days, farmers should irrigate their vegetable field in morning and evening hrs.

- Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. There is a chance of occurrence of **thrip, mites and Aphids** in Brinjal, Tomato, Chilli, cole crops, for control spray imidachloprid 0.3 ml /liter or fipronil 2 ml /liter of water.
- As Onion is shallow rooted crop, so at bulb initiation stage (after 60 days) due to forecasted dry weather farmers are advised to go for irrigation in the moisture sensitive stage.**
- Cabbage and Cauliflower:**
- The transplanted cabbage is at seedling establishment to three week stage. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by **diamond back moth**. To control this apply Spinosad 45SC @ 0.5ml/l.
- Brinjal:**
- Shoot and fruit borer and leaf eating Caterpillar** may observe in brinjal field, to control it spray Emamectin benzoate 5% SG @ 0.4 g/litre of water or Chlorantraniliprole 18.5% SC @ 0.4 ml/litre of water. Spraying should be done during the early morning or evening hours.
- Tomato:**

Due to decrease in minimum temperature and foggy condition there may be chances of late blight in tomato, to avoid late blight precaution must be taken during the initial infection stage. Spray 2.5-gram Metalaxyl + Mancozeb/lit of water. Mix 3 gram Trichoderma with per kg of seed during seed treatment. Avoid excess nitrogen fertilizer and close spacing of plants.

Animal husbandry:

In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-353
Date- 02/12/2022

AGROMET ADVISORY BULLETIN FOR TANGI-CHOUDWAR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0mm rainfall during the last week. The mean daily maximum and minimum temperature was 29.2°C (-0.6) and 17°C (1.33), respectively. The mean relative humidity was 66-74%.

Forecasted period: Today(8.30IST) to 07th December(8.30IST)2022

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 28-30°C and night temperature will be 19-20°C for upcoming 5 days. Maximum and minimum relative humidity will be 53-61% and 32-44% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Tangi-Choudwar (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	02/12/2022	03/12/2022	04/12/2022	05/12/2022	06/12/2022
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	30	29	30	30	28
T-Min(°C)	20	20	20	19	19
Cloud Cover(octa)	0	0	5	1	8
RH Max (%)	61	53	58	58	53
RH Min (%)	44	35	37	32	34
Wind Speed (Kmph)	12	11	12	15	16
Wind Direction (deg)	45	60	60	27	27

Agromet Advisory

General Advisory: Due to forecasted low night temperature coupled with fog during upcoming days, there will be chance of infestation of sucking pests, leaf blight, powdery mildew in the rabi crops and vegetables, monitor the crops and vegetables, if symptoms are noticed, undertake plant protection measures. Provide light irrigation to standing vegetables and crops during morning hours to prevent damage from fog, low temperature and moisture stress. Carry out weeding, hoeing and other intercultural operations in rabi vegetables.

Kharif Rice:

Paddy is at milk dough stage to maturity stage. Drainout filed before 10-15 days of harvesting. **As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop.** For prevention of false smut spray the field before panicle initiation or flowering with propiconazole 25%EC @ 2ml/liter of water.

PULSES:

21-26°C soil temperature is required for proper germination of pulses. Due to forecast of dry weather for optimum germination irrigate the field after sowing. This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture.

Mustard:

Taking the advantage of available soil moisture, land preparation and sowing of the crop should be started to get optimum yield. **Apply irrigation in field if mustard was sown before 15-20 days. Due to increase in day temperature there may be infestation of saw fly and painted bug in Mustard, to control bug spray Dimethote 30 EC @ 1ml in liter of water and to control Saw fly spray imidachloprid 0.3 ml /liter of water.**

VEGETABLES:

As dry weather was prevailed in last 4 weeks and also forecasted for upcoming 5 days, farmers should irrigate their vegetable field in morning and evening hrs.

- Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. There is a chance of occurrence of **thrip, mites and Aphids** in Brinjal, Tomato, Chilli, cole crops, for control spray imidachloprid 0.3 ml /liter or fipronil 2 ml /liter of water.
- As Onion is shallow rooted crop, so at bulb initiation stage (after 60 days) due to forecasted dry weather farmers are advised to go for irrigation in the moisture sensitive stage.**
- Cabbage and Cauliflower:**
- The transplanted cabbage is at seedling establishment to three week stage. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by **diamond back moth**. To control this apply Spinosad 45SC @ 0.5ml/l.
- Brinjal:**
- Shoot and fruit borer and leaf eating Caterpillar** may observe in brinjal field, to control it spray Emamectin benzoate 5% SG @ 0.4 g/litre of water or Chlorantraniliprole 18.5% SC @ 0.4 ml/litre of water. Spraying should be done during the early morning or evening hours.

Tomato:

Due to decrease in minimum temperature and foggy condition there may be chances of late blight in tomato, to avoid late blight precaution must be taken during the initial infection stage. Spray 2.5-gram Metalaxyl + Mancozeb/lit of water. Mix 3 gram Trichoderma with per kg of seed during seed treatment. Avoid excess nitrogen fertilizer and close spacing of plants.

Animal husbandry:

- In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-353
Date- 02/12/2022

AGROMET ADVISORY BULLETIN FOR TIGIRIA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 0mm rainfall during the last week. The mean daily maximum and minimum temperature was 29.2°C (-0.6) and 17°C (1.33), respectively. The mean relative humidity was 66-74%.

Forecasted period: Today(8.30IST) to 07th December (8.30IST)2022

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 29-31°C and night temperature will be 17-18°C for upcoming 5 days. Maximum and minimum relative humidity will be 55-82% and 24-33% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Tigiria (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	02/12/2022	03/12/2022	04/12/2022	05/12/2022	06/12/2022
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	30	30	31	30	29
T-Min(°C)	18	18	18	18	17
Cloud Cover(octa)	0	0	1	2	7
RH Max (%)	82	63	67	63	55
RH Min (%)	33	32	30	31	24
Wind Speed (Kmph)	5	5	5	6	6
Wind Direction (deg)	11	72	19	25	11

Agro met Advisory

General Advisory: Due to forecasted low night temperature coupled with fog during upcoming days, there will be chance of infestation of sucking pests, leaf blight, powdery mildew in the rabi crops and vegetables, monitor the crops and vegetables, if symptoms are noticed, undertake plant protection measures. Provide light irrigation to standing vegetables and crops during morning hours to prevent damage from fog, low temperature and moisture stress. Carry out weeding, hoeing and other intercultural operations in rabi vegetables.

Kharif Rice:

Paddy is at milk dough stage to maturity stage. Drainout filed before 10-15 days of harvesting. As dry weather is forecasted for upcoming 5 days, in early varieties if panicle turns 85% yellow and matured, then go for harvesting of the crop. For prevention of false smut spray the field before panicle initiation or flowering with propiconazole 25%EC @ 2ml/liter of water.

PULSES:

21-26°C soil temperature is required for proper germination of pulses. Due to forecast of dry weather for optimum germination irrigate the field after sowing. This is the time for sowing of pre-rabi Black gram (Biri) and Green gram (Mung). Grow high yielding varieties like T 9, PU 11, PU 19 and PU 30 for black gram; and K 851, PDM 11, PDM 54, OUM 115 for green gram. Use 10 kg seeds/acre. Before sowing, seeds are to be treated with the Rhizobium culture.

Mustard:

Taking the advantage of available soil moisture, land preparation and sowing of the crop should be started to get optimum yield. Apply irrigation in field if mustard was sown before 15-20 days. Due to increase in day temperature there may be infestation of saw fly and painted bug in Mustard, to control bug spray Dimethote 30 EC @ 1ml in liter of water and to control Saw fly spray imidachloprid 0.3 ml /liter of water.

VEGETABLES:

As dry weather was prevailed in last 4 weeks and also forecasted for upcoming 5 days, farmers should irrigate their vegetable field in morning and evening hrs.

- Go for early planting of cole crops like raddish, cabbage, cauliflower, beans, cluster bean and tomato. There is a chance of occurrence of thrip, mites and Aphids in Brinjal, Tomato, Chilli, cole crops, for control spray imidachloprid 0.3 ml /liter or fipronil 2 ml /liter of water.
- As Onion is shallow rooted crop, so at bulb initiation stage (after 60 days) due to forecasted dry weather farmers are advised to go for irrigation in the moisture sensitive stage.
- Cabbage and Cauliflower:**
- The transplanted cabbage is at seedling establishment to three week stage. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by diamond back moth. To control this apply Spinosad 45SC @ 0.5ml/l.
- Brinjal:**
- Shoot and fruit borer and leaf eating Caterpillar may observe in brinjal field, to control it spray Emamectin benzoate 5% SG @ 0.4 g/litre of water or Chlorantraniliprole 18.5% SC @ 0.4 ml/litre of water. Spraying should be done during the early morning or evening hours.
- Tomato:**
- Due to decrease in minimum temperature and foggy condition there may be chances of late blight in tomato, to avoid late blight precaution must be taken during the initial infection stage. Spray 2.5-gram Metalaxyl + Mancozeb/lit of water. Mix 3 gram Trichoderma with per kg of seed during seed treatment. Avoid excess nitrogen fertilizer and close spacing of plants.

Animal husbandry:

- In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-353
Date- 02/12/2022

				
MEGHDOOT APP	DAMINI APP	FACEBOOK (Damu Cuttack)	BLOCK SPECIFIC WHAT'S APP GROUPS	@DAMUCUTTACK

- Kindly install and use **“Meghdoot mobile App”** for Cuttack district weather forecast, warnings and weather based agromet advisory services.
<https://play.google.com/store/apps/details?id=com.aas.meghdoot>
- Kindly install and use **“Damini mobile App”** for location specific for lightening forecast and warnings.
<https://play.google.com/store/apps/details?id=com.lightening.live.damini>
- Follow the **block specific whats app groups** and join in your block wise groups for getting weather based agromet advisory services twice in a week (Tuesday and Friday) and other weather warning based farm management advisories.
- Follow the **Official Website links** to get the weather based agromet advisory service bulletins (**English & Odia**) twice in a week (Tuesday and Friday).

1	ICAR-NRRI, Cuttack, Odisha	https://icar-nrri.in/agromate/	
2	Agrimet division, IMD, Pune	https://imdagrmet.gov.in/imdproject/AGIndex.php	

Dr Sujata Sethy
 (Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427