




Cuttack District Block Wise Agro-met Advisory Service Bulletin

*Friday, 06th January, 2023
(For the period 06th to 10th January, 2023)*



*Issued by
District Agromet Unit,
KrishiVigyan Kendra Cuttack*





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-363
Date- 06/01/2023

STANDARD PRECIPITATION INDEX AND NDVI MAP

Standard Precipitation Index (SPI)	Normalized difference vegetation Index (NDVI)
<p>Standardised Precipitation Index for the period 8th December 2022 to 4th January 2023</p> <p>MINISTRY OF EARTH SCIENCES INDIA METEOROLOGICAL DEPARTMENT HYDROMET SECTION, CRS PUNE</p> <p>STANDARDIZED PRECIPITATION INDEX FOR THE PERIOD 08 DEC 2022 TO 04 JAN 2023</p> <p>SPI VALUES</p> <ul style="list-style-type: none"> <= -2 (Extremely Dry) -1.50 to -1.00 (Moderately Dry) -1.0 to -0.50 (Moderately Dry) 0 to -0.50 (Moderately Dry) 0 to 0.50 (Moderately Wet) 0.50 to 1.00 (Moderately Wet) 1.50 to 2.00 (Extremely Wet) >= 2 (Extremely Wet) No Data 	<p>NOAA/VIIIRS/BLENDED NDVI Composite ending on Week no 52 (24.12.2022 to 30.12.2022) over Agricultural region</p> <p align="center">Odisha</p> <ul style="list-style-type: none"> -0.9999 - 0 0 - 0.05 0.05 - 0.1 0.1 - 0.15 0.15 - 0.2 0.2 - 0.25 0.25 - 0.3 0.3 - 0.35 0.35 - 0.4 0.4 - 0.45 0.45 - 0.5 0.5 - 0.55 0.55 - 0.6 > 0.6 <p>Bare soil, rocks clouds, rain, snow, etc.</p> <p>Moderate</p> <p>Good</p> <p>Very good</p> <p>Agriculture vigour is good over isolated places of the state.</p>
<p>As per SPI value (cumulative 4weeks -08 December2022 to 4 January 2023), mildly wet condition was there in Cuttack district.</p>	<p>According to NDVI map (24.12.2022 to 30.12.2022) the agriculture vigor was light over maximum eastern blocksand light to moderate western blocks of Cuttack district.</p>

LEGEND (Medium range block level rainfall forecast)		
	No Rainfall	00 mm
	Very light rainfall	01-05 mm
	Light rainfall	06-10 mm
	Light to moderate rainfall	11-30 mm
	Moderate to heavy rainfall	31-65 mm
	Heavy to very heavy rainfall	66-115 mm
	Extremely heavy rainfall	>116 mm

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com




Bulletin No.-363
Date- 06/01/2023

Past week (30 Dec, 2022 – 05 Jan, 2023) cumulative rainfall departure% from normal and (31 Dec, 2022 – 06 Jan, 2023) average minimum temperature and its departure from normal for Cuttack district



Medium range (Today (8.30IST) to Wednesday (8.30IST)) Rainfall (mm) forecast for Cuttack district



Dr Sujata Sethy
 (Nodal officer)



**Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area**



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427


















GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-363
Date- 06/01/2023

Weather (Past and Forecasted) based Disease and pest in Major Crops and vegetables

Sl No	Crop	Stage and condition of the crop	Anticipated Disease and pest	
1	Pulses (Green gram and black gram)	Vegetative stage to flowering		Aphid
				Black headed caterpillar
				Powdery mildew
				YMV
2	Mustard	Vegetative stage to flowering		Aphid
				Saw fly
				Leaf webber
				Stink Bug
3	Rabi Groundnut	Vegetative stage to flowering		Leaf Miner
				Aphid and Mites
4	Vegetables (Tomato, Chilli, Brinjal)	Vegetative to fruiting		Thrips
				Mites
				Chili Aphid
5	Brinjal	Fruiting		Fruit and shoot borer
6	Tomato	Vegetative		White fly

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427












GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT

KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-363
Date- 06/01/2023

				Leaf Curl virus
		Fruiting		Late blight
7	Potato	Sprouting/Vegetative stage		Leaf Blight
				Aphid
8	Cole crops (Cauliflower and cabbage)	Vegetative to fruiting		Leaf eating caterpillar
				Aphid
				Fruit borer
				Diamond back moth
				Tobacco Caterpillar

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-363
Date- 06/01/2023

AGROMET ADVISORY BULLETIN FOR ATHAGAD

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 28.4°C (+0.5 from normal) and 16.8°C (+3.4 from normal), respectively. The mean relative humidity was 69-81%.

Forecasted period: Today(8.30IST) to 11th January (8.30IST)2023

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 26-29°C and night temperature will be 10-12°C for upcoming 5 days. Maximum and minimum relative humidity will be 44-67% and 15-31% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Athagad (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	06/01/2023	07/01/2023	08/01/2023	09/01/2023	10/01/2023
Rainfall (mm)	0	0	0	0	0
T-Max (°C)	28	26	27	28	29
T-Min (°C)	11	10	11	11	12
Cloud Cover (octa)	0	0	0	0	0
RH Max (%)	59	45	49	44	67
RH Min (%)	31	15	16	15	15
Wind Speed (kmph)	8	8	6	7	7
Wind Direction (deg)	338	339	345	72	60

Agromet Advisory

General Advisory: Due to likely fall in minimum temperature in upcoming 4-5 days, Provide light irrigation to standing vegetables and crops during morning hours to prevent damage from moisture and low temperature stress and farmers should pause sowing of rabi pulses and oil seeds for upcoming 5 days. Due to low night temperature coupled with fog during the past few days, there is chance of infestation of sucking pests, leaf blight, powdery mildew in the rabi crops and vegetables, monitor the crops and vegetables, if symptoms are noticed, undertake plant protection measures.

Pulses:

In green gram/black gram, chances of infestation of leaf eating caterpillar and black headed caterpillar likely to occur. To control them, spray mixture of 2ml profenofos 50%EC or 2 ml quinalphos 25%EC or 1 ml dichlorvos 76%EC per litre of water @200 litres per acre during afternoon. Cloudy weather may aggravate the infestation of aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew.

Rabi-Groundnut:

Farmers are advised to do intercultivation and apply Gypsum @200 kg/ acre in Groundnut which is at flowering stage. To manage leaf miner spray Neem Based Pesticide (Azadirachtin) 300 PPM @ 1-litre /acre during primary stage of infestation. To manage chemically spray Profenophos 50% EC @ 400 ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 -litre of water. To manage aphids and thrips spray Thiamethoxam 25% W.G @ 40 gram/acre or Imidacloprid 17.8% SL @ 50-ml/acre by mixing it in 200 litre of water.

Mustard:

The mustard is at 15 to 20 days stage. Go for hoeing and weeding. Thin out the plants and maintain optimum plant population. There are chances of Aphid and stink bug infestation in early sown mustard crop. To manage aphids at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids and stink bugs chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly and leaf Webber spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water.

• **Vegetables:**

• **Thrips and mites** are prominent pests in vegetables at present. To control both, spray Bifenthrin@1g/litre or Profenophos@2ml/litre of water. To control the sucking pests like aphid & white fly, spray Imidacloprid @ 5ml/10 lit of water.

• Current weather is favourable for attack of Cabbage webber and tobacco caterpillar in cole crops. To control these insects, go for alternate spray of Neem based formulation 1500ppm@600ml/acre and Emamectin benzoate@70g/acre in 200 lit water. Leaf eating caterpillar and fruit borer may infest cabbage, cauliflower and tomato. Spray fipronil 5% SL 400ml or emamectinbenzoate 5% SG 80gm or indoxacarb14.5% SC mixing in 200lit. of water per acre crop. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by diamond back moth. To control this apply Spinosad 45SC @ 0.5ml/l.

• **POTATO:** There are chances of Leaf Blight disease incidence in potato crop. To manage early leaf blight in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre. To manage late blight in potato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days interval.

• **Onion:** The transplanted seedlings are at seedling establishment to four week stage in main field. Irrigate the crop at 10 days interval. Go for hoeing, weeding and 1 st top dressing with 45kg urea per acre crop area at 4 week stage of crop followed by irrigation.

Animal husbandry:

As the minimum temperature will continue to remain low for five days, the poultry birds should be protected from cold injury using electric bulb. Animals should be given mineral mixture along with salt regularly and wheat grains, jaggery etc. may be provided @ 10%- 20% in the daily ration during winter season to meet the energy requirement of the animals.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-363
Date- 06/01/2023

AGROMET ADVISORY BULLETIN FOR BADAMBA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 28.4°C (+0.5 from normal) and 16.8°C (+3.4 from normal), respectively. The mean relative humidity was 69-81%.

Forecasted period: Today(8.30IST) to 11th January (8.30IST)2023

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 26-30°C and night temperature will be 10-12°C for upcoming 5 days. Maximum and minimum relative humidity will be 41-68% and 14-31% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Badamba (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	06/01/2023	07/01/2023	08/01/2023	09/01/2023	10/01/2023
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	27	26	27	28	30
T-Min(°C)	11	10	11	11	12
Cloud Cover(octa)	0	0	0	0	0
RH Max (%)	63	41	46	45	68
RH Min (%)	31	14	15	15	14
Wind Speed (Kmph)	7	8	6	6	6
Wind Direction (deg)	340	338	345	72	360

Agromet Advisory

General Advisory: Due to likely fall in minimum temperature in upcoming 4-5 days, Provide light irrigation to standing vegetables and crops during morning hours to prevent damage from moisture and low temperature stress and farmers should pause sowing of rabi pulses and oil seeds for upcoming 5 days. Due to low night temperature coupled with fog during the past few days, there is chance of infestation of sucking pests, leaf blight, powdery mildew in the rabi crops and vegetables, monitor the crops and vegetables, if symptoms are noticed, undertake plant protection measures.

Pulses:

In green gram/black gram, chances of infestation of leaf eating caterpillar and black headed caterpillar likely to occur. To control them, spray mixture of 2ml profenofos 50%EC or 2 ml quinalphos 25%EC or 1 ml dichlorvos 76%EC per litre of water @200 litres per acre during afternoon. Cloudy weather may aggravate the infestation of aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew.

Rabi-Groundnut:

Farmers are advised to do intercultivation and apply Gypsum @200 kg/ acre in Groundnut which is at flowering stage. To manage leaf miner spray Neem Based Pesticide (Azadirachtin) 300 PPM @ 1-litre /acre during primary stage of infestation. To manage chemically spray Profenophos 50% EC @ 400 ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 -litre of water. To manage aphids and thrips spray Thiamethoxam 25% W.G @ 40 gram/acre or Imidacloprid 17.8% SL @ 50-ml/acre by mixing it in 200 litre of water.

Mustard:

The mustard is at 15 to 20 days stage. Go for hoeing and weeding. Thin out the plants and maintain optimum plant population. There are chances of Aphid and stink bug infestation in early sown mustard crop. To manage aphids at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids and stink bugs chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly and leaf Webber spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water.

- Vegetables:**
- Thrips and mites** are prominent pests in vegetables at present. To control both, spray Bifenthrin@1g/litre or Profenophos@2ml/litre of water. To control the sucking pests like aphid & white fly, spray Imidacloprid @ 5ml/10 lit of water.
- Current weather is favourable for attack of **Cabbage webber** and **tobacco caterpillar** in cole crops. To control these insects, go for alternate spray of Neem based formulation 1500ppm@600ml/acre and Emamectin benzoate@70g/acre in 200 lit water. **Leaf eating caterpillar and fruit borer** may infest cabbage, cauliflower and tomato. Spray fipronil 5% SL 400ml or emamectinbenzoate 5% SG 80gm or indoxacarb 14.5% SC mixing in 200lit. of water per acre crop. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by **diamond back moth**. To control this apply Spinosad 45SC @ 0.5ml/l.
- POTATO:** There are chances of **Leaf Blight** disease incidence in potato crop. To manage **early leaf blight** in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre. To manage **late blight** in potato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days interval.
- Onion:** The transplanted seedlings are at seedling establishment to four week stage in main field. Irrigate the crop at 10 days interval. Go for hoeing, weeding and 1 st top dressing with 45kg urea per acre crop area at 4 week stage of crop followed by irrigation.

Animal husbandry:

As the minimum temperature will continue to remain low for five days, the poultry birds should be protected from cold injury using electric bulb. Animals should be given mineral mixture along with salt regularly and wheat grains, jaggery etc. may be provided @ 10%- 20% in the daily ration during winter season to meet the energy requirement of the animals.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-363
Date- 06/01/2023

AGROMET ADVISORY BULLETIN FOR BANKI

District Past week average weather condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 28.4°C (+0.5 from normal) and 16.8°C (+3.4 from normal), respectively. The mean relative humidity was 69-81%.

Forecasted period: Today(8.30IST) to 11th January (8.30IST)2023

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 27-30°C and night temperature will be 11-12°C for upcoming 5 days. Maximum and minimum relative humidity will be 42-71% and 14-31% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Banki (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	06/01/2023	07/01/2023	08/01/2023	09/01/2023	10/01/2023
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	28	27	27	28	30
T-Min(°C)	11	11	11	12	12
Cloud Cover(octa)	0	0	0	0	0
RH Max (%)	61	42	48	44	71
RH Min (%)	31	14	15	15	14
Wind Speed (Kmph)	8	8	7	7	6
Wind Direction (deg)	345	338	345	72	60

Agromet Advisory

General Advisory: Due to likely fall in minimum temperature in upcoming 4-5 days, Provide light irrigation to standing vegetables and crops during morning hours to prevent damage from moisture and low temperature stress and farmers should pause sowing of rabi pulses and oil seeds for upcoming 5 days. Due to low night temperature coupled with fog during the past few days, there is chance of infestation of sucking pests, leaf blight, powdery mildew in the rabi crops and vegetables, monitor the crops and vegetables, if symptoms are noticed, undertake plant protection measures.

Pulses:

In green gram/black gram, chances of infestation of leaf eating caterpillar and black headed caterpillar likely to occur. To control them, spray mixture of 2ml profenofos 50%EC or 2 ml quinalphos 25%EC or 1 ml dichlorvos 76%EC per litre of water @200 litres per acre during afternoon. Cloudy weather may aggravate the infestation of aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew.

Rabi-Groundnut:

Farmers are advised to do intercultivation and apply Gypsum @200 kg/ acre in Groundnut which is at flowering stage. To manage leaf miner spray Neem Based Pesticide (Azadirachtin) 300 PPM @ 1-litre /acre during primary stage of infestation. To manage chemically spray Profenophos 50% EC @ 400 ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 -litre of water. To manage aphids and thrips spray Thiamethoxam 25% W.G @ 40 gram/acre or Imidacloprid 17.8% SL @ 50-ml/acre by mixing it in 200 litre of water.

Mustard:

The mustard is at 15 to 20 days stage. Go for hoeing and weeding. Thin out the plants and maintain optimum plant population. There are chances of Aphid and stink bug infestation in early sown mustard crop. To manage aphids at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids and stink bugs chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly and leaf Webber spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water.

- Vegetables:**
- Thrips and mites** are prominent pests in vegetables at present. To control both, spray Bifenthrin@1g/litre or Profenophos@2ml/litre of water. To control the sucking pests like aphid & white fly, spray Imidacloprid @ 5ml/10 lit of water.
- Current weather is favourable for attack of Cabbage webber and tobacco caterpillar in cole crops. To control these insects, go for alternate spray of Neem based formulation 1500ppm@600ml/acre and Emamectin benzoate@70g/acre in 200 lit water. Leaf eating caterpillar and fruit borer may infest cabbage, cauliflower and tomato. Spray fipronil 5% SL 400ml or emamectinbenzoate 5% SG 80gm or indoxacarb 14.5% SC mixing in 200lit. of water per acre crop. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by diamond back moth. To control this apply Spinosad 45SC @ 0.5ml/l.
- POTATO:** There are chances of Leaf Blight disease incidence in potato crop. To manage early leaf blight in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre. To manage late blight in potato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days interval.
- Onion:** The transplanted seedlings are at seedling establishment to four week stage in main field. Irrigate the crop at 10 days interval. Go for hoeing, weeding and 1 st top dressing with 45kg urea per acre crop area at 4 week stage of crop followed by irrigation.

Animal husbandry:

- As the minimum temperature will continue to remain low for five days, the poultry birds should be protected from cold injury using electric bulb. Animals should be given mineral mixture along with salt regularly and wheat grains, jaggery etc. may be provided @ 10%- 20% in the daily ration during winter season to meet the energy requirement of the animals.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-363
Date- 06/01/2023

AGROMET ADVISORY BULLETIN FOR BANKI-DAMPADA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 28.4°C (+0.5 from normal) and 16.8°C (+3.4 from normal), respectively. The mean relative humidity was 69-81%.

Forecasted period: Today(8.30IST) to 11th January (8.30IST)2023

Weather Advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 27-30°C and night temperature will be 11-12°C for upcoming 5 days. Maximum and minimum relative humidity will be 42-70% and 14-31% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Banki-Dampada (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	06/01/2023	07/01/2023	08/01/2023	09/01/2023	10/01/2023
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	27	27	27	28	30
T-Min(°C)	11	11	11	12	12
Cloud Cover(octa)	0	0	0	0	0
RH Max (%)	61	42	49	44	70
RH Min (%)	31	14	15	15	14
Wind Speed (Kmph)	8	8	7	7	6
Wind Direction (deg)	345	338	345	72	60

Agromet Advisory

General Advisory: Due to likely fall in minimum temperature in upcoming 4-5 days, Provide light irrigation to standing vegetables and crops during morning hours to prevent damage from moisture and low temperature stress and farmers should pause sowing of rabi pulses and oil seeds for upcoming 5 days. Due to low night temperature coupled with fog during the past few days, there is chance of infestation of sucking pests, leaf blight, powdery mildew in the rabi crops and vegetables, monitor the crops and vegetables, if symptoms are noticed, undertake plant protection measures.

Pulses:

In green gram/black gram, chances of infestation of leaf eating caterpillar and black headed caterpillar likely to occur. To control them, spray mixture of 2ml profenofos 50%EC or 2 ml quinalphos 25%EC or 1 ml dichlorvos 76%EC per litre of water @200 litres per acre during afternoon. Cloudy weather may aggravate the infestation of aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew.

Rabi-Groundnut:

Farmers are advised to do intercultivation and apply Gypsum @200 kg/ acre in Groundnut which is at flowering stage. To manage leaf miner spray Neem Based Pesticide (Azadirachtin) 300 PPM @ 1-litre /acre during primary stage of infestation. To manage chemically spray Profenophos 50% EC @ 400 ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 -litre of water. To manage aphids and thrips spray Thiamethoxam 25% W.G @ 40 gram/acre or Imidacloprid 17.8% SL @ 50-ml/acre by mixing it in 200 litre of water.

Mustard:

The mustard is at 15 to 20 days stage. Go for hoeing and weeding. Thin out the plants and maintain optimum plant population. There are chances of Aphid and stink bug infestation in early sown mustard crop. To manage aphids at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids and stink bugs chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly and leaf Webber spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water.

- Vegetables:**
- Thrips and mites** are prominent pests in vegetables at present. To control both, spray Bifenthrin@1g/litre or Profenophos@2ml/litre of water. To control the sucking pests like aphid & white fly, spray Imidacloprid @ 5ml/10 lit of water.
- Current weather is favourable for attack of Cabbage webber and tobacco caterpillar in cole crops. To control these insects, go for alternate spray of Neem based formulation 1500ppm@600ml/acre and Emamectin benzoate@70g/acre in 200 lit water. Leaf eating caterpillar and fruit borer may infest cabbage, cauliflower and tomato. Spray fipronil 5% SL 400ml or emamectinbenzoate 5% SG 80gm or indoxacarb14.5% SC mixing in 200lit. of water per acre crop. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by diamond back moth. To control this apply Spinosad 45SC @ 0.5ml/l.
- POTATO:** There are chances of Leaf Blight disease incidence in potato crop. To manage early leaf blight in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre. To manage late blight in potato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days interval.
- Onion:** The transplanted seedlings are at seedling establishment to four week stage in main field. Irrigate the crop at 10 days interval. Go for hoeing, weeding and 1 st top dressing with 45kg urea per acre crop area at 4 week stage of crop followed by irrigation.

Animal husbandry:

As the minimum temperature will continue to remain low for five days, the poultry birds should be protected from cold injury using electric bulb. Animals should be given mineral mixture along with salt regularly and wheat grains, jaggery etc. may be provided @ 10%- 20% in the daily ration during winter season to meet the energy requirement of the animals.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-363
Date- 06/01/2023

AGROMET ADVISORY BULLETIN FOR BARANGA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 28.4°C (+0.5 from normal) and 16.8°C (+3.4 from normal), respectively. The mean relative humidity was 69-81%.

Forecasted period: Today(8.30IST) to 11th January (8.30IST)2023

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 26-29°C and night temperature will be 12-14°C for upcoming 5 days. Maximum and minimum relative humidity will be 40-62% and 15-32% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Baranga (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	06/01/2023	07/01/2023	08/01/2023	09/01/2023	10/01/2023
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	28	26	27	28	29
T-Min(°C)	13	12	12	13	14
Cloud Cover	0	0	0	0	0
RH Max (%)	50	42	47	40	62
RH Min (%)	32	16	16	16	15
Wind Speed (Kmph)	9	8	8	8	8
Wind Direction (deg)	360	341	15	360	360

Agromet Advisory

General Advisory: Due to likely fall in minimum temperature in upcoming 4-5 days, Provide light irrigation to standing vegetables and crops during morning hours to prevent damage from moisture and low temperature stress and farmers should pause sowing of rabi pulses and oil seeds for upcoming 5 days. Due to low night temperature coupled with fog during the past few days, there is chance of infestation of sucking pests, leaf blight, powdery mildew in the rabi crops and vegetables, monitor the crops and vegetables, if symptoms are noticed, undertake plant protection measures.

Pulses:

In green gram/black gram, chances of infestation of leaf eating caterpillar and black headed caterpillar likely to occur. To control them, spray mixture of 2ml profenofos 50%EC or 2 ml quinalphos 25%EC or 1 ml dichlorvos 76%EC per litre of water @200 litres per acre during afternoon. Cloudy weather may aggravate the infestation of aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew.

Rabi-Groundnut:

Farmers are advised to do intercultivation and apply Gypsum @200 kg/ acre in Groundnut which is at flowering stage. To manage leaf miner spray Neem Based Pesticide (Azadirachtin) 300 PPM @ 1-litre /acre during primary stage of infestation. To manage chemically spray Profenophos 50% EC @ 400 ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 -litre of water. To manage aphids and thrips spray Thiamethoxam 25% W.G @ 40 gram/acre or Imidacloprid 17.8% SL @ 50-ml/acre by mixing it in 200 litre of water.

Mustard:

The mustard is at 15 to 20 days stage. Go for hoeing and weeding. Thin out the plants and maintain optimum plant population. There are chances of Aphid and stink bug infestation in early sown mustard crop. To manage aphids at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids and stink bugs chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly and leaf Webber spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water.

- Vegetables:**
- Thrips and mites** are prominent pests in vegetables at present. To control both, spray Bifenthrin@1g/litre or Profenophos@2ml/litre of water. To control the sucking pests like aphid & white fly, spray Imidacloprid @ 5ml/10 lit of water.
- Current weather is favourable for attack of Cabbage webber and tobacco caterpillar in cole crops. To control these insects, go for alternate spray of Neem based formulation 1500ppm@600ml/acre and Emamectin benzoate@70g/acre in 200 lit water. Leaf eating caterpillar and fruit borer may infest cabbage, cauliflower and tomato. Spray fipronil 5% SL 400ml or emamectinbenzoate 5% SG 80gm or indoxacarb 14.5% SC mixing in 200lit. of water per acre crop. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by diamond back moth. To control this apply Spinosad 45SC @ 0.5ml/l.
- POTATO:** There are chances of Leaf Blight disease incidence in potato crop. To manage early leaf blight in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre. To manage late blight in potato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days interval.
- Onion:** The transplanted seedlings are at seedling establishment to four week stage in main field. Irrigate the crop at 10 days interval. Go for hoeing, weeding and 1 st top dressing with 45kg urea per acre crop area at 4 week stage of crop followed by irrigation.

Animal husbandry:

- As the minimum temperature will continue to remain low for five days, the poultry birds should be protected from cold injury using electric bulb. Animals should be given mineral mixture along with salt regularly and wheat grains, jaggery etc. may be provided @ 10%- 20% in the daily ration during winter season to meet the energy requirement of the animals.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-363
Date- 06/01/2023

AGROMET ADVISORY BULLETIN FOR CUTTACK SADAR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 28.4°C (+0.5 from normal) and 16.8°C (+3.4 from normal), respectively. The mean relative humidity was 69-81%.

Forecasted period: Today(8.30IST) to 11th January (8.30IST)2023

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 26-29°C and night temperature will be 12-14°C for upcoming 5 days. Maximum and minimum relative humidity will be 43-60% and 15-31% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Cuttack sadar (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	06/01/2023	07/01/2023	08/01/2023	09/01/2023	10/01/2023
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	28	26	27	28	29
T-Min(°C)	13	12	12	13	14
Cloud Cover(octa)	0	0	0	0	0
RH Max (%)	52	44	48	43	60
RH Min (%)	31	17	17	16	15
Wind Speed (Kmph)	9	7	7	8	7
Wind Direction (deg)	345	341	15	360	360

Agromet Advisory

General Advisory: Due to likely fall in minimum temperature in upcoming 4-5 days, Provide light irrigation to standing vegetables and crops during morning hours to prevent damage from moisture and low temperature stress and farmers should pause sowing of rabi pulses and oil seeds for upcoming 5 days. Due to low night temperature coupled with fog during the past few days, there is chance of infestation of sucking pests, leaf blight, powdery mildew in the rabi crops and vegetables, monitor the crops and vegetables, if symptoms are noticed, undertake plant protection measures.

Pulses:

In green gram/black gram, chances of infestation of leaf eating caterpillar and black headed caterpillar likely to occur. To control them, spray mixture of 2ml profenofos 50%EC or 2 ml quinalphos 25%EC or 1 ml dichlorvos 76%EC per litre of water @200 litres per acre during afternoon. Cloudy weather may aggravate the infestation of aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew.

Rabi-Groundnut:

Farmers are advised to do intercultivation and apply Gypsum @200 kg/ acre in Groundnut which is at flowering stage. To manage leaf miner spray Neem Based Pesticide (Azadirachtin) 300 PPM @ 1-litre /acre during primary stage of infestation. To manage chemically spray Profenophos 50% EC @ 400 ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 -litre of water. To manage aphids and thrips spray Thiamethoxam 25% W.G @ 40 gram/acre or Imidacloprid 17.8% SL @ 50-ml/acre by mixing it in 200 litre of water.

Mustard:

The mustard is at 15 to 20 days stage. Go for hoeing and weeding. Thin out the plants and maintain optimum plant population. There are chances of Aphid and stink bug infestation in early sown mustard crop. To manage aphids at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids and stink bugs chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly and leaf Webber spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water.

Vegetables:

- Thrips and mites are prominent pests in vegetables at present. To control both, spray Bifenthrin@1g/litre or Profenophos@2ml/litre of water. To control the sucking pests like aphid & white fly, spray Imidacloprid @ 5ml/10 lit of water.
- Current weather is favourable for attack of Cabbage webber and tobacco caterpillar in cole crops. To control these insects, go for alternate spray of Neem based formulation 1500ppm@600ml/acre and Emamectin benzoate@70g/acre in 200 lit water. Leaf eating caterpillar and fruit borer may infest cabbage, cauliflower and tomato. Spray fipronil 5% SL 400ml or emamectinbenzoate 5% SG 80gm or indoxacarb 14.5% SC mixing in 200lit. of water per acre crop. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by diamond back moth. To control this apply Spinosad 45SC @ 0.5ml/l.
- POTATO:** There are chances of Leaf Blight disease incidence in potato crop. To manage early leaf blight in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre. To manage late blight in potato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days interval.
- Onion:** The transplanted seedlings are at seedling establishment to four week stage in main field. Irrigate the crop at 10 days interval. Go for hoeing, weeding and 1 st top dressing with 45kg urea per acre crop area at 4 week stage of crop followed by irrigation.

Animal husbandry:

As the minimum temperature will continue to remain low for five days, the poultry birds should be protected from cold injury using electric bulb. Animals should be given mineral mixture along with salt regularly and wheat grains, jaggery etc. may be provided @ 10%- 20% in the daily ration during winter season to meet the energy requirement of the animals.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-363
Date- 06/01/2023

AGROMET ADVISORY BULLETIN FOR KANTAPADA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 28.4°C (+0.5 from normal) and 16.8°C (+3.4 from normal), respectively. The mean relative humidity was 69-81%.

Forecasted period: Today(8.30IST) to 11th January (8.30IST)2023

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 26-29°C and night temperature will be 14-16°C for upcoming 5 days. Maximum and minimum relative humidity will be 40-59% and 16-31% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Kantapada (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	06/01/2023	07/01/2023	08/01/2023	09/01/2023	10/01/2023
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	28	26	27	28	29
T-Min(°C)	14	14	15	16	16
Cloud Cover(octa)	0	0	0	0	0
RH Max (%)	48	41	47	40	59
RH Min (%)	31	16	17	17	16
Wind Speed (Kmph)	10	9	9	9	8
Wind Direction (deg)	23	338	15	360	360

Agromet Advisory

General Advisory: Due to likely fall in minimum temperature in upcoming 4-5 days, Provide light irrigation to standing vegetables and crops during morning hours to prevent damage from moisture and low temperature stress and farmers should pause sowing of rabi pulses and oil seeds for upcoming 5 days. Due to low night temperature coupled with fog during the past few days, there is chance of infestation of sucking pests, leaf blight, powdery mildew in the rabi crops and vegetables, monitor the crops and vegetables, if symptoms are noticed, undertake plant protection measures.

Pulses:

In green gram/black gram, chances of infestation of leaf eating caterpillar and black headed caterpillar likely to occur. To control them, spray mixture of 2ml profenofos 50%EC or 2 ml quinalphos 25%EC or 1 ml dichlorvos 76%EC per litre of water @200 litres per acre during afternoon. Cloudy weather may aggravate the infestation of aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew.

Rabi-Groundnut:

Farmers are advised to do intercultivation and apply Gypsum @200 kg/ acre in Groundnut which is at flowering stage. To manage leaf miner spray Neem Based Pesticide (Azadirachtin) 300 PPM @ 1-litre /acre during primary stage of infestation. To manage chemically spray Profenophos 50% EC @ 400 ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 -litre of water. To manage aphids and thrips spray Thiamethoxam 25% W.G @ 40 gram/acre or Imidacloprid 17.8% SL @ 50-ml/acre by mixing it in 200 litre of water.

Mustard:

The mustard is at 15 to 20 days stage. Go for hoeing and weeding. Thin out the plants and maintain optimum plant population. There are chances of Aphid and stink bug infestation in early sown mustard crop. To manage aphids at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids and stink bugs chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly and leaf Webber spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water.

- Vegetables:**
- Thrips and mites** are prominent pests in vegetables at present. To control both, spray Bifenthrin@1g/litre or Profenophos@2ml/litre of water. To control the sucking pests like aphid & white fly, spray Imidacloprid @ 5ml/10 lit of water.
- Current weather is favourable for attack of **Cabbage webber** and **tobacco caterpillar** in cole crops. To control these insects, go for alternate spray of Neem based formulation 1500ppm@600ml/acre and Emamectin benzoate@70g/acre in 200 lit water. **Leaf eating caterpillar and fruit borer** may infest cabbage, cauliflower and tomato. Spray fipronil 5% SL 400ml or emamectinbenzoate 5% SG 80gm or indoxacarb14.5% SC mixing in 200lit. of water per acre crop. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by **diamond back moth**. To control this apply Spinosad 45SC @ 0.5ml/l.
- POTATO:** There are chances of **Leaf Blight** disease incidence in potato crop. To manage **early leaf blight** in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre. To manage **late blight** in potato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days interval.
- Onion:** The transplanted seedlings are at seedling establishment to four week stage in main field. Irrigate the crop at 10 days interval. Go for hoeing, weeding and 1 st top dressing with 45kg urea per acre crop area at 4 week stage of crop followed by irrigation.

Animal husbandry:

As the minimum temperature will continue to remain low for five days, the poultry birds should be protected from cold injury using electric bulb. Animals should be given mineral mixture along with salt regularly and wheat grains, jaggery etc. may be provided @ 10%- 20% in the daily ration during winter season to meet the energy requirement of the animals.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-363
Date- 06/01/2023

AGROMET ADVISORY BULLETIN FOR MAHANGA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 28.4°C (+0.5 from normal) and 16.8°C (+3.4 from normal), respectively. The mean relative humidity was 69-81%.

Forecasted period: Today(8.30IST) to 11th January (8.30IST)2023

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be and it will be 25-29°C and night temperature will be 12-13°C for upcoming 5 days. Maximum and minimum relative humidity will be 40-47% and 17-32% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Mahanga (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	06/01/2023	07/01/2023	08/01/2023	09/01/2023	10/01/2023
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	27	25	27	28	29
T-Min(°C)	12	12	12	13	13
Cloud Cover(octal)	0	0	0	0	0
RH Max (%)	43	41	43	40	47
RH Min (%)	32	19	19	18	17
Wind Speed (Kmph)	10	7	8	9	8
Wind Direction (deg)	360	338	360	23	360

Agromet Advisory

General Advisory: Due to likely fall in minimum temperature in upcoming 4-5 days, Provide light irrigation to standing vegetables and crops during morning hours to prevent damage from moisture and low temperature stress and farmers should pause sowing of rabi pulses and oil seeds for upcoming 5 days. Due to low night temperature coupled with fog during the past few days, there is chance of infestation of sucking pests, leaf blight, powdery mildew in the rabi crops and vegetables, monitor the crops and vegetables, if symptoms are noticed, undertake plant protection measures.

Pulses:

In green gram/black gram, chances of infestation of leaf eating caterpillar and black headed caterpillar likely to occur. To control them, spray mixture of 2ml profenofos 50%EC or 2 ml quinalphos 25%EC or 1 ml dichlorvos 76%EC per litre of water @200 litres per acre during afternoon. Cloudy weather may aggravate the infestation of aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew.

Rabi-Groundnut:

Farmers are advised to do intercultivation and apply Gypsum @200 kg/ acre in Groundnut which is at flowering stage. To manage leaf miner spray Neem Based Pesticide (Azadirachtin) 300 PPM @ 1-litre /acre during primary stage of infestation. To manage chemically spray Profenophos 50% EC @ 400 ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 -litre of water. To manage aphids and thrips spray Thiamethoxam 25% W.G @ 40 gram/acre or Imidacloprid 17.8% SL @ 50-ml/acre by mixing it in 200 litre of water.

Mustard:

The mustard is at 15 to 20 days stage. Go for hoeing and weeding. Thin out the plants and maintain optimum plant population. There are chances of Aphid and stink bug infestation in early sown mustard crop. To manage aphids at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids and stink bugs chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly and leaf Webber spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water.

- Vegetables:**
- Thrips and mites** are prominent pests in vegetables at present. To control both, spray Bifenthrin@1g/litre or Profenophos@2ml/litre of water. To control the sucking pests like aphid & white fly, spray Imidacloprid @ 5ml/10 lit of water.
- Current weather is favourable for attack of **Cabbage webber** and **tobacco caterpillar** in cole crops. To control these insects, go for alternate spray of Neem based formulation 1500ppm@600ml/acre and Emamectin benzoate@70g/acre in 200 lit water. **Leaf eating caterpillar and fruit borer** may infest cabbage, cauliflower and tomato. Spray fipronil 5% SL 400ml or emamectinbenzoate 5% SG 80gm or indoxacarb14.5% SC mixing in 200lit. of water per acre crop. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by **diamond back moth**. To control this apply Spinosad 45SC @ 0.5ml/l.
- POTATO:** There are chances of **Leaf Blight** disease incidence in potato crop. To manage **early leaf blight** in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre. To manage **late blight** in potato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days interval.
- Onion:** The transplanted seedlings are at seedling establishment to four week stage in main field. Irrigate the crop at 10 days interval. Go for hoeing, weeding and 1 st top dressing with 45kg urea per acre crop area at 4 week stage of crop followed by irrigation.

Animal husbandry:

As the minimum temperature will continue to remain low for five days, the poultry birds should be protected from cold injury using electric bulb. Animals should be given mineral mixture along with salt regularly and wheat grains, jaggery etc. may be provided @ 10%- 20% in the daily ration during winter season to meet the energy requirement of the animals.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-363
Date- 06/01/2023

AGROMET ADVISORY BULLETIN FOR NARASINGHPUR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 28.4°C (+0.5 from normal) and 16.8°C (+3.4 from normal), respectively. The mean relative humidity was 69-81%.

Forecasted period: Today(8.30IST) to 11th January (8.30IST)2023

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 26-30°C and night temperature will be 10-12°C for upcoming 5 days. Maximum and minimum relative humidity will be 39-65% and 14-32% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Narasinghpur (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	06/01/2023	07/01/2023	08/01/2023	09/01/2023	10/01/2023
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	26	26	27	29	30
T-Min(°C)	11	10	11	11	12
Cloud Cover(octa)	0	0	0	0	0
RH Max (%)	65	39	43	44	57
RH Min (%)	32	14	15	15	15
Wind Speed (Kmph)	6	7	5	4	4
Wind Direction (deg)	340	336	23	71	68

Agromet Advisory

General Advisory: Due to likely fall in minimum temperature in upcoming 4-5 days, Provide light irrigation to standing vegetables and crops during morning hours to prevent damage from moisture and low temperature stress and farmers should pause sowing of rabi pulses and oil seeds for upcoming 5 days. Due to low night temperature coupled with fog during the past few days, there is chance of infestation of sucking pests, leaf blight, powdery mildew in the rabi crops and vegetables, monitor the crops and vegetables, if symptoms are noticed, undertake plant protection measures.

Pulses:

In green gram/black gram, chances of infestation of leaf eating caterpillar and black headed caterpillar likely to occur. To control them, spray mixture of 2ml profenofos 50%EC or 2 ml quinalphos 25%EC or 1 ml dichlorvos 76%EC per litre of water @200 litres per acre during afternoon. Cloudy weather may aggravate the infestation of aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew.

Rabi-Groundnut:

Farmers are advised to do intercultivation and apply Gypsum @200 kg/ acre in Groundnut which is at flowering stage. To manage leaf miner spray Neem Based Pesticide (Azadirachtin) 300 PPM @ 1-litre /acre during primary stage of infestation. To manage chemically spray Profenophos 50% EC @ 400 ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 -litre of water. To manage aphids and thrips spray Thiamethoxam 25% W.G @ 40 gram/acre or Imidacloprid 17.8% SL @ 50-ml/acre by mixing it in 200 litre of water.

Mustard:

The mustard is at 15 to 20 days stage. Go for hoeing and weeding. Thin out the plants and maintain optimum plant population. There are chances of Aphid and stink bug infestation in early sown mustard crop. To manage aphids at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids and stink bugs chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly and leaf Webber spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water.

- Vegetables:**
- Thrips and mites** are prominent pests in vegetables at present. To control both, spray Bifenthrin@1g/litre or Profenophos@2ml/litre of water. To control the sucking pests like aphid & white fly, spray Imidacloprid @ 5ml/10 lit of water.
- Current weather is favourable for attack of **Cabbage webber** and **tobacco caterpillar** in cole crops. To control these insects, go for alternate spray of Neem based formulation 1500ppm@600ml/acre and Emamectin benzoate@70g/acre in 200 lit water. **Leaf eating caterpillar and fruit borer** may infest cabbage, cauliflower and tomato. Spray fipronil 5% SL 400ml or emamectinbenzoate 5% SG 80gm or indoxacarb 14.5% SC mixing in 200lit. of water per acre crop. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by **diamond back moth**. To control this apply Spinosad 45SC @ 0.5ml/l.
- POTATO:** There are chances of **Leaf Blight** disease incidence in potato crop. To manage **early leaf blight** in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre. To manage **late blight** in potato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days interval.
- Onion:** The transplanted seedlings are at seedling establishment to four week stage in main field. Irrigate the crop at 10 days interval. Go for hoeing, weeding and 1 st top dressing with 45kg urea per acre crop area at 4 week stage of crop followed by irrigation.

Animal husbandry:

- As the minimum temperature will continue to remain low for five days**, the poultry birds should be protected from cold injury using electric bulb. Animals should be given mineral mixture along with salt regularly and wheat grains, jaggery etc. may be provided @ 10%- 20% in the daily ration during winter season to meet the energy requirement of the animals.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-363
Date- 06/01/2023

AGROMET ADVISORY BULLETIN FOR NIALI

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 28.4°C (+0.5 from normal) and 16.8°C (+3.4 from normal), respectively. The mean relative humidity was 69-81%.

Forecasted period: Today(8.30IST) to 11th January (8.30IST)2023

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 27-29°C and night temperature will be 13-14°C for upcoming 5 days. Maximum and minimum relative humidity will be 43-61% and 16-32% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Niali (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	06/01/2023	07/01/2023	08/01/2023	09/01/2023	10/01/2023
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	28	27	27	28	29
T-Min(°C)	14	13	13	14	14
Cloud Cover(octa)	0	0	0	0	0
RH Max (%)	50	43	50	43	61
RH Min (%)	32	16	17	18	17
Wind Speed(Kmph)	11	9	10	10	9
Wind Direction(deg)	27	338	360	56	105

Agromet Advisory

General Advisory: Due to likely fall in minimum temperature in upcoming 4-5 days, Provide light irrigation to standing vegetables and crops during morning hours to prevent damage from moisture and low temperature stress and farmers should pause sowing of rabi pulses and oil seeds for upcoming 5 days. Due to low night temperature coupled with fog during the past few days, there is chance of infestation of sucking pests, leaf blight, powdery mildew in the rabi crops and vegetables, monitor the crops and vegetables, if symptoms are noticed, undertake plant protection measures.

Pulses:

In green gram/black gram, chances of infestation of leaf eating caterpillar and black headed caterpillar likely to occur. To control them, spray mixture of 2ml profenofos 50%EC or 2 ml quinalphos 25%EC or 1 ml dichlorvos 76%EC per litre of water @200 litres per acre during afternoon. Cloudy weather may aggravate the infestation of aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew.

Rabi-Groundnut:

Farmers are advised to do intercultivation and apply Gypsum @200 kg/ acre in Groundnut which is at flowering stage. To manage leaf miner spray Neem Based Pesticide (Azadirachtin) 300 PPM @ 1-litre /acre during primary stage of infestation. To manage chemically spray Profenophos 50% EC @ 400 ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 -litre of water. To manage aphids and thrips spray Thiamethoxam 25% W.G @ 40 gram/acre or Imidacloprid 17.8% SL @ 50-ml/acre by mixing it in 200 litre of water.

Mustard:

The mustard is at 15 to 20 days stage. Go for hoeing and weeding. Thin out the plants and maintain optimum plant population. There are chances of Aphid and stink bug infestation in early sown mustard crop. To manage aphids at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids and stink bugs chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly and leaf Webber spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water.

- Vegetables:**
- Thrips and mites** are prominent pests in vegetables at present. To control both, spray Bifenthrin@1g/litre or Profenophos@2ml/litre of water. To control the sucking pests like aphid & white fly, spray Imidacloprid @ 5ml/10 lit of water.
- Current weather is favourable for attack of Cabbage webber and tobacco caterpillar in cole crops. To control these insects, go for alternate spray of Neem based formulation 1500ppm@600ml/acre and Emamectin benzoate@70g/acre in 200 lit water. Leaf eating caterpillar and fruit borer may infest cabbage, cauliflower and tomato. Spray fipronil 5% SL 400ml or emamectinbenzoate 5% SG 80gm or indoxacarb14.5% SC mixing in 200lit. of water per acre crop. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by diamond back moth. To control this apply Spinosad 45SC @ 0.5ml/l.
- POTATO:** There are chances of Leaf Blight disease incidence in potato crop. To manage early leaf blight in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre. To manage late blight in potato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days interval.
- Onion:** The transplanted seedlings are at seedling establishment to four week stage in main field. Irrigate the crop at 10 days interval. Go for hoeing, weeding and 1 st top dressing with 45kg urea per acre crop area at 4 week stage of crop followed by irrigation.

Animal husbandry:

- As the minimum temperature will continue to remain low for five days, the poultry birds should be protected from cold injury using electric bulb. Animals should be given mineral mixture along with salt regularly and wheat grains, jaggery etc. may be provided @ 10%- 20% in the daily ration during winter season to meet the energy requirement of the animals.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-363
Date- 06/01/2023

AGROMET ADVISORY BULLETIN FOR NISCHINTAKOILI
District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 28.4°C (+0.5 from normal) and 16.8°C (+3.4 from normal), respectively. The mean relative humidity was 69-81%.

Forecasted period: Today(8.30IST) to 11th January (8.30IST)2023

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 26-29°C and night temperature will be 14-16°C for upcoming 5 days. Maximum and minimum relative humidity will be 39-48% and 17-32% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Nischintakoili (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	06/01/2023	07/01/2023	08/01/2023	09/01/2023	10/01/2023
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	27	26	27	28	29
T-Min(°C)	14	14	15	16	16
Cloud Cover(octa)	0	0	0	0	0
RH Max (%)	43	41	43	39	48
RH Min (%)	32	19	19	18	17
Wind Speed (Kmph)	10	8	9	9	8
Wind Direction (deg)	360	338	360	23	360

Agromet Advisory

General Advisory: Due to likely fall in minimum temperature in upcoming 4-5 days, Provide light irrigation to standing vegetables and crops during morning hours to prevent damage from moisture and low temperature stress and farmers should pause sowing of rabi pulses and oil seeds for upcoming 5 days. Due to low night temperature coupled with fog during the past few days, there is chance of infestation of sucking pests, leaf blight, powdery mildew in the rabi crops and vegetables, monitor the crops and vegetables, if symptoms are noticed, undertake plant protection measures.

Pulses:

In green gram/black gram, chances of infestation of leaf eating caterpillar and black headed caterpillar likely to occur. To control them, spray mixture of 2ml profenofos 50%EC or 2 ml quinalphos 25%EC or 1 ml dichlorvos 76%EC per litre of water @200 litres per acre during afternoon. Cloudy weather may aggravate the infestation of aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew.

Rabi-Groundnut:

Farmers are advised to do intercultivation and apply Gypsum @200 kg/ acre in Groundnut which is at flowering stage. To manage leaf miner spray Neem Based Pesticide (Azadirachtin) 300 PPM @ 1-litre /acre during primary stage of infestation. To manage chemically spray Profenophos 50% EC @ 400 ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 -litre of water. To manage aphids and thrips spray Thiamethoxam 25% W.G @ 40 gram/acre or Imidacloprid 17.8% SL @ 50-ml/acre by mixing it in 200 litre of water.

Mustard:

The mustard is at 15 to 20 days stage. Go for hoeing and weeding. Thin out the plants and maintain optimum plant population. There are chances of Aphid and stink bug infestation in early sown mustard crop. To manage aphids at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids and stink bugs chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly and leaf Webber spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water.

- Vegetables:**
- Thrips and mites** are prominent pests in vegetables at present. To control both, spray Bifenthrin@1g/litre or Profenophos@2ml/litre of water. To control the sucking pests like aphid & white fly, spray Imidacloprid @ 5ml/10 lit of water.
- Current weather is favourable for attack of Cabbage webber and tobacco caterpillar in cole crops. To control these insects, go for alternate spray of Neem based formulation 1500ppm@600ml/acre and Emamectin benzoate@70g/acre in 200 lit water. Leaf eating caterpillar and fruit borer may infest cabbage, cauliflower and tomato. Spray fipronil 5% SL 400ml or emamectinbenzoate 5% SG 80gm or indoxacarb 14.5% SC mixing in 200lit. of water per acre crop. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by diamond back moth. To control this apply Spinosad 45SC @ 0.5ml/l.
- POTATO:** There are chances of Leaf Blight disease incidence in potato crop. To manage early leaf blight in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre. To manage late blight in potato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days interval.
- Onion:** The transplanted seedlings are at seedling establishment to four week stage in main field. Irrigate the crop at 10 days interval. Go for hoeing, weeding and 1 st top dressing with 45kg urea per acre crop area at 4 week stage of crop followed by irrigation.

Animal husbandry:

- As the minimum temperature will continue to remain low for five days, the poultry birds should be protected from cold injury using electric bulb. Animals should be given mineral mixture along with salt regularly and wheat grains, jaggery etc. may be provided @ 10%- 20% in the daily ration during winter season to meet the energy requirement of the animals.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-363
Date- 06/01/2023

AGROMET ADVISORY BULLETIN FOR SALEPUR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 28.4°C (+0.5 from normal) and 16.8°C (+3.4 from normal), respectively. The mean relative humidity was 69-81%.

Forecasted period: Today(8.30IST) to 11th January (8.30IST)2023

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 26-29°C and night temperature will be 14-16°C for upcoming 5 days. Maximum and minimum relative humidity will be 39-50% and 17-32% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Salepur (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	06/01/2023	07/01/2023	08/01/2023	09/01/2023	10/01/2023
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	27	26	27	28	29
T-Min(°C)	14	15	15	16	16
Cloud Cover(octa)	0	0	0	0	0
RH Max (%)	45	41	44	39	50
RH Min (%)	32	18	19	18	17
Wind Speed (Kmph)	10	7	8	9	8
Wind Direction (deg)	23	340	360	23	360

Agromet Advisory

General Advisory: Due to likely fall in minimum temperature in upcoming 4-5 days, Provide light irrigation to standing vegetables and crops during morning hours to prevent damage from moisture and low temperature stress and farmers should pause sowing of rabi pulses and oil seeds for upcoming 5 days. Due to low night temperature coupled with fog during the past few days, there is chance of infestation of sucking pests, leaf blight, powdery mildew in the rabi crops and vegetables, monitor the crops and vegetables, if symptoms are noticed, undertake plant protection measures.

Pulses:

In green gram/black gram, chances of infestation of leaf eating caterpillar and black headed caterpillar likely to occur. To control them, spray mixture of 2ml profenofos 50%EC or 2 ml quinalphos 25%EC or 1 ml dichlorvos 76%EC per litre of water @200 litres per acre during afternoon. Cloudy weather may aggravate the infestation of aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew.

Rabi-Groundnut:

Farmers are advised to do intercultivation and apply Gypsum @200 kg/ acre in Groundnut which is at flowering stage. To manage leaf miner spray Neem Based Pesticide (Azadirachtin) 300 PPM @ 1-litre /acre during primary stage of infestation. To manage chemically spray Profenophos 50% EC @ 400 ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 -litre of water. To manage aphids and thrips spray Thiamethoxam 25% W.G @ 40 gram/acre or Imidacloprid 17.8% SL @ 50-ml/acre by mixing it in 200 litre of water.

Mustard:

The mustard is at 15 to 20 days stage. Go for hoeing and weeding. Thin out the plants and maintain optimum plant population. There are chances of Aphid and stink bug infestation in early sown mustard crop. To manage aphids at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids and stink bugs chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly and leaf Webber spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water.

- Vegetables:**
- Thrips and mites** are prominent pests in vegetables at present. To control both, spray Bifenthrin@1g/litre or Profenophos@2ml/litre of water. To control the sucking pests like aphid & white fly, spray Imidacloprid @ 5ml/10 lit of water.
- Current weather is favourable for attack of **Cabbage webber** and **tobacco caterpillar** in cole crops. To control these insects, go for alternate spray of Neem based formulation 1500ppm@600ml/acre and Emamectin benzoate@70g/acre in 200 lit water. **Leaf eating caterpillar and fruit borer** may infest cabbage, cauliflower and tomato. Spray fipronil 5% SL 400ml or emamectinbenzoate 5% SG 80gm or indoxacarb14.5% SC mixing in 200lit. of water per acre crop. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by **diamond back moth**. To control this apply Spinosad 45SC @ 0.5ml/l.
- POTATO:** There are chances of **Leaf Blight** disease incidence in potato crop. To manage **early leaf blight** in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre. To manage **late blight** in potato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days interval.
- Onion:** The transplanted seedlings are at seedling establishment to four week stage in main field. Irrigate the crop at 10 days interval. Go for hoeing, weeding and 1 st top dressing with 45kg urea per acre crop area at 4 week stage of crop followed by irrigation.

Animal husbandry:

As the minimum temperature will continue to remain low for five days, the poultry birds should be protected from cold injury using electric bulb. Animals should be given mineral mixture along with salt regularly and wheat grains, jaggery etc. may be provided @ 10%- 20% in the daily ration during winter season to meet the energy requirement of the animals.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-363
Date- 06/01/2023

AGROMET ADVISORY BULLETIN FOR TANGI-CHOUDWAR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 28.4°C (+0.5 from normal) and 16.8°C (+3.4 from normal), respectively. The mean relative humidity was 69-81%.

Forecasted period: Today(8.30IST) to 11th January (8.30IST)2023

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 26-29°C and night temperature will be 12-13°C for upcoming 5 days. Maximum and minimum relative humidity will be 43-54% and 22-36% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Tangi-Choudwar (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	06/01/2023	07/01/2023	08/01/2023	09/01/2023	10/01/2023
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	27	26	27	28	29
T-Min(°C)	12	12	12	13	13
Cloud Cover(octa)	0	0	0	0	0
RH Max (%)	47	43	46	44	54
RH Min (%)	36	22	26	27	25
Wind Speed (Kmph)	17	11	12	12	11
Wind Direction (deg)	27	360	45	45	68

Agromet Advisory

General Advisory: Due to likely fall in minimum temperature in upcoming 4-5 days, Provide light irrigation to standing vegetables and crops during morning hours to prevent damage from moisture and low temperature stress and farmers should pause sowing of rabi pulses and oil seeds for upcoming 5 days. Due to low night temperature coupled with fog during the past few days, there is chance of infestation of sucking pests, leaf blight, powdery mildew in the rabi crops and vegetables, monitor the crops and vegetables, if symptoms are noticed, undertake plant protection measures.

Pulses:

In green gram/black gram, chances of infestation of leaf eating caterpillar and black headed caterpillar likely to occur. To control them, spray mixture of 2ml profenofos 50%EC or 2 ml quinalphos 25%EC or 1 ml dichlorvos 76%EC per litre of water @200 litres per acre during afternoon. Cloudy weather may aggravate the infestation of aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew.

Rabi-Groundnut:

Farmers are advised to do intercultivation and apply Gypsum @200 kg/ acre in Groundnut which is at flowering stage. To manage leaf miner spray Neem Based Pesticide (Azadirachtin) 300 PPM @ 1-litre /acre during primary stage of infestation. To manage chemically spray Profenophos 50% EC @ 400 ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 -litre of water. To manage aphids and thrips spray Thiamethoxam 25% W.G @ 40 gram/acre or Imidacloprid 17.8% SL @ 50-ml/acre by mixing it in 200 litre of water.

Mustard:

The mustard is at 15 to 20 days stage. Go for hoeing and weeding. Thin out the plants and maintain optimum plant population. There are chances of Aphid and stink bug infestation in early sown mustard crop. To manage aphids at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids and stink bugs chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly and leaf Webber spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water.

- Vegetables:**
- Thrips and mites** are prominent pests in vegetables at present. To control both, spray Bifenthrin@1g/litre or Profenophos@2ml/litre of water. To control the sucking pests like aphid & white fly, spray Imidacloprid @ 5ml/10 lit of water.
- Current weather is favourable for attack of **Cabbage webber** and **tobacco caterpillar** in cole crops. To control these insects, go for alternate spray of Neem based formulation 1500ppm@600ml/acre and Emamectin benzoate@70g/acre in 200 lit water. **Leaf eating caterpillar and fruit borer** may infest cabbage, cauliflower and tomato. Spray fipronil 5% SL 400ml or emamectinbenzoate 5% SG 80gm or indoxacarb14.5% SC mixing in 200lit. of water per acre crop. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by **diamond back moth**. To control this apply Spinosad 45SC @ 0.5ml/l.
- POTATO:** There are chances of **Leaf Blight** disease incidence in potato crop. To manage **early leaf blight** in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre. To manage **late blight** in potato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days interval.
- Onion:** The transplanted seedlings are at seedling establishment to four week stage in main field. Irrigate the crop at 10 days interval. Go for hoeing, weeding and 1 st top dressing with 45kg urea per acre crop area at 4 week stage of crop followed by irrigation.

Animal husbandry:

- As the minimum temperature will continue to remain low for five days**, the poultry birds should be protected from cold injury using electric bulb. Animals should be given mineral mixture along with salt regularly and wheat grains, jaggery etc. may be provided @ 10%- 20% in the daily ration during winter season to meet the energy requirement of the animals.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-363
Date- 06/01/2023

AGROMET ADVISORY BULLETIN FOR TIGIRIA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 28.4°C (+0.5 from normal) and 16.8°C (+3.4 from normal), respectively. The mean relative humidity was 69-81%.

Forecasted period: Today(8.30IST) to 11th January (8.30IST)2023

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 26-30°C and night temperature will be 10-12°C for upcoming 5 days. Maximum and minimum relative humidity will be 44-68% and 15-31% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Tigiria (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	06/01/2023	07/01/2023	08/01/2023	09/01/2023	10/01/2023
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	28	26	27	28	30
T-Min(°C)	11	10	11	11	12
Cloud Cover(octa)	0	0	0	0	0
RH Max (%)	61	44	48	44	68
RH Min (%)	31	15	16	15	15
Wind Speed (Kmph)	8	8	6	6	6
Wind Direction (deg)	343	338	345	72	23

Agro met Advisory

General Advisory: Due to likely fall in minimum temperature in upcoming 4-5 days, Provide light irrigation to standing vegetables and crops during morning hours to prevent damage from moisture and low temperature stress and farmers should pause sowing of rabi pulses and oil seeds for upcoming 5 days. Due to low night temperature coupled with fog during the past few days, there is chance of infestation of sucking pests, leaf blight, powdery mildew in the rabi crops and vegetables, monitor the crops and vegetables, if symptoms are noticed, undertake plant protection measures.

Pulses:

In green gram/black gram, chances of infestation of leaf eating caterpillar and black headed caterpillar likely to occur. To control them, spray mixture of 2ml profenofos 50%EC or 2 ml quinalphos 25%EC or 1 ml dichlorvos 76%EC per litre of water @200 litres per acre during afternoon. Cloudy weather may aggravate the infestation of aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew.

Rabi-Groundnut:

Farmers are advised to do intercultivation and apply Gypsum @200 kg/ acre in Groundnut which is at flowering stage. To manage leaf miner spray Neem Based Pesticide (Azadirachtin) 300 PPM @ 1-litre /acre during primary stage of infestation. To manage chemically spray Profenophos 50% EC @ 400 ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 -litre of water. To manage aphids and thrips spray Thiamethoxam 25% W.G @ 40 gram/acre or Imidacloprid 17.8% SL @ 50-ml/acre by mixing it in 200 litre of water.

Mustard:

The mustard is at 15 to 20 days stage. Go for hoeing and weeding. Thin out the plants and maintain optimum plant population. There are chances of Aphid and stink bug infestation in early sown mustard crop. To manage aphids at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids and stink bugs chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly and leaf Webber spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water.

- Vegetables:**
- Thrips and mites** are prominent pests in vegetables at present. To control both, spray Bifenthrin@1g/litre or Profenophos@2ml/litre of water. To control the sucking pests like aphid & white fly, spray Imidacloprid @ 5ml/10 lit of water.
- Current weather is favourable for attack of Cabbage webber and tobacco caterpillar in cole crops. To control these insects, go for alternate spray of Neem based formulation 1500ppm@600ml/acre and Emamectin benzoate@70g/acre in 200 lit water. Leaf eating caterpillar and fruit borer may infest cabbage, cauliflower and tomato. Spray fipronil 5% SL 400ml or emamectinbenzoate 5% SG 80gm or indoxacarb 14.5% SC mixing in 200lit. of water per acre crop. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Cole crops may get affected by diamond back moth. To control this apply Spinosad 45SC @ 0.5ml/l.
- POTATO:** There are chances of Leaf Blight disease incidence in potato crop. To manage early leaf blight in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre. To manage late blight in potato spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre at 10 days interval.
- Onion:** The transplanted seedlings are at seedling establishment to four week stage in main field. Irrigate the crop at 10 days interval. Go for hoeing, weeding and 1 st top dressing with 45kg urea per acre crop area at 4 week stage of crop followed by irrigation.

Animal husbandry:

- As the minimum temperature will continue to remain low for five days, the poultry birds should be protected from cold injury using electric bulb. Animals should be given mineral mixture along with salt regularly and wheat grains, jaggery etc. may be provided @ 10%- 20% in the daily ration during winter season to meet the energy requirement of the animals.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-363
Date- 06/01/2023

				
MEGHDOOT APP	DAMINI APP	FACEBOOK (Damu Cuttack)	BLOCK SPECIFIC WHAT'S APP GROUPS	@DAMUCUTTACK



- Kindly install and use **“Meghdoot mobile App”** for Cuttack district weather forecast, warnings and weather based agromet advisory services.

<https://play.google.com/store/apps/details?id=com.aas.meghdoot>

- Kindly install and use **“Damini mobile App”** for location specific for lightening forecast and warnings.

<https://play.google.com/store/apps/details?id=com.lightening.live.damini>

- Follow the **block specific whats app groups** and join in your block wise groups for getting weather based agromet advisory services twice in a week (Tuesday and Friday) and other weather warning based farm management advisories.
- Follow the **Official Website links** to get the weather based agromet advisory service bulletins (**English & Odia**) twice in a week (Tuesday and Friday).

1	ICAR-NRRI, Cuttack, Odisha	https://icar-nrri.in/agromate/	
2	Agromet division, IMD, Pune	https://imdagrmet.gov.in/imdproject/AGIndex.php	

Dr Sujata Sethy
 (Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427