




Cuttack District Block Wise Agro-met Advisory Service Bulletin

Tuesday, 21st February, 2023
(For the period 21st to 25th February, 2023)



Issued by
District Agromet Unit,
KrishiVigyan Kendra Cuttack






GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-376
Date- 21/02/2023

STANDARD PRECIPITATION INDEX AND NDVI MAP



LEGEND (Medium range block level rainfall forecast)		
	No Rainfall	00 mm
	Very light rainfall	01-05 mm
	Light rainfall	06-10 mm
	Light to moderate rainfall	11-30 mm
	Moderate to heavy rainfall	31-65 mm
	Heavy to very heavy rainfall	66-115 mm
	Extremely heavy rainfall	>116 mm

LEGEND (Past week temperature)		
Color code	Temperature	Departure
	Below Normal	< 2°C from normal
	Normal	-2°C to +2°C from normal
	Above Normal	>2°C from normal

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com




Bulletin No.-376
Date- 21/02/2023

Past week (14-20 Feb, 2023) cumulative rainfall departure% from normal and (15-21 Feb, 2023) average minimum temperature and its departure from normal for Cuttack district



Medium range (Today (8.30IST) to Sunday (8.30IST)) Rainfall (mm) forecast for Cuttack district



Dr Sujata Sethy
 (Nodal officer)



**Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area**




Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



EXTENDED RANGE RAINFALL FORECAST FOR UPCOMING 2 WEEKS


INDIA METEOROLOGICAL DEPARTMENT
NUMERICAL WEATHER PREDICTION DIVISION, NEW DELHI
STATE: ODISHA



PERIOD: WEEK-1 (17-02-2023 To 23-02-2023)



INDIA METEOROLOGICAL DEPARTMENT
NUMERICAL WEATHER PREDICTION DIVISION, NEW DELHI
STATE: ODISHA



PERIOD: WEEK-2 (24-02-2023 To 02-03-2023)



As per extended range outlook there will be dry weather is forecasted in first and 2nd week for Odisha condition. So Farmers are advised to make their arrangement of irrigation of rabi crops like Pulses and groundnut, Mustard especially at flowering stage and they may go for sowing activity with utilization of rain water. To conserve soil moisture they are advised to cover the crop with mulching material (straw material) and should add rotted FYM in soil.

















GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT

KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-376
Date- 21/02/2023

Weather (Past and Forecasted) based Disease and pest in Major Crops and vegetables

Sl No	Crop	Stage and condition of the crop	Anticipated Disease and pest	
1	Pulses (Green gram and black gram)	Flowering to pod development		YMV
2	Mustard	Flowering to maturity		Saw Fly
				Leaf webber
3	Rabi Groundnut	Flowering to maturity		Spodotera/Tobacco caterpillar
				Red hairy caterpillar
6	Tomato	Flowering to Fruiting		Bacterial Canker
				Sun burn
				Flower drop
9	Cucurbits	Flowering to Fruiting stage		Whitefly
				Late blight
				Tender fruit drop
10	Mango	Flowering to fruiting		Mango fruit drop

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT

KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-376
Date- 21/02/2023

11	Cashew	Flowering to fruiting		Tea mosquito bug
----	--------	-----------------------	--	------------------

Dr Sujata Sethy
(Nodal officer)



**Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area**



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-376
Date- 21/02/2023

AGROMET ADVISORY BULLETIN FOR ATHAGAD

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.8°C (-0.4 from normal) and 19.5°C (+1.13 from normal), respectively. The mean relative humidity was 53-79%.

Forecasted period: Today(8.30IST) to 26th February(8.30IST)2023

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 33-35°C and night temperature will be 20-22°C for upcoming 5 days. Maximum and minimum relative humidity will be 90-95% and 10-19% respectively. There will be clear sky for next 5 days.

Block	Athagad (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	21/02/2023	22/02/2023	23/02/2023	24/02/2023	25/02/2023
Rainfall (mm)	0	0	0	0	0
T-Max (°C)	33	33	34	35	35
T-Min (°C)	22	20	21	21	21
Cloud Cover (octa)	0	0	0	0	0
RH Max (%)	95	90	94	94	93
RH Min (%)	10	19	12	17	14
Wind Speed (kmph)	9	14	8	8	9
Wind Direction (deg)	199	206	68	154	169

Agromet Advisory

General Advisory: Provide light and frequent irrigation to standing vegetables and crops during morning and evening hours to prevent damage from moisture and high temperature stress.

- Foliar spray of 2%DAP +1% KCL (MOP) during critical stages of flowering and grain formation stage.
- Use of biofertilizer like Azotobacter, Azospirillum and phospho bacteria @10 packets (500g)/ha along with 25 kg of soil or FYM.
- To avoid flower and fruit drop spray planofix @ 1ml/4.5lit of water.
- Ridge and furrow irrigation should be practiced to conserve soil moisture in the main field.

Pulses:

November to December sown crop is at pod formation to seed development (nine to twelve week) stage. Flowering and pod development stage is sensitive to moisture stress; carry out irrigation as per the status of soil moisture. In case of YMV infestation uproots and destroys the infected plant and spray imidacloprid 50ml/acre mixing with 200 lit of water.

Rabi-Groundnut:

Do not disturb the soil after flower initiation in groundnut. Do irrigation in the field based upon soil moisture. Increase in temperature may enhance red hairy caterpillar and tobacco caterpillar infestation in groundnut; to control, spray Profenophos 50% EC @ 2 ml/litre of water or Flubendiamide 480 SC @ 0.2 ml/litre of water.

Mustard:

Rainfed mustard sown during October is at maturity stage. Harvest the mustard crop as soon as the pods turn yellowish-brown. Present weather may favour the infestation of saw fly, leaf Webber. To control saw fly, spray 400ml Endosulphan/acre, to control leaf webber spray 200 ml Ithofenprox or Deltamethrin per acre in 200 lit water.

Vegetables:

Tomato: Transplanted seedlings are at 6-8 week stage in main field. Due to water stress, deficiency of boron or due to hormonal disbalance flower drop occurs. Control: Apply timely and required amount of irrigation. Spray Boron 20% 2 gram/lit during early flowering and 15 days later. Due to increase in temperature there may be chances of bacterial canker and Sun burn disease in tomato at fruiting stage. To control bacterial canker Farmers should spray Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water. They should use shade net to protect the crop from sun burn.

Potato: The main crop is ready for harvest within 75-85 days of planting depending upon the soil type and variety sown in our climatic condition of Odisha. The crop is ready to harvest when majority of the leaves turn yellowish brown. Irrigation should be stopped 15 days before harvesting.

Onion: Do inter culture and hand weeding may be carried out at 45-50 DAT and application of 30-kg of urea and 15-kg of MOP fertilizer should be done at this stage. Onion being a shallow rooted crop, needs frequent light irrigation to maintain optimum soil moisture for proper growth and bulb development.

Fruits:

Mango: The present weather is appropriate for blossoming of flowers. Spray NAA @ 200 ppm per tree, with the help of ring method of irrigation apply 25 kg FYM, 500 gram of urea and 1 kg of potash and phosphate each.

Cashew: Prevailing conditions are suitable for the incidence of Tea mosquito bug. Farmers are advised to go for spraying of Neem oil @ 5ml/L + Monocrotophos @ 1.6ml/L at new blush stage and Chlorpyrifos @ 2ml/L at flowering stage. If inflorescence have flourished stop spraying of any insecticides. Hormones like NAA @ 2 ppm can be sprayed. FYM @ 25 kg per tree can be applied.

Mushroom:

Initiate cultivation of paddy straw mushroom. As hot weather condition is prevailed in Cuttack district, Spray water if the mushrooms are at fruiting stage. Cover the Mushroom unit with water soaked Gunny bags to maintain the inside temperature.

Animal husbandry:

As temperature is slightly increasing hence, it will impact physiological, metabolic behavior as well as egg production of poultry bird. Farmers should provide sufficient clean drinking water to birds and make proper shelter and feed management.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-376
Date- 21/02/2023

AGROMET ADVISORY BULLETIN FOR BADAMBA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.8°C (-0.4 from normal) and 19.5°C (+1.13 from normal), respectively. The mean relative humidity was 53-79%.

Forecasted period: Today(8.30IST) to 26th February(8.30IST)2023

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 36-37°C and night temperature will be 19-20°C for upcoming 5 days. Maximum and minimum relative humidity will be 87-96% and 11-17% respectively. There will be clear sky for next 5 days.

Block	Badamba (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	21/02/2023	22/02/2023	23/02/2023	24/02/2023	25/02/2023
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	36	36	36	36	37
T-Min(°C)	20	19	19	20	20
Cloud Cover(octa)	0	0	0	0	0
RH Max (%)	96	87	92	93	90
RH Min (%)	11	17	11	15	13
Wind Speed (Kmph)	8	12	7	7	9
Wind Direction (deg)	199	200	15	158	169

Agromet Advisory

General Advisory: Provide light and frequent irrigation to standing vegetables and crops during morning and evening hours to prevent damage from moisture and high temperature stress.

- Foliar spray of 2%DAP +1% KCL (MOP) during critical stages of flowering and grain formation stage.
- Use of biofertilizer like Azotobacter, Azospirillum and phospho bacteria @10 packets (500g)/ha along with 25 kg of soil or FYM.
- To avoid flower and fruit drop spray planofix @ 1ml/4.5lit of water.
- Ridge and furrow irrigation should be practiced to conserve soil moisture in the main field.

Pulses:

November to December sown crop is at pod formation to seed development (nine to twelve week) stage. Flowering and pod development stage is sensitive to moisture stress; carry out irrigation as per the status of soil moisture. In case of YMV infestation uproots and destroys the infected plant and spray imidacloprid 50ml/acre mixing with 200 lit of water.

Rabi-Groundnut:

Do not disturb the soil after flower initiation in groundnut. Do irrigation in the field based upon soil moisture. Increase in temperature may enhance red hairy caterpillar and tobacco caterpillar infestation in groundnut; to control, spray Profenophos 50% EC @ 2 ml/litre of water or Flubendiamide 480 SC @ 0.2 ml/litre of water.

Mustard:

Rained mustard sown during October is at maturity stage. Harvest the mustard crop as soon as the pods turn yellowish-brown. Present weather may favour the infestation of saw fly, leaf Webber. To control saw fly, spray 400ml Endosulphan/acre, to control leaf webber spray 200 ml Ithofenprox or Deltamethrin per acre in 200 lit water.

Vegetables:

Tomato: Transplanted seedlings are at 6-8 week stage in main field. Due to water stress, deficiency of boron or due to hormonal disbalance flower drop occurs. Control: Apply timely and required amount of irrigation. Spray Boron 20% 2 gram/lit during early flowering and 15 days later. Due to increase in temperature there may be chances of bacterial canker and Sun burn disease in tomato at fruiting stage. To control bacterial canker Darners should spray Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water. They should use shade net to protect the crop from sun burn.

Potato: The main crop is ready for harvest within 75-85 days of planting depending upon the soil type and variety sown in our climatic condition of Odisha. The crop is ready to harvest when majority of the leaves turn yellowish brown. Irrigation should be stopped 15 days before harvesting.

Onion: Do inter culture and hand weeding may be carried out at 45-50 DAT and application of 30-kg of urea and 15-kg of MOP fertilizer should be done at this stage. Onion being a shallow rooted crop, needs frequent light irrigation to maintain optimum soil moisture for proper growth and bulb development.

Fruits:

Mango: The present weather is appropriate for blossoming of flowers. Spray NAA @ 200 ppm per tree, with the help of ring method of irrigation apply 25 kg FYM, 500 gram of urea and 1 kg of potash and phosphate each.

Cashew: Prevailing conditions are suitable for the incidence of Tea mosquito bug. Farmers are advised to go for spraying of Neem oil @ 5ml/L + Monocrotophos @ 1.6ml/L at new blush stage and Chlorpyrifos @ 2ml/L at flowering stage. If inflorescence have flourished stop spraying of any insecticides. Hormones like NAA @ 2 ppm can be sprayed. FYM @ 25 kg per tree can be applied.

Mushroom:

Initiate cultivation of paddy straw mushroom. As hot weather condition is prevailed in Cuttack district, Spray water if the mushrooms are at fruiting stage. Cover the Mushroom unit with water soaked Gunny bags to maintain the inside temperature.

Animal husbandry:

As temperature is slightly increasing hence, it will impact physiological, metabolic behavior as well as egg production of poultry bird. Farmers should provide sufficient clean drinking water to birds and make proper shelter and feed management.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-376
Date- 21/02/2023

AGROMET ADVISORY BULLETIN FOR BANKI

District Past week average weather condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.8°C (-0.4 from normal) and 19.5°C (+1.13 from normal), respectively. The mean relative humidity was 53-79%.

Forecasted period: Today(8.30IST) to 26th February(8.30IST)2023

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 35-36°C and night temperature will be 20-21°C for upcoming 5 days. Maximum and minimum relative humidity will be 89-95% and 11-21% respectively. There will be clear sky for next 5 days.

Block	Banki (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	21/02/2023	22/02/2023	23/02/2023	24/02/2023	25/02/2023
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	36	36	36	35	35
T-Min(°C)	21	20	20	20	20
Cloud Cover(octa)	0	0	0	0	0
RH Max (%)	95	89	92	92	91
RH Min (%)	11	21	13	19	13
Wind Speed (Kmph)	9	14	9	9	10
Wind Direction (deg)	195	206	210	180	189

Agromet Advisory

General Advisory: Provide light and frequent irrigation to standing vegetables and crops during morning and evening hours to prevent damage from moisture and high temperature stress.

- Foliar spray of 2%DAP +1% KCL (MOP) during critical stages of flowering and grain formation stage.
- Use of biofertilizer like Azotobacter, Azospirillum and phospho bacteria @10 packets (500g)/ha along with 25 kg of soil or FYM.
- To avoid flower and fruit drop spray planofix @ 1ml/4.5lit of water.
- Ridge and furrow irrigation should be practiced to conserve soil moisture in the main field.

Pulses:

November to December sown crop is at pod formation to seed development (nine to twelve week) stage. Flowering and pod development stage is sensitive to moisture stress; carry out irrigation as per the status of soil moisture. In case of YMV infestation uproots and destroys the infected plant and spray imidacloprid 50ml/acre mixing with 200 lit of water.

Rabi-Groundnut:

Do not disturb the soil after flower initiation in groundnut. Do irrigation in the field based upon soil moisture. Increase in temperature may enhance red hairy caterpillar and tobacco caterpillar infestation in groundnut; to control, spray Profenophos 50% EC @ 2 ml/litre of water or Flubendiamide 480 SC @ 0.2 ml/litre of water.

Mustard:

Rainfed mustard sown during October is at maturity stage. Harvest the mustard crop as soon as the pods turn yellowish-brown. Present weather may favour the infestation of saw fly, leaf webber. To control saw fly, spray 400ml Endosulphan/acre, to control leaf webber spray 200 ml Ithofenprox or Deltamethrin per acre in 200 lit water.

Vegetables:

Tomato: Transplanted seedlings are at 6-8 week stage in main field. Due to water stress, deficiency of boron or due to hormonal disbalance flower drop occurs. Control: Apply timely and required amount of irrigation. Spray Boron 20% 2 gram/lit during early flowering and 15 days later. Due to increase in temperature there may be chances of bacterial canker and Sun burn disease in tomato at fruiting stage. To control bacterial canker farmers should spray Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water. They should use shade net to protect the crop from sun burn.

Potato: The main crop is ready for harvest within 75-85 days of planting depending upon the soil type and variety sown in our climatic condition of Odisha. The crop is ready to harvest when majority of the leaves turn yellowish brown. Irrigation should be stopped 15 days before harvesting.

Onion: Do inter culture and hand weeding may be carried out at 45-50 DAT and application of 30-kg of urea and 15-kg of MOP fertilizer should be done at this stage. Onion being a shallow rooted crop, needs frequent light irrigation to maintain optimum soil moisture for proper growth and bulb development.

Fruits:

Mango: The present weather is appropriate for blossoming of flowers. Spray NAA @ 200 ppm per tree, with the help of ring method of irrigation apply 25 kg FYM, 500 gram of urea and 1 kg of potash and phosphate each.

Cashew: Prevailing conditions are suitable for the incidence of Tea mosquito bug. Farmers are advised to go for spraying of Neem oil @ 5ml/L + Monocrotophos @ 1.6ml/L at new blush stage and Chlorpyrifos @ 2ml/L at flowering stage. If inflorescence have flourished stop spraying of any insecticides. Hormones like NAA @ 2 ppm can be sprayed. FYM @25 kg per tree can be applied.

Mushroom:

Initiate cultivation of paddy straw mushroom. As hot weather condition is prevailed in Cuttack district, Spray water if the mushrooms are at fruiting stage. Cover the Mushroom unit with water soaked Gunny bags to maintain the inside temperature.

Animal husbandry:

- As temperature is slightly increasing hence, it will impact physiological, metabolic behavior as well as egg production of poultry bird. Farmers should provide sufficient clean drinking water to birds and make proper shelter and feed management.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-376
Date- 21/02/2023

AGROMET ADVISORY BULLETIN FOR BANKI-DAMPADA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.8°C (-0.4 from normal) and 19.5°C (+1.13 from normal), respectively. The mean relative humidity was 53-79%.

Forecasted period: Today(8.30IST) to 26th February(8.30IST)2023

Weather Advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 35-36°C and night temperature will be 20-21°C for upcoming 5 days. Maximum and minimum relative humidity will be 88-95% and 11-21% respectively. There will be clear sky for next 5 days.

Block	Banki-Dampada (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	21/02/2023	22/02/2023	23/02/2023	24/02/2023	25/02/2023
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	36	35	36	35	35
T-Min(°C)	21	20	20	20	20
Cloud Cover(octa)	0	0	0	0	0
RH Max (%)	95	88	93	92	92
RH Min (%)	11	21	14	19	13
Wind Speed (Kmph)	10	14	9	9	10
Wind Direction (deg)	195	206	210	169	189

Agromet Advisory

General Advisory: Provide light and frequent irrigation to standing vegetables and crops during morning and evening hours to prevent damage from moisture and high temperature stress.

➤ **Foliar spray of 2%DAP +1% KCL (MOP) during critical stages of flowering and grain formation stage.**

➤ **Use of biofertilizer like Azotobacter, Azospirillum and phospho bacteria @10 packets (500g)/ha along with 25 kg of soil or FYM.**

➤ **To avoid flower and fruit drop spray planofix @ 1ml/4.5lit of water.**

➤ **Ridge and furrow irrigation should be practiced to conserve soil moisture in the main field.**

Pulses:

November to December sown crop is at pod formation to seed development (nine to twelve week) stage. **Flowering and pod development stage is sensitive to moisture stress; carry out irrigation as per the status of soil moisture.** In case of **YMV infestation** uproots and destroys the infected plant and spray imidacloprid 50ml/acre mixing with 200 lit of water.

Rabi-Groundnut:

Do not disturb the soil after flower initiation in groundnut. **Do irrigation in the field based upon soil moisture.** Increase in temperature may enhance **red hairy caterpillar and tobacco caterpillar** infestation in groundnut; to control, spray Profenophos 50% EC @ 2 ml/litre of water or Flubendiamide 480 SC @ 0.2 ml/litre of water.

Mustard:

Rainfed mustard sown during October is at maturity stage. Harvest the mustard crop as soon as the pods turn yellowish-brown. Present weather may favour the infestation of **saw fly, leaf Webber**. To control saw fly, spray 400ml Endosulphan/acre, to control leaf webber spray 200 ml Ithofenprox or Deltamethrin per acre in 200 litr water.

Vegetables:

Tomato: Transplanted seedlings are at 6-8 week stage in main field. **Due to water stress, deficiency of boron or due to hormonal disbalance flower drop occurs.** Control: Apply timely and required amount of irrigation. Spray Boron 20% 2 gram/lit during early flowering and 15 days later. Due to increase in temperature there may be chances of **bacterial canker and Sun burn** disease in tomato at fruiting stage. To control bacterial canker Farmers should spray Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water. They should use shade net to protect the crop from sun burn.

Potato: The main crop is ready for harvest within 75-85 days of planting depending upon the soil type and variety sown in our climatic condition of odisha. The crop is ready to harvest when majority of the leaves turn yellowish brown. Irrigation should be stopped 15 days before harvesting.

Onion: Do inter culture and hand weeding may be carried out at 45-50 DAT and application of 30-kg of urea and 15-kg of MOP fertilizer should be done at this stage. Onion being a shallow rooted crop, **needs frequent light irrigation to maintain optimum soil moisture for proper growth and bulb development.**

Fruits:

Mango: The present weather is appropriate for blossoming of flowers. Spray NAA @ 200 ppm per tree, **with the help of ring method of irrigation** apply 25 kg FYM, 500 gram of urea and 1 kg of potash and phosphate each.

Cashew: Prevailing conditions are suitable for the incidence of **Tea mosquito bug**. Farmers are advised to go for spraying of Neem oil @ 5ml/L + Monocrotophos @ 1.6ml/L at new blush stage and Chlorpyrifos @ 2ml/L at flowering stage. If inflorescence have flourished stop spraying of any insecticides. Hormones like NAA @ 2 ppm can be sprayed. FYM @25 kg per tree can be applied.

Mushroom:

Initiate cultivation of paddy straw mushroom. As hot weather condition is prevailed in Cuttack district, Spray water if the mushrooms are at fruiting stage. Cover the Mushroom unit with water soaked Gunny bags to maintain the inside temperature.

Animal husbandry:

- As temperature is slightly increasing hence, it will impact physiological, metabolic behavior as well as egg production of poultry bird. Farmers should provide sufficient clean drinking water to birds and make proper shelter and feed management.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-376
Date- 21/02/2023

AGROMET ADVISORY BULLETIN FOR BARANGA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.8°C (-0.4 from normal) and 19.5°C (+1.13 from normal), respectively. The mean relative humidity was 53-79%.

Forecasted period: Today(8.30IST) to 26th February(8.30IST)2023

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 34-36°C and night temperature will be 21-22°C for upcoming 5 days. Maximum and minimum relative humidity will be 90-95% and 13-26% respectively. There will be clear sky for next 5 days.

Block	Baranga (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	21/02/2023	22/02/2023	23/02/2023	24/02/2023	25/02/2023
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	36	35	35	34	35
T-Min(°C)	22	21	21	21	21
Cloud Cover	0	0	0	0	0
RH Max (%)	95	90	95	94	93
RH Min (%)	13	26	17	23	17
Wind Speed (Kmph)	11	16	11	10	10
Wind Direction (deg)	200	204	200	180	180

Agromet Advisory

General Advisory: Provide light and frequent irrigation to standing vegetables and crops during morning and evening hours to prevent damage from moisture and high temperature stress.

➤ **Foliar spray of 2%DAP +1% KCL (MOP) during critical stages of flowering and grain formation stage.**

➤ **Use of biofertilizer like Azotobacter, Azospirillum and phospho bacteria @10 packets (500g)/ha along with 25 kg of soil or FYM.**

➤ **To avoid flower and fruit drop spray planofix @ 1ml/4.5lit of water.**

➤ **Ridge and furrow irrigation should be practiced to conserve soil moisture in the main field.**

Pulses:

November to December sown crop is at pod formation to seed development (nine to twelve week) stage. **Flowering and pod development stage is sensitive to moisture stress; carry out irrigation as per the status of soil moisture.** In case of **YMV infestation** uproots and destroys the infected plant and spray imidacloprid 50ml/acre mixing with 200 lit of water.

Rabi-Groundnut:

Do not disturb the soil after flower initiation in groundnut. **Do irrigation in the field based upon soil moisture.** Increase in temperature may enhance **red hairy caterpillar and tobacco caterpillar** infestation in groundnut; to control, spray Profenophos 50% EC @ 2 ml/litre of water or Flubendiamide 480 SC @ 0.2 ml/litre of water.

Mustard:

Rainfed mustard sown during October is at maturity stage. Harvest the mustard crop as soon as the pods turn yellowish-brown. Present weather may favour the infestation of **saw fly, leaf Webber**. To control saw fly, spray 400ml Endosulphan/acre, to control leaf webber spray 200 ml Ithofenprox or Deltamethrin per acre in 200 lit water.

Vegetables:

Tomato: Transplanted seedlings are at 6-8 week stage in main field. **Due to water stress, deficiency of boron or due to hormonal disbalance flower drop occurs.** Control: Apply timely and required amount of irrigation. Spray Boron 20% 2 gram/lit during early flowering and 15 days later. Due to increase in temperature there may be chances of **bacterial canker and Sun burn** disease in tomato at fruiting stage. To control bacterial canker farmers should spray Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water. They should use shade net to protect the crop from sun burn.

Potato: The main crop is ready for harvest within 75-85 days of planting depending upon the soil type and variety sown in our climatic condition of Odisha. The crop is ready to harvest when majority of the leaves turn yellowish brown. Irrigation should be stopped 15 days before harvesting.

Onion: Do inter culture and hand weeding may be carried out at 45-50 DAT and application of 30-kg of urea and 15-kg of MOP fertilizer should be done at this stage. Onion being a shallow rooted crop, **needs frequent light irrigation to maintain optimum soil moisture for proper growth and bulb development.**

Fruits:

Mango: The present weather is appropriate for blossoming of flowers. Spray NAA @ 200 ppm per tree, **with the help of ring method of irrigation** apply 25 kg FYM, 500 gram of urea and 1 kg of potash and phosphate each.

Cashew: Prevailing conditions are suitable for the incidence of **Tea mosquito bug**. Farmers are advised to go for spraying of Neem oil @ 5ml/L + Monocrotophos @ 1.6ml/L at new blush stage and Chlorpyrifos @ 2ml/L at flowering stage. If inflorescence have flourished stop spraying of any insecticides. Hormones like NAA @ 2 ppm can be sprayed. FYM @ 25 kg per tree can be applied.

Mushroom:

Initiate cultivation of paddy straw mushroom. As hot weather condition is prevailed in Cuttack district, Spray water if the mushrooms are at fruiting stage. Cover the Mushroom unit with water soaked Gunny bags to maintain the inside temperature.

Animal husbandry:

As temperature is slightly increasing hence, it will impact physiological, metabolic behavior as well as egg production of poultry bird. Farmers should provide sufficient clean drinking water to birds and make proper shelter and feed management.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-376
Date- 21/02/2023

AGROMET ADVISORY BULLETIN FOR CUTTACK SADAR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.8°C (-0.4 from normal) and 19.5°C (+1.13 from normal), respectively. The mean relative humidity was 53-79%.

Forecasted period: Today(8.30IST) to 26th February(8.30IST)2023

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 34-35°C and night temperature will be 21-22°C for upcoming 5 days. Maximum and minimum relative humidity will be 91-95% and 11-23% respectively. There will be clear sky for next 5 days.

Block	Cuttack sadar (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	21/02/2023	22/02/2023	23/02/2023	24/02/2023	25/02/2023
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	34	34	35	35	3
T-Min(°C)	22	21	21	22	21
Cloud Cover(octa)	0	0	0	0	0
RH Max (%)	95	91	95	95	93
RH Min (%)	11	23	14	20	16
Wind Speed (Kmph)	11	15	10	9	9
Wind Direction (deg)	200	207	207	169	189

Agromet Advisory

General Advisory: Provide light and frequent irrigation to standing vegetables and crops during morning and evening hours to prevent damage from moisture and high temperature stress.

➤ **Foliar spray of 2%DAP +1% KCL (MOP) during critical stages of flowering and grain formation stage.**

➤ **Use of biofertilizer like Azotobacter, Azospirillum and phospho bacteria @10 packets (500g)/ha along with 25 kg of soil or FYM.**

➤ **To avoid flower and fruit drop spray planofix @ 1ml/4.5lit of water.**

➤ **Ridge and furrow irrigation should be practiced to conserve soil moisture in the main field.**

Pulses:

November to December sown crop is at pod formation to seed development (nine to twelve week) stage. **Flowering and pod development stage is sensitive to moisture stress; carry out irrigation as per the status of soil moisture.** In case of **YMV infestation** uproots and destroys the infected plant and spray imidacloprid 50ml/acre mixing with 200 lit of water.

Rabi-Groundnut:

Do not disturb the soil after flower initiation in groundnut. **Do irrigation in the field based upon soil moisture.** Increase in temperature may enhance **red hairy caterpillar and tobacco caterpillar** infestation in groundnut; to control, spray Profenophos 50% EC @ 2 ml/litre of water or Flubendiamide 480 SC @ 0.2 ml/litre of water.

Mustard:

Rainfed mustard sown during October is at maturity stage. Harvest the mustard crop as soon as the pods turn yellowish-brown. Present weather may favour the infestation of **saw fly, leaf Webber**. To control saw fly, spray 400ml Endosulphan/acre, to control leaf webber spray 200 ml Ithofenprox or Deltamethrin per acre in 200 lit water.

Vegetables:

Tomato: Transplanted seedlings are at 6-8 week stage in main field. **Due to water stress, deficiency of boron or due to hormonal disbalance flower drop occurs.** Control: Apply timely and required amount of irrigation. Spray Boron 20% 2 gram/lit during early flowering and 15 days later. Due to increase in temperature there may be chances of **bacterial canker and Sun burn** disease in tomato at fruiting stage. To control bacterial canker farmers should spray Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water. They should use shade net to protect the crop from sun burn.

Potato: The main crop is ready for harvest within 75-85 days of planting depending upon the soil type and variety sown in our climatic condition of Odisha. The crop is ready to harvest when majority of the leaves turn yellowish brown. Irrigation should be stopped 15 days before harvesting.

Onion: Do inter culture and hand weeding may be carried out at 45-50 DAT and application of 30-kg of urea and 15-kg of MOP fertilizer should be done at this stage. Onion being a shallow rooted crop, **needs frequent light irrigation to maintain optimum soil moisture for proper growth and bulb development.**

Fruits:

Mango: The present weather is appropriate for blossoming of flowers. Spray NAA @ 200 ppm per tree, **with the help of ring method of irrigation** apply 25 kg FYM, 500 gram of urea and 1 kg of potash and phosphate each.

Cashew: Prevailing conditions are suitable for the incidence of **Tea mosquito bug**. Farmers are advised to go for spraying of Neem oil @ 5ml/L + Monocrotophos @ 1.6ml/L at new blush stage and Chlorpyrifos @ 2ml/L at flowering stage. If inflorescence have flourished stop spraying of any insecticides. Hormones like NAA @ 2 ppm can be sprayed. FYM @ 25 kg per tree can be applied.

Mushroom:

Initiate cultivation of paddy straw mushroom. As hot weather condition is prevailed in Cuttack district, Spray water if the mushrooms are at fruiting stage. Cover the Mushroom unit with water soaked Gunny bags to maintain the inside temperature.

Animal husbandry:

- As temperature is slightly increasing hence, it will impact physiological, metabolic behavior as well as egg production of poultry bird. Farmers should provide sufficient clean drinking water to birds and make proper shelter and feed management.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-376
Date- 21/02/2023

AGROMET ADVISORY BULLETIN FOR KANTAPADA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.8°C (-0.4 from normal) and 19.5°C (+1.13 from normal), respectively. The mean relative humidity was 53-79%.

Forecasted period: Today(8.30IST) to 26th February(8.30IST)2023

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 34-35°C and night temperature will be 21-23°C for upcoming 5 days. Maximum and minimum relative humidity will be 89-95% and 20-31% respectively. There will be clear sky for next 5 days.

Block	Kantapada (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	21/02/2023	22/02/2023	23/02/2023	24/02/2023	25/02/2023
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	34	34	35	34	35
T-Min(°C)	23	22	21	22	21
Cloud Cover(octa)	0	0	0	0	0
RH Max (%)	95	89	95	94	93
RH Min (%)	20	31	23	31	20
Wind Speed (Kmph)	12	17	13	11	10
Wind Direction (deg)	199	203	203	198	191

Agromet Advisory

General Advisory: Provide light and frequent irrigation to standing vegetables and crops during morning and evening hours to prevent damage from moisture and high temperature stress.

- Foliar spray of 2%DAP +1% KCL (MOP) during critical stages of flowering and grain formation stage.
- Use of biofertilizer like Azotobacter, Azospirillum and phospho bacteria @10 packets (500g)/ha along with 25 kg of soil or FYM.
- To avoid flower and fruit drop spray planofix @ 1ml/4.5lit of water.
- Ridge and furrow irrigation should be practiced to conserve soil moisture in the main field.

Pulses:

November to December sown crop is at pod formation to seed development (nine to twelve week) stage. Flowering and pod development stage is sensitive to moisture stress; carry out irrigation as per the status of soil moisture. In case of YMV infestation uproots and destroys the infected plant and spray imidacloprid 50ml/acre mixing with 200 lit of water.

Rabi-Groundnut:

Do not disturb the soil after flower initiation in groundnut. Do irrigation in the field based upon soil moisture. Increase in temperature may enhance red hairy caterpillar and tobacco caterpillar infestation in groundnut; to control, spray Profenophos 50% EC @ 2 ml/litre of water or Flubendiamide 480 SC @ 0.2 ml/litre of water.

Mustard:

Rained mustard sown during October is at maturity stage. Harvest the mustard crop as soon as the pods turn yellowish-brown. Present weather may favour the infestation of saw fly, leaf Webber. To control saw fly, spray 400ml Endosulphan/acre, to control leaf webber spray 200 ml Ithofenprox or Deltamethrin per acre in 200 litr water.

Vegetables:

Tomato: Transplanted seedlings are at 6-8 week stage in main field. Due to water stress, deficiency of boron or due to hormonal disbalance flower drop occurs. Control: Apply timely and required amount of irrigation. Spray Boron 20% 2 gram/lit during early flowering and 15 days later. Due to increase in temperature there may be chances of bacterial canker and Sun burn disease in tomato at fruiting stage. To control bacterial canker Farmers should spray Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water. They should use shade net to protect the crop from sun burn.

Potato: The main crop is ready for harvest within 75-85 days of planting depending upon the soil type and variety sown in our climatic condition of odisha. The crop is ready to harvest when majority of the leaves turn yellowish brown. Irrigation should be stopped 15 days before harvesting.

Onion: Do inter culture and hand weeding may be carried out at 45-50 DAT and application of 30-kg of urea and 15-kg of MOP fertilizer should be done at this stage. Onion being a shallow rooted crop, needs frequent light irrigation to maintain optimum soil moisture for proper growth and bulb development.

Fruits:

Mango: The present weather is appropriate for blossoming of flowers. Spray NAA @ 200 ppm per tree, with the help of ring method of irrigation apply 25 kg FYM, 500 gram of urea and 1 kg of potash and phosphate each.

Cashew: Prevailing conditions are suitable for the incidence of Tea mosquito bug. Farmers are advised to go for spraying of Neem oil @ 5ml/L + Monocrotophos @ 1.6ml/L at new blush stage and Chlorpyrifos @ 2ml/L at flowering stage. If inflorescence have flourished stop spraying of any insecticides. Hormones like NAA @ 2 ppm can be sprayed. FYM @25 kg per tree can be applied.

Mushroom:

Initiate cultivation of paddy straw mushroom. As hot weather condition is prevailed in Cuttack district, Spray water if the mushrooms are at fruiting stage. Cover the Mushroom unit with water soaked Gunny bags to maintain the inside temperature.

Animal husbandry:

- As temperature is slightly increasing hence, it will impact physiological, metabolic behavior as well as egg production of poultry bird. Farmers should provide sufficient clean drinking water to birds and make proper shelter and feed management.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-376
Date- 21/02/2023

AGROMET ADVISORY BULLETIN FOR MAHANGA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.8°C (-0.4 from normal) and 19.5°C (+1.13 from normal), respectively. The mean relative humidity was 53-79%.

Forecasted period: Today(8.30IST) to 26th February(8.30IST)2023

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be and it will be 34-35°C and night temperature will be 21-22°C for upcoming 5 days. Maximum and minimum relative humidity will be 92-96% and 14-27% respectively.

There will be clear sky for next 5 days.

Block	Mahanga (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	21/02/2023	22/02/2023	23/02/2023	24/02/2023	25/02/2023
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	34	35	35	34	34
T-Min(°C)	22	21	21	21	21
Cloud Cover(octal)	0	0	0	0	0
RH Max (%)	96	92	95	95	93
RH Min (%)	14	27	18	25	21
Wind Speed (Kmph)	12	17	11	9	9
Wind Direction (deg)	200	204	203	244	162

Agromet Advisory

General Advisory: Provide light and frequent irrigation to standing vegetables and crops during morning and evening hours to prevent damage from moisture and high temperature stress.

➤ Foliar spray of 2%DAP +1% KCL (MOP) during critical stages of flowering and grain formation stage.

➤ Use of biofertilizer like Azotobacter, Azospirillum and phospho bacteria @10 packets (500g)/ha along with 25 kg of soil or FYM.

➤ To avoid flower and fruit drop spray planofix @ 1ml/4.5lit of water.

➤ Ridge and furrow irrigation should be practiced to conserve soil moisture in the main field.

Pulses:

November to December sown crop is at pod formation to seed development (nine to twelve week) stage. Flowering and pod development stage is sensitive to moisture stress; carry out irrigation as per the status of soil moisture. In case of YMV infestation uproots and destroys the infected plant and spray imidacloprid 50ml/acre mixing with 200 lit of water.

Rabi-Groundnut:

Do not disturb the soil after flower initiation in groundnut. Do irrigation in the field based upon soil moisture. Increase in temperature may enhance red hairy caterpillar and tobacco caterpillar infestation in groundnut; to control, spray Profenophos 50% EC @ 2 ml/litre of water or Flubendiamide 480 SC @ 0.2 ml/litre of water.

Mustard:

Rained mustard sown during October is at maturity stage. Harvest the mustard crop as soon as the pods turn yellowish-brown. Present weather may favour the infestation of saw fly, leaf Webber. To control saw fly, spray 400ml Endosulphan/acre, to control leaf webber spray 200 ml Ithofenprox or Deltamethrin per acre in 200 litr water.

Vegetables:

Tomato: Transplanted seedlings are at 6-8 week stage in main field. Due to water stress, deficiency of boron or due to hormonal disbalance flower drop occurs. Control: Apply timely and required amount of irrigation. Spray Boron 20% 2 gram/lit during early flowering and 15 days later. Due to increase in temperature there may be chances of bacterial canker and Sun burn disease in tomato at fruiting stage. To control bacterial canker Farmers should spray Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water. They should use shade net to protect the crop from sun burn.

Potato: The main crop is ready for harvest within 75-85 days of planting depending upon the soil type and variety sown in our climatic condition of Odisha. The crop is ready to harvest when majority of the leaves turn yellowish brown. Irrigation should be stopped 15 days before harvesting.

Onion: Do inter culture and hand weeding may be carried out at 45-50 DAT and application of 30-kg of urea and 15-kg of MOP fertilizer should be done at this stage. Onion being a shallow rooted crop, needs frequent light irrigation to maintain optimum soil moisture for proper growth and bulb development.

Fruits:

Mango: The present weather is appropriate for blossoming of flowers. Spray NAA @ 200 ppm per tree, with the help of ring method of irrigation apply 25 kg FYM, 500 gram of urea and 1 kg of potash and phosphate each.

Cashew: Prevailing conditions are suitable for the incidence of Tea mosquito bug. Farmers are advised to go for spraying of Neem oil @ 5ml/L + Monocrotophos @ 1.6ml/L at new blush stage and Chlorpyrifos @ 2ml/L at flowering stage. If inflorescence have flourished stop spraying of any insecticides. Hormones like NAA @ 2 ppm can be sprayed. FYM @ 25 kg per tree can be applied.

Mushroom:

Initiate cultivation of paddy straw mushroom. As hot weather condition is prevailed in Cuttack district, Spray water if the mushrooms are at fruiting stage. Cover the Mushroom unit with water soaked Gunny bags to maintain the inside temperature.

Animal husbandry:

- As temperature is slightly increasing hence, it will impact physiological, metabolic behavior as well as egg production of poultry bird. Farmers should provide sufficient clean drinking water to birds and make proper shelter and feed management.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-376
Date- 21/02/2023

AGROMET ADVISORY BULLETIN FOR NARASINGHPUR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.8°C (-0.4 from normal) and 19.5°C (+1.13 from normal), respectively. The mean relative humidity was 53-79%.

Forecasted period: Today(8.30IST) to 26th February(8.30IST)2023

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 35-37°C and night temperature will be 18-19°C for upcoming 5 days. Maximum and minimum relative humidity will be 84-97% and 09-13% respectively. There will be clear sky for next 5 days.

Block	Narasinghpur (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	21/02/2023	22/02/2023	23/02/2023	24/02/2023	25/02/2023
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	36	37	37	37	37
T-Min(°C)	18	18	18	19	18
Cloud Cover(octa)	0	0	0	0	0
RH Max (%)	97	84	88	94	88
RH Min (%)	11	12	9	13	13
Wind Speed (Kmph)	6	8	5	5	6
Wind Direction (deg)	203	200	15	79	167

Agromet Advisory

General Advisory: Provide light and frequent irrigation to standing vegetables and crops during morning and evening hours to prevent damage from moisture and high temperature stress.

- Foliar spray of 2%DAP +1% KCL (MOP) during critical stages of flowering and grain formation stage.
- Use of biofertilizer like Azotobacter, Azospirillum and phospho bacteria @10 packets (500g)/ha along with 25 kg of soil or FYM.
- To avoid flower and fruit drop spray planofix @ 1ml/4.5lit of water.
- Ridge and furrow irrigation should be practiced to conserve soil moisture in the main field.

Pulses:

November to December sown crop is at pod formation to seed development (nine to twelve week) stage. Flowering and pod development stage is sensitive to moisture stress; carry out irrigation as per the status of soil moisture. In case of YMV infestation uproots and destroys the infected plant and spray imidacloprid 50ml/acre mixing with 200 lit of water.

Rabi-Groundnut:

Do not disturb the soil after flower initiation in groundnut. Do irrigation in the field based upon soil moisture. Increase in temperature may enhance red hairy caterpillar and tobacco caterpillar infestation in groundnut; to control, spray Profenophos 50% EC @ 2 ml/litre of water or Flubendiamide 480 SC @ 0.2 ml/litre of water.

Mustard:

Rainfed mustard sown during October is at maturity stage. Harvest the mustard crop as soon as the pods turn yellowish-brown. Present weather may favour the infestation of saw fly, leaf Webber. To control saw fly, spray 400ml Endosulphan/acre, to control leaf webber spray 200 ml Ithofenprox or Deltamethrin per acre in 200 lit water.

Vegetables:

Tomato: Transplanted seedlings are at 6-8 week stage in main field. Due to water stress, deficiency of boron or due to hormonal disbalance flower drop occurs. Control: Apply timely and required amount of irrigation. Spray Boron 20% 2 gram/lit during early flowering and 15 days later. Due to increase in temperature there may be chances of bacterial canker and Sun burn disease in tomato at fruiting stage. To control bacterial canker farmers should spray Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water. They should use shade net to protect the crop from sun burn.

Potato: The main crop is ready for harvest within 75-85 days of planting depending upon the soil type and variety sown in our climatic condition of Odisha. The crop is ready to harvest when majority of the leaves turn yellowish brown. Irrigation should be stopped 15 days before harvesting.

Onion: Do inter culture and hand weeding may be carried out at 45-50 DAT and application of 30-kg of urea and 15-kg of MOP fertilizer should be done at this stage. Onion being a shallow rooted crop, needs frequent light irrigation to maintain optimum soil moisture for proper growth and bulb development.

Fruits:

Mango: The present weather is appropriate for blossoming of flowers. Spray NAA @ 200 ppm per tree, with the help of ring method of irrigation apply 25 kg FYM, 500 gram of urea and 1 kg of potash and phosphate each.

Cashew: Prevailing conditions are suitable for the incidence of Tea mosquito bug. Farmers are advised to go for spraying of Neem oil @ 5ml/L + Monocrotophos @ 1.6ml/L at new blush stage and Chlorpyrifos @ 2ml/L at flowering stage. If inflorescence have flourished stop spraying of any insecticides. Hormones like NAA @ 2 ppm can be sprayed. FYM @ 25 kg per tree can be applied.

Mushroom:

Initiate cultivation of paddy straw mushroom. As hot weather condition is prevailed in Cuttack district, Spray water if the mushrooms are at fruiting stage. Cover the Mushroom unit with water soaked Gunny bags to maintain the inside temperature.

Animal husbandry:

- As temperature is slightly increasing hence, it will impact physiological, metabolic behavior as well as egg production of poultry bird. Farmers should provide sufficient clean drinking water to birds and make proper shelter and feed management.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-376
Date- 21/02/2023

AGROMET ADVISORY BULLETIN FOR NIALI

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.8°C (-0.4 from normal) and 19.5°C (+1.13 from normal), respectively. The mean relative humidity was 53-79%.

Forecasted period: Today(8.30IST) to 26th February(8.30IST)2023

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 33-35°C and night temperature will be 21-22°C for upcoming 5 days. Maximum and minimum relative humidity will be 86-94% and 24-38% respectively. There will be clear sky for next 5 days.

Block	Niali (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	21/02/2023	22/02/2023	23/02/2023	24/02/2023	25/02/2023
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	35	33	34	33	35
T-Min(°C)	22	22	22	22	21
Cloud Cover(octa)	0	0	0	0	0
RH Max (%)	94	86	94	93	92
RH Min (%)	30	36	30	38	24
Wind Speed(Kmph)	13	18	14	12	11
Wind Direction(deg)	203	203	203	193	191

Agromet Advisory

General Advisory: Provide light and frequent irrigation to standing vegetables and crops during morning and evening hours to prevent damage from moisture and high temperature stress.

- Foliar spray of 2%DAP +1% KCL (MOP) during critical stages of flowering and grain formation stage.
- Use of biofertilizer like Azotobacter, Azospirillum and phospho bacteria @10 packets (500g)/ha along with 25 kg of soil or FYM.
- To avoid flower and fruit drop spray planofix @ 1ml/4.5lit of water.
- Ridge and furrow irrigation should be practiced to conserve soil moisture in the main field.

Pulses:

November to December sown crop is at pod formation to seed development (nine to twelve week) stage. Flowering and pod development stage is sensitive to moisture stress; carry out irrigation as per the status of soil moisture. In case of YMV infestation uproots and destroys the infected plant and spray imidacloprid 50ml/acre mixing with 200 lit of water.

Rabi-Groundnut:

Do not disturb the soil after flower initiation in groundnut. Do irrigation in the field based upon soil moisture. Increase in temperature may enhance red hairy caterpillar and tobacco caterpillar infestation in groundnut; to control, spray Profenophos 50% EC @ 2 ml/litre of water or Flubendiamide 480 SC @ 0.2 ml/litre of water.

Mustard:

Rainfed mustard sown during October is at maturity stage. Harvest the mustard crop as soon as the pods turn yellowish-brown. Present weather may favour the infestation of saw fly, leaf Webber. To control saw fly, spray 400ml Endosulphan/acre, to control leaf webber spray 200 ml Ithofenprox or Deltamethrin per acre in 200 lit water.

Vegetables:

Tomato: Transplanted seedlings are at 6-8 week stage in main field. Due to water stress, deficiency of boron or due to hormonal disbalance flower drop occurs. Control: Apply timely and required amount of irrigation. Spray Boron 20% 2 gram/lit during early flowering and 15 days later. Due to increase in temperature there may be chances of bacterial canker and Sun burn disease in tomato at fruiting stage. To control bacterial canker Farmers should spray Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water. They should use shade net to protect the crop from sun burn.

Potato: The main crop is ready for harvest within 75-85 days of planting depending upon the soil type and variety sown in our climatic condition of Odisha. The crop is ready to harvest when majority of the leaves turn yellowish brown. Irrigation should be stopped 15 days before harvesting.

Onion: Do inter culture and hand weeding may be carried out at 45-50 DAT and application of 30-kg of urea and 15-kg of MOP fertilizer should be done at this stage. Onion being a shallow rooted crop, needs frequent light irrigation to maintain optimum soil moisture for proper growth and bulb development.

Fruits:

Mango: The present weather is appropriate for blossoming of flowers. Spray NAA @ 200 ppm per tree, with the help of ring method of irrigation apply 25 kg FYM, 500 gram of urea and 1 kg of potash and phosphate each.

Cashew: Prevailing conditions are suitable for the incidence of Tea mosquito bug. Farmers are advised to go for spraying of Neem oil @ 5ml/L + Monocrotophos @ 1.6ml/L at new blush stage and Chlorpyrifos @ 2ml/L at flowering stage. If inflorescence have flourished stop spraying of any insecticides. Hormones like NAA @ 2 ppm can be sprayed. FYM @ 25 kg per tree can be applied.

Mushroom:

Initiate cultivation of paddy straw mushroom. As hot weather condition is prevailed in Cuttack district, Spray water if the mushrooms are at fruiting stage. Cover the Mushroom unit with water soaked Gunny bags to maintain the inside temperature.

Animal husbandry:

- As temperature is slightly increasing hence, it will impact physiological, metabolic behavior as well as egg production of poultry bird. Farmers should provide sufficient clean drinking water to birds and make proper shelter and feed management.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-376
Date- 21/02/2023

AGROMET ADVISORY BULLETIN FOR NISCHINTAKOILI
District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.8°C (-0.4 from normal) and 19.5°C (+1.13 from normal), respectively. The mean relative humidity was 53-79%.

Forecasted period: Today(8.30IST) to 26th February(8.30IST)2023

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 34-35°C and night temperature will be 21-22°C for upcoming 5 days. Maximum and minimum relative humidity will be 92-96% and 15-29% respectively. There will be clear sky for next 5 days.

Block	Nischintakoili (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	21/02/2023	22/02/2023	23/02/2023	24/02/2023	25/02/2023
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	34	34	35	34	34
T-Min(°C)	22	21	21	21	21
Cloud Cover(octa)	0	0	0	0	0
RH Max (%)	96	92	95	95	93
RH Min (%)	15	29	20	27	21
Wind Speed (Kmph)	12	17	12	10	9
Wind Direction (deg)	200	203	205	240	162

Agromet Advisory

General Advisory: Provide light and frequent irrigation to standing vegetables and crops during morning and evening hours to prevent damage from moisture and high temperature stress.

- Foliar spray of 2%DAP +1% KCL (MOP) during critical stages of flowering and grain formation stage.
- Use of biofertilizer like Azotobacter, Azospirillum and phospho bacteria @10 packets (500g)/ha along with 25 kg of soil or FYM.
- To avoid flower and fruit drop spray planofix @ 1ml/4.5lit of water.
- Ridge and furrow irrigation should be practiced to conserve soil moisture in the main field.

Pulses:

November to December sown crop is at pod formation to seed development (nine to twelve week) stage. Flowering and pod development stage is sensitive to moisture stress; carry out irrigation as per the status of soil moisture. In case of YMV infestation uproots and destroys the infected plant and spray imidacloprid 50ml/acre mixing with 200 lit of water.

Rabi-Groundnut:

Do not disturb the soil after flower initiation in groundnut. Do irrigation in the field based upon soil moisture. Increase in temperature may enhance red hairy caterpillar and tobacco caterpillar infestation in groundnut; to control, spray Profenophos 50% EC @ 2 ml/litre of water or Flubendiamide 480 SC @ 0.2 ml/litre of water.

Mustard:

Rainfed mustard sown during October is at maturity stage. Harvest the mustard crop as soon as the pods turn yellowish-brown. Present weather may favour the infestation of saw fly, leaf webber. To control saw fly, spray 400ml Endosulphan/acre, to control leaf webber spray 200 ml Ithofenprox or Deltamethrin per acre in 200 lit water.

Vegetables:

Tomato: Transplanted seedlings are at 6-8 week stage in main field. Due to water stress, deficiency of boron or due to hormonal disbalance flower drop occurs. Control: Apply timely and required amount of irrigation. Spray Boron 20% 2 gram/lit during early flowering and 15 days later. Due to increase in temperature there may be chances of bacterial canker and Sun burn disease in tomato at fruiting stage. To control bacterial canker farmers should spray Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water. They should use shade net to protect the crop from sun burn.

Potato: The main crop is ready for harvest within 75-85 days of planting depending upon the soil type and variety sown in our climatic condition of Odisha. The crop is ready to harvest when majority of the leaves turn yellowish brown. Irrigation should be stopped 15 days before harvesting.

Onion: Do inter culture and hand weeding may be carried out at 45-50 DAT and application of 30-kg of urea and 15-kg of MOP fertilizer should be done at this stage. Onion being a shallow rooted crop, needs frequent light irrigation to maintain optimum soil moisture for proper growth and bulb development.

Fruits:

Mango: The present weather is appropriate for blossoming of flowers. Spray NAA @ 200 ppm per tree, with the help of ring method of irrigation apply 25 kg FYM, 500 gram of urea and 1 kg of potash and phosphate each.

Cashew: Prevailing conditions are suitable for the incidence of Tea mosquito bug. Farmers are advised to go for spraying of Neem oil @ 5ml/L + Monocrotophos @ 1.6ml/L at new blush stage and Chlorpyrifos @ 2ml/L at flowering stage. If inflorescence have flourished stop spraying of any insecticides. Hormones like NAA @ 2 ppm can be sprayed. FYM @25 kg per tree can be applied.

Mushroom:

Initiate cultivation of paddy straw mushroom. As hot weather condition is prevailed in Cuttack district, Spray water if the mushrooms are at fruiting stage. Cover the Mushroom unit with water soaked Gunny bags to maintain the inside temperature.

Animal husbandry:

- As temperature is slightly increasing hence, it will impact physiological, metabolic behavior as well as egg production of poultry bird. Farmers should provide sufficient clean drinking water to birds and make proper shelter and feed management.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-376
Date- 21/02/2023

AGROMET ADVISORY BULLETIN FOR SALEPUR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.8°C (-0.4 from normal) and 19.5°C (+1.13 from normal), respectively. The mean relative humidity was 53-79%.

Forecasted period: Today(8.30IST) to 26th February(8.30IST)2023

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 34-35°C and night temperature will be 21-22°C for upcoming 5 days. Maximum and minimum relative humidity will be 92-96% and 14-28% respectively. There will be clear sky for next 5 days.

Block	Salepur (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	21/02/2023	22/02/2023	23/02/2023	24/02/2023	25/02/2023
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	34	34	35	34	34
T-Min(°C)	22	21	21	21	21
Cloud Cover(octa)	0	0	0	0	0
RH Max (%)	96	92	95	95	93
RH Min (%)	14	28	18	25	20
Wind Speed (Kmph)	12	17	12	9	9
Wind Direction (deg)	200	204	205	169	162

Agromet Advisory

General Advisory: Provide light and frequent irrigation to standing vegetables and crops during morning and evening hours to prevent damage from moisture and high temperature stress.

- Foliar spray of 2%DAP +1% KCL (MOP) during critical stages of flowering and grain formation stage.
- Use of biofertilizer like Azotobacter, Azospirillum and phospho bacteria @10 packets (500g)/ha along with 25 kg of soil or FYM.
- To avoid flower and fruit drop spray planofix @ 1ml/4.5lit of water.
- Ridge and furrow irrigation should be practiced to conserve soil moisture in the main field.

Pulses:

November to December sown crop is at pod formation to seed development (nine to twelve week) stage. Flowering and pod development stage is sensitive to moisture stress; carry out irrigation as per the status of soil moisture. In case of YMV infestation uproots and destroys the infected plant and spray imidacloprid 50ml/acre mixing with 200 lit of water.

Rabi-Groundnut:

Do not disturb the soil after flower initiation in groundnut. Do irrigation in the field based upon soil moisture. Increase in temperature may enhance red hairy caterpillar and tobacco caterpillar infestation in groundnut; to control, spray Profenophos 50% EC @ 2 ml/litre of water or Flubendiamide 480 SC @ 0.2 ml/litre of water.

Mustard:

Rained mustard sown during October is at maturity stage. Harvest the mustard crop as soon as the pods turn yellowish-brown. Present weather may favour the infestation of saw fly, leaf Webber. To control saw fly, spray 400ml Endosulphan/acre, to control leaf webber spray 200 ml Ithofenprox or Deltamethrin per acre in 200 litr water.

Vegetables:

Tomato: Transplanted seedlings are at 6-8 week stage in main field. Due to water stress, deficiency of boron or due to hormonal disbalance flower drop occurs. Control: Apply timely and required amount of irrigation. Spray Boron 20% 2 gram/lit during early flowering and 15 days later. Due to increase in temperature there may be chances of bacterial canker and Sun burn disease in tomato at fruiting stage. To control bacterial canker Farmers should spray Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water. They should use shade net to protect the crop from sun burn.

Potato: The main crop is ready for harvest within 75-85 days of planting depending upon the soil type and variety sown in our climatic condition of Odisha. The crop is ready to harvest when majority of the leaves turn yellowish brown. Irrigation should be stopped 15 days before harvesting.

Onion: Do inter culture and hand weeding may be carried out at 45-50 DAT and application of 30-kg of urea and 15-kg of MOP fertilizer should be done at this stage. Onion being a shallow rooted crop, needs frequent light irrigation to maintain optimum soil moisture for proper growth and bulb development.

Fruits:

Mango: The present weather is appropriate for blossoming of flowers. Spray NAA @ 200 ppm per tree, with the help of ring method of irrigation apply 25 kg FYM, 500 gram of urea and 1 kg of potash and phosphate each.

Cashew: Prevailing conditions are suitable for the incidence of Tea mosquito bug. Farmers are advised to go for spraying of Neem oil @ 5ml/L + Monocrotophos @ 1.6ml/L at new blush stage and Chlorpyrifos @ 2ml/L at flowering stage. If inflorescence have flourished stop spraying of any insecticides. Hormones like NAA @ 2 ppm can be sprayed. FYM @25 kg per tree can be applied.

Mushroom:

Initiate cultivation of paddy straw mushroom. As hot weather condition is prevailed in Cuttack district, Spray water if the mushrooms are at fruiting stage. Cover the Mushroom unit with water soaked Gunny bags to maintain the inside temperature.

Animal husbandry:

- As temperature is slightly increasing hence, it will impact physiological, metabolic behavior as well as egg production of poultry bird. Farmers should provide sufficient clean drinking water to birds and make proper shelter and feed management.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-376
Date- 21/02/2023

AGROMET ADVISORY BULLETIN FOR TANGI-CHOUDWAR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.8°C (-0.4 from normal) and 19.5°C (+1.13 from normal), respectively. The mean relative humidity was 53-79%.

Forecasted period: Today(8.30IST) to 26th February(8.30IST)2023

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 33-35°C and night temperature will be 23-24°C for upcoming 5 days. Maximum and minimum relative humidity will be 83-92% and 41-47% respectively. There will be clear sky for next 5 days.

Block	Tangi-Choudwar (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	21/02/2023	22/02/2023	23/02/2023	24/02/2023	25/02/2023
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	34	34	34	35	35
T-Min(°C)	24	24	23	23	23
Cloud Cover(octa)	0	0	0	0	0
RH Max (%)	92	83	91	90	90
RH Min (%)	46	47	41	47	42
Wind Speed (Kmph)	16	20	16	13	11
Wind Direction (deg)	198	203	204	169	161

Agromet Advisory

General Advisory: Provide light and frequent irrigation to standing vegetables and crops during morning and evening hours to prevent damage from moisture and high temperature stress.

- Foliar spray of 2%DAP +1% KCL (MOP) during critical stages of flowering and grain formation stage.
- Use of biofertilizer like Azotobacter, Azospirillum and phospho bacteria @10 packets (500g)/ha along with 25 kg of soil or FYM.
- To avoid flower and fruit drop spray planofix @ 1ml/4.5lit of water.
- Ridge and furrow irrigation should be practiced to conserve soil moisture in the main field.

Pulses:

November to December sown crop is at pod formation to seed development (nine to twelve week) stage. Flowering and pod development stage is sensitive to moisture stress; carry out irrigation as per the status of soil moisture. In case of YMV infestation uproots and destroys the infected plant and spray imidacloprid 50ml/acre mixing with 200 lit of water.

Rabi-Groundnut:

Do not disturb the soil after flower initiation in groundnut. Do irrigation in the field based upon soil moisture. Increase in temperature may enhance red hairy caterpillar and tobacco caterpillar infestation in groundnut; to control, spray Profenophos 50% EC @ 2 ml/litre of water or Flubendiamide 480 SC @ 0.2 ml/litre of water.

Mustard:

Rained mustard sown during October is at maturity stage. Harvest the mustard crop as soon as the pods turn yellowish-brown. Present weather may favour the infestation of saw fly, leaf Webber. To control saw fly, spray 400ml Endosulphan/acre, to control leaf webber spray 200 ml Ithofenprox or Deltamethrin per acre in 200 litr water.

Vegetables:

Tomato: Transplanted seedlings are at 6-8 week stage in main field. Due to water stress, deficiency of boron or due to hormonal disbalance flower drop occurs. Control: Apply timely and required amount of irrigation. Spray Boron 20% 2 gram/lit during early flowering and 15 days later. Due to increase in temperature there may be chances of bacterial canker and Sun burn disease in tomato at fruiting stage. To control bacterial canker Darners should spray Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water. They should use shade net to protect the crop from sun burn.

Potato: The main crop is ready for harvest within 75-85 days of planting depending upon the soil type and variety sown in our climatic condition of Odisha. The crop is ready to harvest when majority of the leaves turn yellowish brown. Irrigation should be stopped 15 days before harvesting.

Onion: Do inter culture and hand weeding may be carried out at 45-50 DAT and application of 30-kg of urea and 15-kg of MOP fertilizer should be done at this stage. Onion being a shallow rooted crop, needs frequent light irrigation to maintain optimum soil moisture for proper growth and bulb development.

Fruits:

Mango: The present weather is appropriate for blossoming of flowers. Spray NAA @ 200 ppm per tree, with the help of ring method of irrigation apply 25 kg FYM, 500 gram of urea and 1 kg of potash and phosphate each.

Cashew: Prevailing conditions are suitable for the incidence of Tea mosquito bug. Farmers are advised to go for spraying of Neem oil @ 5ml/L + Monocrotophos @ 1.6ml/L at new blush stage and Chlorpyrifos @ 2ml/L at flowering stage. If inflorescence have flourished stop spraying of any insecticides. Hormones like NAA @ 2 ppm can be sprayed. FYM @25 kg per tree can be applied.

Mushroom:

Initiate cultivation of paddy straw mushroom. As hot weather condition is prevailed in Cuttack district, Spray water if the mushrooms are at fruiting stage. Cover the Mushroom unit with water soaked Gunny bags to maintain the inside temperature.

Animal husbandry:

- As temperature is slightly increasing hence, it will impact physiological, metabolic behavior as well as egg production of poultry bird. Farmers should provide sufficient clean drinking water to birds and make proper shelter and feed management.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-376
Date- 21/02/2023

AGROMET ADVISORY BULLETIN FOR TIGIRIA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.8°C (-0.4 from normal) and 19.5°C (+1.13 from normal), respectively. The mean relative humidity was 53-79%.

Forecasted period: Today(8.30IST) to 26th February(8.30IST)2023

Weather advisory: Dry weather is forecasted for upcoming 5 days in the block. The day temperature will be 33-36°C and night temperature will be 20-21°C for upcoming 5 days. Maximum and minimum relative humidity will be 89-95% and 10-18% respectively. There will be clear sky for next 5 days.

Block	Tigiria (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	21/02/2023	22/02/2023	23/02/2023	24/02/2023	25/02/2023
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	33	34	34	35	36
T-Min(°C)	21	20	20	20	20
Cloud Cover(octa)	0	0	0	0	0
RH Max (%)	95	89	93	93	92
RH Min (%)	10	18	11	15	13
Wind Speed (Kmph)	9	13	7	7	9
Wind Direction (deg)	199	203	23	154	169

Agro met Advisory

General Advisory: Provide light and frequent irrigation to standing vegetables and crops during morning and evening hours to prevent damage from moisture and high temperature stress.

- Foliar spray of 2%DAP +1% KCL (MOP) during critical stages of flowering and grain formation stage.
- Use of biofertilizer like Azotobacter, Azospirillum and phospho bacteria @10 packets (500g)/ha along with 25 kg of soil or FYM.
- To avoid flower and fruit drop spray planofix @ 1ml/4.5lit of water.
- Ridge and furrow irrigation should be practiced to conserve soil moisture in the main field.

Pulses:

November to December sown crop is at pod formation to seed development (nine to twelve week) stage. Flowering and pod development stage is sensitive to moisture stress; carry out irrigation as per the status of soil moisture. In case of YMV infestation uproots and destroys the infected plant and spray imidacloprid 50ml/acre mixing with 200 lit of water.

Rabi-Groundnut:

Do not disturb the soil after flower initiation in groundnut. Do irrigation in the field based upon soil moisture. Increase in temperature may enhance red hairy caterpillar and tobacco caterpillar infestation in groundnut; to control, spray Profenophos 50% EC @ 2 ml/litre of water or Flubendiamide 480 SC @ 0.2 ml/litre of water.

Mustard:

Rainfed mustard sown during October is at maturity stage. Harvest the mustard crop as soon as the pods turn yellowish-brown. Present weather may favour the infestation of saw fly, leaf Webber. To control saw fly, spray 400ml Endosulphan/acre, to control leaf webber spray 200 ml Ithofenprox or Deltamethrin per acre in 200 lit water.

Vegetables:

Tomato: Transplanted seedlings are at 6-8 week stage in main field. Due to water stress, deficiency of boron or due to hormonal disbalance flower drop occurs. Control: Apply timely and required amount of irrigation. Spray Boron 20% 2 gram/lit during early flowering and 15 days later. Due to increase in temperature there may be chances of bacterial canker and Sun burn disease in tomato at fruiting stage. To control bacterial canker farmers should spray Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water. They should use shade net to protect the crop from sun burn.

Potato: The main crop is ready for harvest within 75-85 days of planting depending upon the soil type and variety sown in our climatic condition of Odisha. The crop is ready to harvest when majority of the leaves turn yellowish brown. Irrigation should be stopped 15 days before harvesting.

Onion: Do inter culture and hand weeding may be carried out at 45-50 DAT and application of 30-kg of urea and 15-kg of MOP fertilizer should be done at this stage. Onion being a shallow rooted crop, needs frequent light irrigation to maintain optimum soil moisture for proper growth and bulb development.

Fruits:

Mango: The present weather is appropriate for blossoming of flowers. Spray NAA @ 200 ppm per tree, with the help of ring method of irrigation apply 25 kg FYM, 500 gram of urea and 1 kg of potash and phosphate each.

Cashew: Prevailing conditions are suitable for the incidence of Tea mosquito bug. Farmers are advised to go for spraying of Neem oil @ 5ml/L + Monocrotophos @ 1.6ml/L at new blush stage and Chlorpyrifos @ 2ml/L at flowering stage. If inflorescence have flourished stop spraying of any insecticides. Hormones like NAA @ 2 ppm can be sprayed. FYM @25 kg per tree can be applied.

Mushroom:

Initiate cultivation of paddy straw mushroom. As hot weather condition is prevailed in Cuttack district, Spray water if the mushrooms are at fruiting stage. Cover the Mushroom unit with water soaked Gunny bags to maintain the inside temperature.

Animal husbandry:

- As temperature is slightly increasing hence, it will impact physiological, metabolic behavior as well as egg production of poultry bird. Farmers should provide sufficient clean drinking water to birds and make proper shelter and feed management.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-376
Date- 21/02/2023

				
MEGHDOOT APP	DAMINI APP	FACEBOOK (Damu Cuttack)	BLOCK SPECIFIC WHAT'S APP GROUPS	@DAMUCUTTACK

- Kindly install and use **“Meghdoot mobile App”** for Cuttack district weather forecast, warnings and weather based agromet advisory services.
<https://play.google.com/store/apps/details?id=com.aas.meghdoot>
- Kindly install and use **“Damini mobile App”** for location specific for lightening forecast and warnings.
<https://play.google.com/store/apps/details?id=com.lightening.live.damini>
- Follow the **block specific whats app groups** and join in your block wise groups for getting weather based agromet advisory services twice in a week (Tuesday and Friday) and other weather warning based farm management advisories.
- Follow the **Official Website links** to get the weather based agromet advisory service bulletins **(English & Odia)** twice in a week (Tuesday and Friday).

1	ICAR-NRRI, Cuttack, Odisha	https://icar-nrri.in/agromate/	
2	Agrimet division, IMD, Pune	https://imdagrmet.gov.in/imdproject/AGIndex.php	

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427