





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-413
Date- 30/06/2023

STANDARD PRECIPITATION INDEX AND NDVI MAP

Standard Precipitation Index (SPI)	Normalized difference vegetation Index (NDVI)
<p>Standardised Precipitation Index for the period 25th May to 21st June 2023</p>  <p>As per SPI value (cumulative 4weeks -25May to 21 June 2023), Mildly dry condition was there in Cuttack district.</p>	<p>NOAA/VIIRS/BLENDED NDVI Composite ending on Week no 25 (18.06.2023 to 24.06.2023) over Agricultural region</p> <p style="text-align: center;">Odisha</p>  <p>Agriculture vigour is moderate over the state.</p> <p>According to NDVI map (18.06.2023 to 24.06.2023) the agriculture vigor was moderate over isolated places of Narsinghpur, Banki-dampada, Badambaa, Athgad blocks of Cuttack district and very light over other places of other blocks.</p>

LEGEND (Medium range block level rainfall forecast)		
	No Rainfall	00 mm
	Very light rainfall	01-05 mm
	Light rainfall	06-10 mm
	Light to moderate rainfall	11-30 mm
	Moderate to heavy rainfall	31-65 mm
	Heavy to very heavy rainfall	66-115 mm
	Extremely heavy rainfall	>116 mm

LEGEND (Past week temperature)		
Color code	Temperature	Departure
	Below Normal	< 2°C from normal
	Normal	-2°C to +2°C from normal
	Above Normal	>2°C from normal

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427
















GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-413
Date- 30/06/2023

Weather (Past and Forecasted) based Disease and pest in Major Crops and vegetables

Sl No	Crop	Stage and condition of the crop	Anticipated Disease and pest	
1	Sugarcane	Vegetative		Red rot
				Wilt
2	Jute	Sowing		Weed
3	Brinjal	Fruiting		Phomopsis Blight
				Fruit and shoot borer
4	Tomato	Fruting		Bacterial wilt
				Seedling rot
5	Cowpea			Rust
6	Cucurbits	Flowering to Fruiting stage		Downy mildew
7	Okra	Fruiting		YMV
8	Banana			Banana leaf spot/sigatoka
				Panama wilt
9	Papaya	Fruting		phytophthora crown rot or foot rot

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA INDIA METEOROLOGICAL DEPARTMENT DISTRICT AGROMET UNIT KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-413

Date- 30/06/2023

District Past week average weather condition and Block past week average Rainfall condition

The block received average 22mm and district received average 10mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.42°C (-1.62°C from normal) and 26°C (+1.8°C from normal), respectively. Mean relative humidity of the district was around 87-90%.

Forecasted period: Today (8.30IST) to 05th July (8.30IST) 2023

Weather advisory: Very light to light rainfall is forecasted for today to Sunday and Moderate rainfall/ thundershower is forecasted for other 2 days in the block. The day temperature will be 34-36°C and night temperature will be 27-29°C for upcoming 5 days. Maximum and minimum relative humidity will be 61-81% and 37-43% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Athagad (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	30/06/2023	01/07/2023	02/07/2023	03/07/2023	04/07/2023
Rainfall (mm)	5	4	10	20	30
T-Max (°C)	34	35	35	36	36
T-Min (°C)	28	29	29	28	27
Cloud Cover (octa)	6	5	7	6	5
RH Max (%)	72	67	61	67	81
RH Min (%)	43	38	37	37	37
Wind Speed (kmph)	14	13	13	15	16
Wind Direction (deg)	205	210	205	198	167

Agromet Advisory

General advisory: Farmer should wait for sufficient amount of rainfall (75-100 mm in a week) for sowing of Karif Rice.

Kharif rice:

Sowing should be started soon after receive of good amount of early monsoon showers (75-100 mm in a week) where direct seeding is practiced. Incorporate FYM or compost @ 2000 Kg/acre. Apply 27 kg DAP and 20 kg MoP per acre during last ploughing. Before sowing seed should be treated with Trichoderma viride @ 10 g or Carbendazim 50% WP @ 1.5 g/kg seed.

Transplanting rice:

After receive of sufficient rainfall, farmers are advised for sowing of dhanicha seeds @ 12 kg/acre prior to the transplanting of paddy so that the dhanicha crop will be 3-4 weeks old before 10-15 days of transplanting of rice. Apply all phosphatic fertilizers of Kharif rice to this green manuring crop in advance. Farmers who are sowing paddy nurseries are advised to treat the seeds with carbendizem @ 3 g/kg seed followed by shade drying for 24 hours before sowing in case of dry nursery and in case of wet nursery seeds should be treated with carbendizem @ 1 g/l of water by soaking for 24 hrs and another 24 hrs for sprouting and the seeds should be sown in wet nurseries. Raise the nursery for transplanted kharif rice. 400 m² area is required for one acre land. Prepare a 3 feet width and 6 inch raised bed with suitable length keeping 1 feet between two beds for irrigation and drainage. 40 baskets of FYM, 12 kg SSP and 2 kg MoP is required for 400 m² nursery area.

Maize: It is time for sowing of maize for green cob purpose under irrigated condition. Grow high yielding varieties like Navjot, Shakti, Decan-107, Decan-109, Use 6 kg seeds/acre. Apply one third of N and all P & K of the recommended dose of 50:24:24 kg. NPK/acre. Sow the seeds in lines with spacing of 45-60X20 cm at 5 cm soil depth.

VEGETABLES AND FRUITS:

- Fruit and shoot borer** in **Brinjal**- Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@300g/acre at 15 days interval.
- Due to Past rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- After cessation of rain, **downy mildew disease** will appear in **cucurbits**. To control it, go for alternate spray of Metalaxyl+Mancozeb@2g/litre and Azoxystrobin+Tebuconazole @0.6g/litre of water at 10 days interval.
- Rust** disease is likely to appear in **Cowpea**. Spray Chlorothalonil@2g/litre of water.
- Wilting** may observe in **tomato and brinjal** field, to control it drenching with Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water.
- Current weather favours spread of **YMV** in **Ladies finger**. In case of appearance of diseases, rogue out affected plants and spray Thiometoxam 25WG @ 60g/acre.

Fruit:

Banana: For leaf spot/sigatoka as a precautionary measure spray carbendizim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalonil @ 2.0 g / litre of water. Drenching soil with Metiram 75% WG @ 1.5g/4 lit or carbendazim 12%+ mancozeb 63% @ 2g/lit to prevent Panama wilt.

Mango: Harvest the mature mango fruits during morning or evening hours. Keep the harvested fruits in shade to prevent from spongy tissue disease and heat.

Papaya: After rain due to water logging there may be increase in the infestation of **phytophthora crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

Animal Husbandry: Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination. Keep the animal shed clean and dry.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA INDIA METEOROLOGICAL DEPARTMENT DISTRICT AGROMET UNIT KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-413

Date- 30/06/2023

AGROMET ADVISORY BULLETIN FOR BADAMBA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 07mm and district received average 10mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.42°C (-1.62°C from normal) and 26°C (+1.8°C from normal), respectively. Mean relative humidity of the district was around 87-90%.

Forecasted period: Today (8.30IST) to 05th July (8.30IST) 2023

Weather advisory: Very light to light rainfall is forecasted for today to Sunday and Moderate rainfall/ thundershower is forecasted for other 2 days in the block. The day temperature will be 34-36°C and night temperature will be 26-29°C for upcoming 5 days. Maximum and minimum relative humidity will be 62-81% and 36-42% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Badamba (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	30/06/2023	01/07/2023	02/07/2023	03/07/2023	04/07/2023
Rainfall (mm)	5	4	10	20	30
T-Max (°C)	34	35	36	36	36
T-Min (°C)	27	28	29	28	26
Cloud Cover (octa)	6	4	7	6	5
RH Max (%)	71	66	62	65	81
RH Min (%)	42	38	37	36	38
Wind Speed (Kmph)	13	12	13	14	15
Wind Direction (deg)	207	210	205	203	167

Agromet Advisory

General advisory: Farmer should wait for sufficient amount of rainfall (75-100 mm in a week) for sowing of Kharif Rice.

Kharif rice:

Sowing should be started soon after receive of good amount of early monsoon showers (75-100 mm in a week) where direct seeding is practiced. Incorporate FYM or compost @ 2000 Kg/acre. Apply 27 kg DAP and 20 kg MoP per acre during last ploughing. Before sowing seed should be treated with Trichoderma viride @ 10 g or Carbendazim 50% WP @ 1.5 g/kg seed.

Transplanting rice:

After receive of sufficient rainfall, farmers are advised for sowing of dhanicha seeds @ 12 kg/acre prior to the transplanting of paddy so that the dhanicha crop will be 3-4 weeks old before 10-15 days of transplanting of rice. Apply all phosphatic fertilizers of Kharif rice to this green manuring crop in advance. Farmers who are sowing paddy nurseries are advised to treat the seeds with carbendazim @ 3 g/kg seed followed by shade drying for 24 hours before sowing in case of dry nursery and in case of wet nursery seeds should be treated with carbendazim @ 1 g/l of water by soaking for 24 hrs and another 24 hrs for sprouting and the seeds should be sown in wet nurseries. Raise the nursery for transplanted kharif rice. 400 m² area is required for one acre land. Prepare a 3 feet width and 6 inch raised bed with suitable length keeping 1 feet between two beds for irrigation and drainage. 40 baskets of FYM, 12 kg SSP and 2 kg MoP is required for 400 m² nursery area.

Maize: It is time for sowing of maize for green cob purpose under irrigated condition. Grow high yielding varieties like Navjot, Shakti, Decan-107, Decan-109. Use 6 kg seeds/acre. Apply one third of N and all P & K of the recommended dose of 50:24:24 kg. NPK/acre. Sow the seeds in lines with spacing of 45-60X20 cm at 5 cm soil depth.

VEGETABLES AND FRUITS:

- Fruit and shoot borer** in **Brinjal**- Spray Neem based pesticide 1500ppm @ 600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC @ 70ml/acre and Thiodicarb @ 300g/acre at 15 days interval.
- Due to Past rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- After cessation of rain, **downy mildew disease** will appear in **cucurbits**. To control it, go for alternate spray of Metalaxyl+Mancozeb @ 2g/litre and Azoxystrobin+Tebuconazole @ 0.6g/litre of water at 10 days interval.
- Rust** disease is likely to appear in **Cowpea**. Spray Chlorothalonil @ 2g/litre of water.
- Wilting** may observe in **tomato and brinjal** field, to control it drenching with Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water.
- Current weather favours spread of **YMV** in **Ladies finger**. In case of appearance of diseases, rogue out affected plants and spray Thiometoxam 25WG @ 60g/acre.

Fruit:

Banana: For leaf spot/sigatoka as a precautionary measure spray carbendazim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalonil @ 2.0 g / litre of water. Drenching soil with Metiram 75% WG @ 1.5g/4 lit or carbendazim 12%+ mancozeb 63% @ 2g/lit to prevent Panama wilt.

Mango: Harvest the mature mango fruits during morning or evening hours. Keep the harvested fruits in shade to prevent from spongy tissue disease and heat.

Papaya: After rain due to water logging there may be increase in the infestation of **phytophthora crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

Animal Husbandry: Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination. Keep the animal shed clean and dry.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-413
Date- 30/06/2023

AGROMET ADVISORY BULLETIN FOR BANKI

District Past week average weather condition

The district received average 10mm rainfall during the last week. The mean daily maximum and minimum temperature was **31.42°C(-1.62°C from normal)** and **26°C(+1.8°C from normal)**, respectively. Mean relative humidity of the district was around 87-90%.

Forecasted period: Today(8.30IST) to 05th July (8.30IST)2023

Weather advisory: *Very light to light rainfall is forecasted for today to Sunday and Moderate rainfall/ thundershower is forecasted for other 2 days in the block.* The day temperature will be **34-36°C** and night temperature will be **26-29°C** for upcoming 5 days. Maximum and minimum relative humidity will be 61-81% and 35-41% respectively. **There will be partly cloudy to overcast sky for next 5 days.**

Block	Banki (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	30/06/2023	01/07/2023	02/07/2023	03/07/2023	04/07/2023
Rainfall(mm)	5	4	10	20	30
T-Max(°C)	34	35	35	36	36
T-Min(°C)	27	28	29	28	26
Cloud Cover(octa)	6	4	7	6	5
RH Max (%)	71	66	61	65	81
RH Min (%)	41	37	36	35	37
Wind Speed (Kmph)	14	14	14	16	17
Wind Direction (deg)	205	210	205	203	180

Agromet Advisory

General advisory: **Farmer should wait for sufficient amount of rainfall (75-100 mm in a week) for sowing of Karif Rice.**

Kharif rice:

Sowing should be started soon after receive of good amount of early monsoon showers (**75-100 mm in a week**) where direct seeding is practiced. Incorporate FYM or compost @ 2000 Kg/acre. Apply 27 kg DAP and 20 kg MoP per acre during last ploughing. Before sowing seed should be treated with Trichoderma viride @ 10 g or Carbendazim 50% WP @ 1.5 g/kg seed.

Transplanting rice:

After receive of sufficient rainfall, farmers are advised for sowing of dhanicha seeds @ 12 kg/acre prior to the transplanting of paddy so that the dhanicha crop will be 3-4 weeks old before 10-15 days of transplanting of rice. Apply all phosphatic fertilizers of Kharif rice to this green manuring crop in advance. Farmers who are sowing paddy nurseries are advised to treat the seeds with carbendizim @3 g/kg seed followed by shade drying for 24 hours before sowing in case of dry nursery and in case of wet nursery seeds should be treated with carbendizim @ 1 g/l of water bysoaking for 24 hrs and another 24 hrs for sprouting and the seeds should be sown in wet nurseries. Raise the nursery for transplanted kharif rice. 400 m2 area is required for one acre land. Prepare a 3 feet width and 6 inch raised bed with suitable length keeping 1 feet between two beds for irrigation and drainage. 40 baskets of FYM, 12 kg SSP and 2 kg MoP is required for 400 m2 nursery area.

Maize: It is time for sowing of maize for green cob purpose under irrigated condition Grow high yielding varieties like Navjot, Shakti, Decan-107, Decan-109, Use 6 kg seeds/acre. Apply one third of N and all P & K of the recommended dose of 50:24:24 kg. NPK/acre. Sow the seeds in lines with spacing of 45-60X20 cm at 5 cm soil depth.

VEGETABLES AND FRUITS:

- Fruit and shoot borer** in **Brinjal**- Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@300g/acre at 15 days interval.
- Due to Past rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- After cessation of rain, **downy mildew disease** will appear in **cucurbits**. To control it, go for alternate spray of Metalaxyl+Mancozeb@2g/litre and Azoxystrobin+Tebuconazole @0.6g/litre of water at 10 days interval.
- Rust** disease is likely to appear in **Cowpea**. Spray Chlorothalonil@2g/litre of water.
- Wilting** may observe in **tomato and brinjal** field, to control it drenching with Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water.
- Current weather favours spread of **YMV** in **Ladies finger**. In case of appearance of diseases, rogue out affected plants and spray Thiomethoxam 25WG @ 60g/acre.

Fruit:

Banana: For leaf spot/sigatoka as a precautionary measure spray carbendizim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalonil @ 2.0 g/ litre of water. Drenching soil with Metiram 75% WG @1.5g/4 lit or carbendazim 12%+ mancozeb 63% @ 2g/lit to prevent **Panama wilt**.

Mango: Harvest the mature mango fruits during morning or evening hours. Keep the harvested fruits in shade to prevent from spongy tissue disease and heat.

Papaya: After rain due to water logging there may be increase in the infestation of **phytophthora crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

Animal Husbandry: Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination. Keep the animal shed clean and dry.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA INDIA METEOROLOGICAL DEPARTMENT DISTRICT AGROMET UNIT KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-413

Date- 30/06/2023

AGROMET ADVISORY BULLETIN FOR BANKI-DAMPARA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 07mm and district received average 10mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.42°C (-1.62°C from normal) and 26°C (+1.8°C from normal), respectively. Mean relative humidity of the district was around 87-90%.

Forecasted period: Today (8.30IST) to 05th July (8.30IST) 2023

Weather Advisory: Very light to light rainfall is forecasted for today to Sunday and Moderate rainfall/ thundershower is forecasted for other 2 days in the block. The day temperature will be 34-36°C and night temperature will be 26-29°C for upcoming 5 days. Maximum and minimum relative humidity will be 61-81% and 35-42% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Banki-Dampada (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	30/06/2023	01/07/2023	02/07/2023	03/07/2023	04/07/2023
Rainfall(mm)	5	4	10	20	30
T-Max(°C)	34	35	35	36	36
T-Min(°C)	27	28	29	28	26
Cloud Cover(octa)	6	4	7	6	5
RH Max (%)	71	66	61	65	81
RH Min (%)	42	37	36	35	37
Wind Speed (Kmph)	14	13	14	16	17
Wind Direction (deg)	205	210	205	203	180

Agromet Advisory

General advisory: Farmer should wait for sufficient amount of rainfall (75-100 mm in a week) for sowing of Kharif Rice.

Kharif rice:

Sowing should be started soon after receive of good amount of early monsoon showers (75-100 mm in a week) where direct seeding is practiced. Incorporate FYM or compost @ 2000 Kg/acre. Apply 27 kg DAP and 20 kg MoP per acre during last ploughing. Before sowing seed should be treated with Trichoderma viride @ 10 g or Carbendazim 50% WP @ 1.5 g/kg seed.

Transplanting rice:

After receive of sufficient rainfall, farmers are advised for sowing of dhanicha seeds @ 12 kg/acre prior to the transplanting of paddy so that the dhanicha crop will be 3-4 weeks old before 10-15 days of transplanting of rice. Apply all phosphatic fertilizers of Kharif rice to this green manuring crop in advance. Farmers who are sowing paddy nurseries are advised to treat the seeds with carbendazim @ 3 g/kg seed followed by shade drying for 24 hours before sowing in case of dry nursery and in case of wet nursery seeds should be treated with carbendazim @ 1 g/l of water by soaking for 24 hrs and another 24 hrs for sprouting and the seeds should be sown in wet nurseries. Raise the nursery for transplanted kharif rice. 400 m² area is required for one acre land. Prepare a 3 feet width and 6 inch raised bed with suitable length keeping 1 feet between two beds for irrigation and drainage. 40 baskets of FYM, 12 kg SSP and 2 kg MoP is required for 400 m² nursery area.

Maize: It is time for sowing of maize for green cob purpose under irrigated condition. Grow high yielding varieties like Navjot, Shakti, Decan-107, Decan-109. Use 6 kg seeds/acre. Apply one third of N and all P & K of the recommended dose of 50:24:24 kg. NPK/acre. Sow the seeds in lines with spacing of 45-60X20 cm at 5 cm soil depth.

VEGETABLES AND FRUITS:

- Fruit and shoot borer** in Brinjal- Spray Neem based pesticide 1500ppm @ 600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC @ 70ml/acre and Thiodicarb @ 300g/acre at 15 days interval.
- Due to Past rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- After cessation of rain, **downy mildew disease** will appear in **cucurbits**. To control it, go for alternate spray of Metalaxyl+Mancozeb @ 2g/litre and Azoxystrobin+Tebuconazole @ 0.6g/litre of water at 10 days interval.
- Rust** disease is likely to appear in **Cowpea**. Spray Chlorothalonil @ 2g/litre of water.
- Wilting** may observe in **tomato and brinjal** field, to control it drenching with Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water.
- Current weather favours spread of **YMV** in **Ladies finger**. In case of appearance of diseases, rogue out affected plants and spray Thiometoxam 25WG @ 60g/acre.

Fruit:

Banana: For leaf spot/sigatoka as a precautionary measure spray carbendazim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalonil @ 2.0 g / litre of water. Drenching soil with Metiram 75% WG @ 1.5g/4 lit or carbendazim 12%+ mancozeb 63% @ 2g/lit to prevent

Panama wilt.

Mango: Harvest the mature mango fruits during morning or evening hours. Keep the harvested fruits in shade to prevent from spongy tissue disease and heat.

Papaya: After rain due to water logging there may be increase in the infestation of **phytophthora crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

Animal Husbandry: Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination. Keep the animal shed clean and dry.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-413
Date- 30/06/2023

AGROMET ADVISORY BULLETIN FOR BARANGA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 12mm and district received average 10mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.42°C (-1.62°C from normal) and 26°C (+1.8°C from normal), respectively. Mean relative humidity of the district was around 87-90%.

Forecasted period: Today (8.30IST) to 05th July (8.30IST) 2023

Weather advisory: Very light to light rainfall is forecasted for today to Sunday and Moderate rainfall/ thundershower is forecasted for other 2 days in the block. The day temperature will be 34-36°C and night temperature will be 27-30°C for upcoming 5 days. Maximum and minimum relative humidity will be 61-81% and 35-41% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Baranga (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	30/06/2023	01/07/2023	02/07/2023	03/07/2023	04/07/2023
Rainfall (mm)	5	4	10	20	30
T-Max (°C)	34	35	35	36	36
T-Min (°C)	28	29	30	28	27
Cloud Cover	5	4	7	6	5
RH Max (%)	70	65	61	68	81
RH Min (%)	41	36	37	35	36
Wind Speed (Kmph)	16	14	14	17	18
Wind Direction (deg)	205	208	203	200	180

Agromet Advisory

General advisory: Farmer should wait for sufficient amount of rainfall (75-100 mm in a week) for sowing of Kharif Rice.

Kharif rice:

Sowing should be started soon after receive of good amount of early monsoon showers (75-100 mm in a week) where direct seeding is practiced. Incorporate FYM or compost @ 2000 Kg/acre. Apply 27 kg DAP and 20 kg MoP per acre during last ploughing. Before sowing seed should be treated with Trichoderma viride @ 10 g or Carbendazim 50% WP @ 1.5 g/kg seed.

Transplanting rice:

After receive of sufficient rainfall, farmers are advised for sowing of dhanicha seeds @ 12 kg/acre prior to the transplanting of paddy so that the dhanicha crop will be 3-4 weeks old before 10-15 days of transplanting of rice. Apply all phosphatic fertilizers of Kharif rice to this green manuring crop in advance. Farmers who are sowing paddy nurseries are advised to treat the seeds with carbendazim @ 3 g/kg seed followed by shade drying for 24 hours before sowing in case of dry nursery and in case of wet nursery seeds should be treated with carbendazim @ 1 g/l of water by soaking for 24 hrs and another 24 hrs for sprouting and the seeds should be sown in wet nurseries. Raise the nursery for transplanted kharif rice. 400 m² area is required for one acre land. Prepare a 3 feet width and 6 inch raised bed with suitable length keeping 1 feet between two beds for irrigation and drainage. 40 baskets of FYM, 12 kg SSP and 2 kg MoP is required for 400 m² nursery area.

Maize: It is time for sowing of maize for green cob purpose under irrigated condition. Grow high yielding varieties like Navjot, Shakti, Decan-107, Decan-109. Use 6 kg seeds/acre. Apply one third of N and all P & K of the recommended dose of 50:24:24 kg. NPK/acre. Sow the seeds in lines with spacing of 45-60X20 cm at 5 cm soil depth.

VEGETABLES AND FRUITS:

- Fruit and shoot borer** in **Brinjal**- Spray Neem based pesticide 1500ppm @ 600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC @ 70ml/acre and Thiodicarb @ 300g/acre at 15 days interval.
- Due to Past rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- After cessation of rain, **downy mildew disease** will appear in **cucurbits**. To control it, go for alternate spray of Metalaxyl+Mancozeb @ 2g/litre and Azoxystrobin+Tebuconazole @ 0.6g/litre of water at 10 days interval.
- Rust** disease is likely to appear in **Cowpea**. Spray Chlorothalonil @ 2g/litre of water.
- Wilting** may observe in **tomato and brinjal** field, to control it drenching with Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water.
- Current weather favours spread of **YMV** in **Ladies finger**. In case of appearance of diseases, rogue out affected plants and spray Thiometoxam 25WG @ 60g/acre.

Fruit:

Banana: For leaf spot/sigatoka as a precautionary measure spray carbendazim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalonil @ 2.0 g / litre of water. Drenching soil with Metiram 75% WG @ 1.5g/4 lit or carbendazim 12%+ mancozeb 63% @ 2g/lit to prevent Panama wilt.

Mango: Harvest the mature mango fruits during morning or evening hours. Keep the harvested fruits in shade to prevent from spongy tissue disease and heat.

Papaya: After rain due to water logging there may be increase in the infestation of **phytophthora crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

Animal Husbandry: Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination. Keep the animal shed clean and dry.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)



Bulletin No.-413
Date- 30/06/2023

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

AGROMET ADVISORY BULLETIN FOR CUTTACK SADAR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 06mm and district received average 10mm rainfall during the last week. The mean daily maximum and minimum temperature was **31.42°C(-1.62°C from normal)** and **26°C(+1.8°C from normal)**, respectively. Mean relative humidity of the district was around 87-90%.

Forecasted period: Today(8.30IST) to 05th July (8.30IST)2023

Weather advisory: *Very light to light rainfall is forecasted for today to Sunday and Moderate rainfall/ thundershower is forecasted for other 2 days in the block.* The day temperature will be **34-36°C** and night temperature will be **27-30°C** for upcoming 5 days. Maximum and minimum relative humidity will be 61-81% and 36-42% respectively. **There will be partly cloudy to overcast sky for next 5 days.**

Block	Cuttack sadar (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	30/06/2023	01/07/2023	02/07/2023	03/07/2023	04/07/2023
Rainfall(mm)	5	4	10	20	30
T-Max(°C)	34	35	35	36	36
T-Min(°C)	28	29	30	28	27
Cloud Cover(octa)	6	4	7	6	5
RH Max (%)	71	66	61	69	81
RH Min (%)	42	37	36	36	37
Wind Speed (Kmph)	15	14	14	16	17
Wind Direction (deg)	205	208	203	198	180

Agromet Advisory

General advisory: **Farmer should wait for sufficient amount of rainfall (75-100 mm in a week) for sowing of Karif Rice.**

Kharif rice:

Sowing should be started soon after received of good amount of early monsoon showers (20mm per week) after 15-16 June where direct seeding is practiced. Incorporate FYM or compost @ 2000 Kg/acre. Apply 27 kg DAP and 20 kg MoP per acre during last ploughing. Before sowing seed should be treated with Trichoderma viride @ 10 g or Carbendazim 50% WP @ 1.5 g/kg seed.

Transplanting rice:

After received of sufficient rainfall, farmers are advised for sowing of dhanicha seeds @ 12 kg/acre prior to the transplanting of paddy so that the dhanicha crop will be 3-4 weeks old before 10-15 days of transplanting of rice. Apply all phosphatic fertilizers of Kharif rice to this green manuring crop in advance. Farmers who are sowing paddy nurseries are advised to treat the seeds with carbendizem @ 3 g/kg seed followed by shade drying for 24 hours before sowing in case of dry nursery and in case of wet nursery seeds should be treated with carbendizem @ 1 g/l of water by soaking for 24 hrs and another 24 hrs for sprouting and the seeds should be sown in wet nurseries. Raise the nursery for transplanted kharif rice. 400 m² area is required for one acre land. Prepare a 3 feet width and 6 inch raised bed with suitable length keeping 1 feet between two beds for irrigation and drainage. 40 baskets of FYM, 12 kg SSP and 2 kg MoP is required for 400 m² nursery area.

Suagrcane:

For **Red rot and wilt disease**, spray the crop with Mancozeb 0.3% or carbendazim 0.1% or Azoxystrobin 18.2%SC/ Difenconazole 11.45SC @ 1ml per liter water in dry days.

VEGETABLES AND FRUITS:

- Fruit and shoot borer** in **Brinjal**- Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@300g/acre at 15 days interval.
- Due to Past rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- After cessation of rain, **downy mildew disease** will appear in **cucurbits**. To control it, go for alternate spray of Metalaxyl+Mancozeb@2g/litre and Azoxystrobin+Tebuconazole @0.6g/litre of water at 10 days interval.
- Rust** disease is likely to appear in **Cowpea**. Spray Chlorothalonil@2g/litre of water.
- Wilting** may observe in **tomato and brinjal** field, to control it drenching with Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water.
- Current weather favours spread of **YMV** in **Ladies finger**. In case of appearance of diseases, rogue out affected plants and spray Thiometoxam 25WG @ 60g/acre.

Fruit:

Banana: For **leaf spot/sigatoka** as a precautionary measure spray carbendizim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalonil @ 2.0 g / litre of water. Drenching soil with Metiram 75% WG @ 1.5g/4 lit or carbendazim 12%+ mancozeb 63% @ 2g/lit to prevent **Panama wilt**.

Mango: Harvest the mature mango fruits during morning or evening hours. Keep the harvested fruits in shade to prevent from spongy tissue disease and heat.

Papaya: After rain due to water logging there may be increase in the infestation of **phytophthora crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

Animal Husbandry:

Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination. Keep the animal shed clean and dry.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)



Bulletin No.-413
Date- 30/06/2023

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

AGROMET ADVISORY BULLETIN FOR KANTAPADA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 15mm and district received average 10mm rainfall during the last week. The mean daily maximum and minimum temperature was **31.42°C(-1.62°C from normal)** and **26°C(+1.8°C from normal)**, respectively. Mean relative humidity of the district was around 87-90%.

Forecasted period: Today(8.30IST) to 05th July (8.30IST)2023

Weather advisory: Very light to light rainfall is forecasted for today to Sunday and Moderate rainfall/ thundershower is forecasted for other 2 days in the block. The day temperature will be **34-36°C** and night temperature will be **27-30°C** for upcoming 5 days. Maximum and minimum relative humidity will be 62-81% and 35-41% respectively. **There will be partly cloudy to overcast sky for next 5days.**

Block	Kantapada (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	30/06/2023	01/07/2023	02/07/2023	03/07/2023	04/07/2023
Rainfall(mm)	5	4	10	20	30
T-Max(°C)	34	36	35	36	36
T-Min(°C)	29	29	30	28	27
Cloud Cover(octa)	5	3	7	6	6
RH Max (%)	71	65	62	71	81
RH Min (%)	41	35	37	35	38
Wind Speed (Kmph)	17	16	15	18	18
Wind Direction (deg)	207	205	198	203	180

Agromet Advisory

General advisory: Farmer should wait for sufficient amount of rainfall (75-100 mm in a week) for sowing of Karif Rice.

Kharif rice:

Sowing should be started soon after received of good amount of early monsoon showers (20mm per week) after 15-16 June where direct seeding is practiced. Incorporate FYM or compost @ 2000 Kg/acre. Apply 27 kg DAP and 20 kg MoP per acre during last ploughing. Before sowing seed should be treated with Trichoderma viride @ 10 g or Carbendazim 50% WP @ 1.5 g/kg seed.

Transplanting rice:

After received of sufficient rainfall, farmers are advised for sowing of dhanicha seeds @ 12 kg/acre prior to the transplanting of paddy so that the dhanicha crop will be 3-4 weeks old before 10-15 days of transplanting of rice. Apply all phosphatic fertilizers of Kharif rice to this green manuring crop in advance. Farmers who are sowing paddy nurseries are advised to treat the seeds with carbendazim @ 3 g/kg seed followed by shade drying for 24 hours before sowing in case of dry nursery and in case of wet nursery seeds should be treated with carbendazim @ 1 g/l of water bysoaking for 24 hrs and another 24 hrs for sprouting and the seeds should be sown in wet nurseries. Raise the nursery for transplanted kharif rice. 400 m2 area is required for one acre land. Prepare a 3 feet width and 6 inch raised bed with suitable length keeping 1 feet between two beds for irrigation and drainage. 40 baskets of FYM, 12 kg SSP and 2 kg MoP is required for 400 m2 nursery area.

Suagrcane:

For **Red rot and wilt disease**, spray the crop with Mancozeb 0.3% or carbendazim 0.1% or Azoxystrobin 18.2%SC/ Difenconazole 11.45SC @ 1ml per liter water in dry days.

VEGETABLES AND FRUITS:

- Fruit and shoot borer** in **Brinjal**- Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@300g/acre at 15 days interval.
- Due to Past rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- After cessation of rain, **downy mildew disease** will appear in **cucurbits**. To control it, go for alternate spray of Metalaxyl+Mancozeb@2g/litre and Azoxystrobin+Tebuconazole @0.6g/litre of water at 10 days interval.
- Rust** disease is likely to appear in **Cowpea**. Spray Chlorothalonil@2g/litre of water.
- Wilting** may observe in **tomato and brinjal** field, to control it drenching with Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water.
- Current weather favours spread of **YMV** in **Ladies finger**. In case of appearance of diseases, rogue out affected plants and spray Thiometoxam 25WG @ 60g/acre.

Fruit:

Banana: For **leaf spot/sigatoka** as a precautionary measure spray carbendazim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalonil @ 2.0 g / litre of water. Drenching soil with Metiram 75% WG @ 1.5g/4 lit or carbendazim 12%+ mancozeb 63% @ 2g/lit to prevent **Panama wilt**.

Mango: Harvest the mature mango fruits during morning or evening hours. Keep the harvested fruits in shade to prevent from spongy tissue disease and heat.

Papaya: After rain due to water logging there may be increase in the infestation of **phytophthora crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

Animal Husbandry:

Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination. Keep the animal shed clean and dry.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA INDIA METEOROLOGICAL DEPARTMENT DISTRICT AGROMET UNIT KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

AGROMET ADVISORY BULLETIN FOR MAHANGA



Bulletin No.-413

Date- 30/06/2023

District Past week average weather condition and Block past week average Rainfall condition

The block received average 06mm and district received average 10mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.42°C (-1.62°C from normal) and 26°C (+1.8°C from normal), respectively. Mean relative humidity of the district was around 87-90%.

Forecasted period: Today (8.30 IST) to 05th July (8.30 IST) 2023

Weather advisory: Very light to light rainfall is forecasted for today to Sunday and Moderate rainfall/ thundershower is forecasted for other 2 days in the block. The day temperature will be and it will be 34-37°C and night temperature will be 27-29°C for upcoming 5 days. Maximum and minimum relative humidity will be 64-81% and 37-41% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Mahanga (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	30/06/2023	01/07/2023	02/07/2023	03/07/2023	04/07/2023
Rainfall (mm)	5	4	10	20	30
T-Max (°C)	34	35	35	36	37
T-Min (°C)	28	29	29	28	27
Cloud Cover (octal)	5	4	7	6	6
RH Max (%)	71	65	64	72	81
RH Min (%)	41	38	39	37	40
Wind Speed (Kmph)	16	15	16	17	18
Wind Direction (deg)	207	203	198	189	167

Agromet Advisory

General advisory: Farmer should wait for sufficient amount of rainfall (75-100 mm in a week) for sowing of Kharif Rice.

Kharif rice:

Sowing should be started soon after received of good amount of early monsoon showers (20mm per week) after 15-16 June where direct seeding is practiced. Incorporate FYM or compost @ 2000 Kg/acre. Apply 27 kg DAP and 20 kg MoP per acre during last ploughing. Before sowing seed should be treated with Trichoderma viride @ 10 g or Carbendazim 50% WP @ 1.5 g/kg seed.

Transplanting rice:

After received of sufficient rainfall, farmers are advised for sowing of dhanicha seeds @ 12 kg/acre prior to the transplanting of paddy so that the dhanicha crop will be 3-4 weeks old before 10-15 days of transplanting of rice. Apply all phosphatic fertilizers of Kharif rice to this green manuring crop in advance. Farmers who are sowing paddy nurseries are advised to treat the seeds with carbendazim @ 3 g/kg seed followed by shade drying for 24 hours before sowing in case of dry nursery and in case of wet nursery seeds should be treated with carbendazim @ 1 g/l of water by soaking for 24 hrs and another 24 hrs for sprouting and the seeds should be sown in wet nurseries. Raise the nursery for transplanted kharif rice. 400 m² area is required for one acre land. Prepare a 3 feet width and 6 inch raised bed with suitable length keeping 1 feet between two beds for irrigation and drainage. 40 baskets of FYM, 12 kg SSP and 2 kg MoP is required for 400 m² nursery area.

Suagrcane: For Red rot and wilt disease, spray the crop with Mancozeb 0.3% or carbendazim 0.1% or Azoxystrobin 18.2% SC/ Difenconazole 11.45 SC @ 1ml per liter water in dry days.

VEGETABLES AND FRUITS:

- Fruit and shoot borer** in Brinjal- Spray Neem based pesticide 1500ppm @ 600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45% SC @ 70ml/acre and Thiodicarb @ 300g/acre at 15 days interval.
- Due to Past rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- After cessation of rain, **downy mildew disease** will appear in **cucurbits**. To control it, go for alternate spray of Metalaxyl + Mancozeb @ 2g/litre and Azoxystrobin + Tebuconazole @ 0.6g/litre of water at 10 days interval.
- Rust** disease is likely to appear in **Cowpea**. Spray Chlorothalonil @ 2g/litre of water.
- Wilting** may observe in **tomato and brinjal** field, to control it drenching with Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water.
- Current weather favours spread of **YMV** in **Ladies finger**. In case of appearance of diseases, rogue out affected plants and spray Thiometoxam 25WG @ 60g/acre.

Fruit:

Banana: For leaf spot/sigatoka as a precautionary measure spray carbendazim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalonil @ 2.0 g / litre of water. Drenching soil with Metiram 75% WG @ 1.5g/4 lit or carbendazim 12% + mancozeb 63% @ 2g/lit to prevent Panama wilt.

Mango: Harvest the mature mango fruits during morning or evening hours. Keep the harvested fruits in shade to prevent from spongy tissue disease and heat.

Papaya: After rain due to water logging there may be increase in the infestation of **phytophthora crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70% wp 2-4g/ltr of water.

Animal Husbandry:

Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination. Keep the animal shed clean and dry.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)



Bulletin No.-413
Date- 30/06/2023

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

AGROMET ADVISORY BULLETIN FOR NARASINGHPUR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 13mm and district received average 10mm rainfall during the last week. The mean daily maximum and minimum temperature was **31.42°C(-1.62°C from normal)** and **26°C(+1.8°C from normal)**, respectively. Mean relative humidity of the district was around 87-90%.

Forecasted period: Today(8.30IST) to 05th July (8.30IST)2023

Weather advisory: Very light to light rainfall is forecasted for today to Sunday and Moderate rainfall/ thundershower is forecasted for other 2 days in the block. The day temperature will be **34-37°C** and night temperature will be **25-28°C** for upcoming 5 days. Maximum and minimum relative humidity will be 64-83% and 36-41% respectively. **There will be partly cloudy to overcast sky for next 5 days.**

Block	Narasinghpur (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	30/06/2023	01/07/2023	02/07/2023	03/07/2023	04/07/2023
Rainfall(mm)	5	4	10	20	30
T-Max(°C)	34	35	37	36	37
T-Min(°C)	26	27	28	27	25
Cloud Cover(octa)	6	4	7	6	6
RH Max (%)	72	67	64	68	83
RH Min (%)	41	38	37	36	38
Wind Speed (Kmph)	10	10	10	10	9
Wind Direction (deg)	169	225	210	198	167

Agromet Advisory

General advisory: Farmer should wait for sufficient amount of rainfall (75-100 mm in a week) for sowing of Kharif Rice.

Kharif rice:

Sowing should be started soon after receive of good amount of early monsoon showers (**75-100 mm in a week**) where direct seeding is practiced. Incorporate FYM or compost @ 2000 Kg/acre. Apply 27 kg DAP and 20 kg MoP per acre during last ploughing. Before sowing seed should be treated with Trichoderma viride @ 10 g or Carbendazim 50% WP @ 1.5 g/kg seed.

Transplanting rice:

After receive of sufficient rainfall, farmers are advised for sowing of dhanicha seeds @ 12 kg/acre prior to the transplanting of paddy so that the dhanicha crop will be 3-4 weeks old before 10-15 days of transplanting of rice. Apply all phosphatic fertilizers of Kharif rice to this green manuring crop in advance. Farmers who are sowing paddy nurseries are advised to treat the seeds with carbendazim @ 3 g/kg seed followed by shade drying for 24 hours before sowing in case of dry nursery and in case of wet nursery seeds should be treated with carbendazim @ 1 g/l of water by soaking for 24 hrs and another 24 hrs for sprouting and the seeds should be sown in wet nurseries. Raise the nursery for transplanted kharif rice. 400 m² area is required for one acre land. Prepare a 3 feet width and 6 inch raised bed with suitable length keeping 1 feet between two beds for irrigation and drainage. 40 baskets of FYM, 12 kg SSP and 2 kg MoP is required for 400 m² nursery area.

Maize: It is time for sowing of maize for green cob purpose under irrigated condition. Grow high yielding varieties like Navjot, Shakti, Decan-107, Decan-109, Use 6 kg seeds/acre. Apply one third of N and all P & K of the recommended dose of 50:24:24 kg. NPK/acre. Sow the seeds in lines with spacing of 45-60X20 cm at 5 cm soil depth.

VEGETABLES AND FRUITS:

- Fruit and shoot borer** in **Brinjal**- Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@300g/acre at 15 days interval.
- Due to Past rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- After cessation of rain, **downy mildew disease** will appear in **cucurbits**. To control it, go for alternate spray of Metalaxyl+Mancozeb@2g/litre and Azoxystrobin+Tebuconazole @0.6g/litre of water at 10 days interval.
- Rust** disease is likely to appear in **Cowpea**. Spray Chlorothalonil@2g/litre of water.
- Wilting** may observe in **tomato and brinjal** field, to control it drenching with Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water.
- Current weather favours spread of **YMV** in **Ladies finger**. In case of appearance of diseases, rogue out affected plants and spray Thiometoxam 25WG @ 60g/acre.

Fruit:

Banana: For leaf spot/sigatoka as a precautionary measure spray carbendazim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalonil @ 2.0 g / litre of water. Drenching soil with Metiram 75% WG @ 1.5g/4 lit or carbendazim 12%+ mancozeb 63% @ 2g/lit to prevent Panama wilt.

Mango: Harvest the mature mango fruits during morning or evening hours. Keep the harvested fruits in shade to prevent from spongy tissue disease and heat.

Papaya: After rain due to water logging there may be increase in the infestation of **phytophthora crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

Animal Husbandry: Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination. Keep the animal shed clean and dry.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-413
Date- 30/06/2023

AGROMET ADVISORY BULLETIN FOR NIALI

District Past week average weather condition and Block past week average Rainfall condition

The block received average 06mm and district received average 10mm rainfall during the last week. The mean daily maximum and minimum temperature was **31.42°C (-1.62°C from normal)** and **26°C (+1.8°C from normal)**, respectively. Mean relative humidity of the district was around 87-90%.

Forecasted period: Today (8.30IST) to 05th July (8.30IST) 2023

Weather advisory: *Very light to light rainfall is forecasted for today to Sunday and Moderate rainfall/ thundershower is forecasted for other 2 days in the block.* The day temperature will be **34-37°C** and night temperature will be **27-30°C** for upcoming 5 days. Maximum and minimum relative humidity will be 65-81% and 37-43% respectively. **There will be partly cloudy to overcast sky for next 5 days.**

Block	Niali (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	30/06/2023	01/07/2023	02/07/2023	03/07/2023	04/07/2023
Rainfall (mm)	5	4	10	20	30
T-Max (°C)	34	36	36	37	36
T-Min (°C)	28	29	30	28	27
Cloud Cover (octa)	5	3	7	6	6
RH Max (%)	72	67	65	73	81
RH Min (%)	43	37	37	37	41
Wind Speed (Kmph)	17	16	16	19	18
Wind Direction (deg)	204	208	203	199	180

Agromet Advisory

General advisory: **Farmer should wait for sufficient amount of rainfall (75-100 mm in a week) for sowing of Kharif Rice.**

Kharif rice:

Sowing should be started soon after received of good amount of early monsoon showers (20mm per week) after 15-16 June where direct seeding is practiced. Incorporate FYM or compost @ 2000 Kg/acre. Apply 27 kg DAP and 20 kg MoP per acre during last ploughing. Before sowing seed should be treated with Trichoderma viride @ 10 g or Carbendazim 50% WP @ 1.5 g/kg seed.

Transplanting rice:

After received of sufficient rainfall, farmers are advised for sowing of dhanicha seeds @ 12 kg/acre prior to the transplanting of paddy so that the dhanicha crop will be 3-4 weeks old before 10-15 days of transplanting of rice. Apply all phosphatic fertilizers of Kharif rice to this green manuring crop in advance. Farmers who are sowing paddy nurseries are advised to treat the seeds with carbendazim @ 3 g/kg seed followed by shade drying for 24 hours before sowing in case of dry nursery and in case of wet nursery seeds should be treated with carbendazim @ 1 g/l of water by soaking for 24 hrs and another 24 hrs for sprouting and the seeds should be sown in wet nurseries. Raise the nursery for transplanted kharif rice. 400 m² area is required for one acre land. Prepare a 3 feet width and 6 inch raised bed with suitable length keeping 1 feet between two beds for irrigation and drainage. 40 baskets of FYM, 12 kg SSP and 2 kg MoP is required for 400 m² nursery area.

Suagrcane:

For **Red rot and wilt disease**, spray the crop with Mancozeb 0.3% or carbendazim 0.1% or Azoxystrobin 18.2% SC/ Difenconazole 11.45SC @ 1ml per liter water in dry days.

VEGETABLES AND FRUITS:

- Fruit and shoot borer** in **Brinjal**- Spray Neem based pesticide 1500ppm @ 600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45% SC @ 70ml/acre and Thiodicarb @ 300g/acre at 15 days interval.
- Due to Past rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- After cessation of rain, **downy mildew disease** will appear in **cucurbits**. To control it, go for alternate spray of Metalaxyl+Mancozeb @ 2g/litre and Azoxystrobin+Tebuconazole @ 0.6g/litre of water at 10 days interval.
- Rust** disease is likely to appear in **Cowpea**. Spray Chlorothalonil @ 2g/litre of water.
- Wilting** may observe in **tomato and brinjal** field, to control it drenching with Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water.
- Current weather favours spread of **YMV** in **Ladies finger**. In case of appearance of diseases, rogue out affected plants and spray Thiometoxam 25WG @ 60g/acre.

Fruit:

Banana: For **leaf spot/sigatoka** as a precautionary measure spray carbendazim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalonil @ 2.0 g / litre of water. Drenching soil with Metiram 75% WG @ 1.5g/4 lit or carbendazim 12%+ mancozeb 63% @ 2g/lit to prevent **Panama wilt**.

Mango: Harvest the mature mango fruits during morning or evening hours. Keep the harvested fruits in shade to prevent from spongy tissue disease and heat.

Papaya: After rain due to water logging there may be increase in the infestation of **phytophthora crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

Animal Husbandry: Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination. Keep the animal shed clean and dry.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-413
Date- 30/06/2023

District Past week average weather condition and Block past week average Rainfall condition

The block received average 05mm and district received average 10mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.42°C (-1.62°C from normal) and 26°C (+1.8°C from normal), respectively. Mean relative humidity of the district was around 87-90%.

Forecasted period: Today(8.30IST) to 05th July (8.30IST)2023

Weather advisory: *Very light to light rainfall is forecasted for today to Sunday and Moderate rainfall/ thundershower is forecasted for other 2 days in the block.* The day temperature will be 34-37°C and night temperature will be 27-29°C for upcoming 5 days. Maximum and minimum relative humidity will be 64-81% and 36-41% respectively. **There will be partly cloudy to overcast sky for next 5 days.**

Block	Nischintakoili (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	30/06/2023	01/07/2023	02/07/2023	03/07/2023	04/07/2023
Rainfall(mm)	5	4	10	20	30
T-Max(°C)	34	35	35	36	37
T-Min(°C)	28	29	29	28	27
Cloud Cover(octa)	5	4	7	6	5
RH Max (%)	70	65	64	71	81
RH Min (%)	41	37	39	36	39
Wind Speed (Kmph)	16	15	16	18	18
Wind Direction (deg)	207	203	198	189	167

Agromet Advisory

General advisory: **Farmer should wait for sufficient amount of rainfall (75-100 mm in a week) for sowing of Karif Rice.**

Kharif rice:

Sowing should be started soon after received of good amount of early monsoon showers (20mm per week) after 15-16 June where direct seeding is practiced. Incorporate FYM or compost @ 2000 Kg/acre. Apply 27 kg DAP and 20 kg MoP per acre during last ploughing. Before sowing seed should be treated with Trichoderma viride @ 10 g or Carbendazim 50% WP @ 1.5 g/kg seed.

Transplanting rice:

After received of sufficient rainfall, farmers are advised for sowing of dhanicha seeds @ 12 kg/acre prior to the transplanting of paddy so that the dhanicha crop will be 3-4 weeks old before 10-15 days of transplanting of rice. Apply all phosphatic fertilizers of Kharif rice to this green manuring crop in advance. Farmers who are sowing paddy nurseries are advised to treat the seeds with carbendazim @ 3 g/kg seed followed by shade drying for 24 hours before sowing in case of dry nursery and in case of wet nursery seeds should be treated with carbendazim @ 1 g/l of water by soaking for 24 hrs and another 24 hrs for sprouting and the seeds should be sown in wet nurseries. Raise the nursery for transplanted kharif rice. 400 m² area is required for one acre land. Prepare a 3 feet width and 6 inch raised bed with suitable length keeping 1 feet between two beds for irrigation and drainage. 40 baskets of FYM, 12 kg SSP and 2 kg MoP is required for 400 m² nursery area.

Suagrcane:

For **Red rot and wilt disease**, spray the crop with Mancozeb 0.3% or carbendazim 0.1% or Azoxystrobin 18.2%SC/ Difenoconazole 11.45SC @ 1ml per liter water in dry days.

VEGETABLES AND FRUITS:

- **Fruit and shoot borer** in **Brinjal**- Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@300g/acre at 15 days interval.
- Due to Past rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- After cessation of rain, **downy mildew disease** will appear in **cucurbits**. To control it, go for alternate spray of Metalaxyl+Mancozeb@2g/litre and Azoxystrobin+Tebuconazole @0.6g/litre of water at 10 days interval.
- **Rust** disease is likely to appear in **Cowpea**. Spray Chlorothalonil@2g/litre of water.
- **Wilting** may observe in **tomato and brinjal** field, to control it drenching with Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water.
- Current weather favours spread of **YMV** in **Ladies finger**. In case of appearance of diseases, rogue out affected plants and spray Thiomethoxam 25WG @ 60g/acre.

Fruit:

Banana: For **leaf spot/sigatoka** as a precautionary measure spray carbendazim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalonil @ 2.0 g/ litre of water. Drenching soil with Metiram 75% WG @ 1.5g/4 lit or carbendazim 12%+ mancozeb 63% @ 2g/lit to prevent **Panama wilt**.

Mango: Harvest the mature mango fruits during morning or evening hours. Keep the harvested fruits in shade to prevent from spongy tissue disease and heat.

Papaya: After rain due to water logging there may be increase in the infestation of **phytophthora crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

Animal Husbandry: Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination. Keep the animal shed clean and dry.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-413
Date- 30/06/2023

AGROMET ADVISORY BULLETIN FOR SALEPUR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 04mm and district received average 10mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.42°C (-1.62°C from normal) and 26°C (+1.8°C from normal), respectively. Mean relative humidity of the district was around 87-90%.

Forecasted period: Today(8.30IST) to 05th July (8.30IST)2023

Weather advisory: Very light to light rainfall is forecasted for today to Sunday and Moderate rainfall/ thundershower is forecasted for other 2 days in the block. The day temperature will be 34-37°C and night temperature will be 27-30°C for upcoming 5 days. Maximum and minimum relative humidity will be 64-81% and 36-41% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Salepur (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	30/06/2023	01/07/2023	02/07/2023	03/07/2023	04/07/2023
Rainfall(mm)	5	4	10	20	30
T-Max(°C)	34	35	35	36	37
T-Min(°C)	29	29	30	28	27
Cloud Cover(octa)	5	4	7	6	5
RH Max (%)	70	65	64	71	81
RH Min (%)	41	37	38	36	39
Wind Speed (Kmph)	16	15	15	17	18
Wind Direction (deg)	207	203	198	189	180

Agromet Advisory

General advisory: Farmer should wait for sufficient amount of rainfall (75-100 mm in a week) for sowing of Karif Rice.

Kharif rice:

Sowing should be started soon after receive of good amount of early monsoon showers (75-100 mm in a week) where direct seeding is practiced. Incorporate FYM or compost @ 2000 Kg/acre. Apply 27 kg DAP and 20 kg MoP per acre during last ploughing. Before sowing seed should be treated with Trichoderma viride @ 10 g or Carbendazim 50% WP @ 1.5 g/kg seed.

Transplanting rice:

After receive of sufficient rainfall, farmers are advised for sowing of dhanicha seeds @ 12 kg/acre prior to the transplanting of paddy so that the dhanicha crop will be 3-4 weeks old before 10-15 days of transplanting of rice. Apply all phosphatic fertilizers of Kharif rice to this green manuring crop in advance. Farmers who are sowing paddy nurseries are advised to treat the seeds with carbendazim @ 3 g/kg seed followed by shade drying for 24 hours before sowing in case of dry nursery and in case of wet nursery seeds should be treated with carbendazim @ 1 g/l of water by soaking for 24 hrs and another 24 hrs for sprouting and the seeds should be sown in wet nurseries. Raise the nursery for transplanted kharif rice. 400 m² area is required for one acre land. Prepare a 3 feet width and 6 inch raised bed with suitable length keeping 1 feet between two beds for irrigation and drainage. 40 baskets of FYM, 12 kg SSP and 2 kg MoP is required for 400 m² nursery area.

Maize: It is time for sowing of maize for green cob purpose under irrigated condition. Grow high yielding varieties like Navjot, Shakti, Decan-107, Decan-109. Use 6 kg seeds/acre. Apply one third of N and all P & K of the recommended dose of 50:24:24 kg. NPK/acre. Sow the seeds in lines with spacing of 45-60X20 cm at 5 cm soil depth.

VEGETABLES AND FRUITS:

- Fruit and shoot borer** in **Brinjal**- Spray Neem based pesticide 1500ppm @ 600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC @ 70ml/acre and Thiodicarb @ 300g/acre at 15 days interval.
- Due to Past rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- After cessation of rain, **downy mildew disease** will appear in **cucurbits**. To control it, go for alternate spray of Metalaxyl+Mancozeb @ 2g/litre and Azoxystrobin+Tebuconazole @ 0.6g/litre of water at 10 days interval.
- Rust** disease is likely to appear in **Cowpea**. Spray Chlorothalonil @ 2g/litre of water.
- Wilting** may observe in **tomato and brinjal** field, to control it drenching with Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water.
- Current weather favours spread of **YMV** in **Ladies finger**. In case of appearance of diseases, rogue out affected plants and spray Thiometoxam 25WG @ 60g/acre.

Fruit:

Banana: For leaf spot/sigatoka as a precautionary measure spray carbendazim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalonil @ 2.0 g / litre of water. Drenching soil with Metiram 75% WG @ 1.5g/4 lit or carbendazim 12%+ mancozeb 63% @ 2g/lit to prevent Panama wilt.

Mango: Harvest the mature mango fruits during morning or evening hours. Keep the harvested fruits in shade to prevent from spongy tissue disease and heat.

Papaya: After rain due to water logging there may be increase in the infestation of **phytophthora crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

Animal Husbandry: Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination. Keep the animal shed clean and dry.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA INDIA METEOROLOGICAL DEPARTMENT DISTRICT AGROMET UNIT KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

AGROMET ADVISORY BULLETIN FOR TANGI-CHOUDWAR



Bulletin No.-413

Date- 30/06/2023

District Past week average weather condition and Block past week average Rainfall condition

The block received average 08mm and district received average 10mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.42°C (-1.62°C from normal) and 26°C (+1.8°C from normal), respectively. Mean relative humidity of the district was around 87-90%.

Forecasted period: Today (8.30IST) to 05th July (8.30IST) 2023

Weather advisory: Very light to light rainfall is forecasted for today to Sunday and Moderate rainfall/ thundershower is forecasted for other 2 days in the block. The day temperature will be 36-38°C and night temperature will be 29-31°C for upcoming 5 days. Maximum and minimum relative humidity will be 70-83% and 45-52% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Tangi-Choudwar (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	30/06/2023	01/07/2023	02/07/2023	03/07/2023	04/07/2023
Rainfall(mm)	5	4	10	20	30
T-Max(°C)	36	38	38	38	37
T-Min(°C)	30	31	31	30	29
Cloud Cover(octa)	5	3	6	5	6
RH Max (%)	76	70	72	79	83
RH Min (%)	48	45	45	46	52
Wind Speed (Kmph)	20	19	20	23	20
Wind Direction (deg)	204	199	198	198	180

Agromet Advisory

General advisory: Farmer should wait for sufficient amount of rainfall (75-100 mm in a week) for sowing of Kharif Rice.

Kharif rice:

Sowing should be started soon after receive of good amount of early monsoon showers (75-100 mm in a week) where direct seeding is practiced. Incorporate FYM or compost @ 2000 Kg/acre. Apply 27 kg DAP and 20 kg MoP per acre during last ploughing. Before sowing seed should be treated with Trichoderma viride @ 10 g or Carbendazim 50% WP @ 1.5 g/kg seed.

Transplanting rice:

After receive of sufficient rainfall, farmers are advised for sowing of dhanicha seeds @ 12 kg/acre prior to the transplanting of paddy so that the dhanicha crop will be 3-4 weeks old before 10-15 days of transplanting of rice. Apply all phosphatic fertilizers of Kharif rice to this green manuring crop in advance. Farmers who are sowing paddy nurseries are advised to treat the seeds with carbendazim @ 3 g/kg seed followed by shade drying for 24 hours before sowing in case of dry nursery and in case of wet nursery seeds should be treated with carbendazim @ 1 g/l of water by soaking for 24 hrs and another 24 hrs for sprouting and the seeds should be sown in wet nurseries. Raise the nursery for transplanted kharif rice. 400 m² area is required for one acre land. Prepare a 3 feet width and 6 inch raised bed with suitable length keeping 1 feet between two beds for irrigation and drainage. 40 baskets of FYM, 12 kg SSP and 2 kg MoP is required for 400 m² nursery area.

Maize:

It is time for sowing of maize for green cob purpose under irrigated condition. Grow high yielding varieties like Navjot, Shakti, Decan-107, Decan-109, Use 6 kg seeds/acre. Apply one third of N and all P & K of the recommended dose of 50:24:24 kg. NPK/acre. Sow the seeds in lines with spacing of 45-60X20 cm at 5 cm soil depth.

VEGETABLES AND FRUITS:

- Fruit and shoot borer** in **Brinjal**- Spray Neem based pesticide 1500ppm @ 600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC @ 70ml/acre and Thiodicarb @ 300g/acre at 15 days interval.
- Due to Past rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- After cessation of rain, **downy mildew disease** will appear in **cucurbits**. To control it, go for alternate spray of Metalaxyl+Mancozeb @ 2g/litre and Azoxystrobin+Tebuconazole @ 0.6g/litre of water at 10 days interval.
- Rust** disease is likely to appear in **Cowpea**. Spray Chlorothalonil @ 2g/litre of water.
- Wilting** may observe in **tomato and brinjal** field, to control it drenching with Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water.
- Current weather favours spread of **YMV** in **Ladies finger**. In case of appearance of diseases, rogue out affected plants and spray Thiometoxam 25WG @ 60g/acre.

Fruit:

Banana: For leaf spot/sigatoka as a precautionary measure spray carbendazim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalonil @ 2.0 g / litre of water. Drenching soil with Metiram 75% WG @ 1.5g/4 lit or carbendazim 12%+ mancozeb 63% @ 2g/lit to prevent Panama wilt.

Mango: Harvest the mature mango fruits during morning or evening hours. Keep the harvested fruits in shade to prevent from spongy tissue disease and heat.

Papaya: After rain due to water logging there may be increase in the infestation of **phytophthora crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

Animal Husbandry: Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination. Keep the animal shed clean and dry.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA INDIA METEOROLOGICAL DEPARTMENT DISTRICT AGROMET UNIT KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-413

Date- 30/06/2023

AGROMET ADVISORY BULLETIN FOR TIGIRIA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 17mm and district received average 10mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.42°C (-1.62°C from normal) and 26°C (+1.8°C from normal), respectively. Mean relative humidity of the district was around 87-90%.

Forecasted period: Today (8.30 IST) to 05th July (8.30 IST) 2023

Weather advisory: Very light to light rainfall is forecasted for today to Sunday and Moderate rainfall/ thundershower is forecasted for other 2 days in the block. The day temperature will be 34-36°C and night temperature will be 26-29°C for upcoming 5 days. Maximum and minimum relative humidity will be 62-81% and 37-43% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Tigiria (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	30/06/2023	01/07/2023	02/07/2023	03/07/2023	04/07/2023
Rainfall (mm)	5	4	10	20	30
T-Max (°C)	34	35	35	36	36
T-Min (°C)	27	28	29	28	26
Cloud Cover (octa)	7	5	7	6	5
RH Max (%)	72	67	62	66	81
RH Min (%)	43	38	37	37	38
Wind Speed (Kmph)	13	12	13	14	16
Wind Direction (deg)	205	210	205	198	167

Agro met Advisory

General advisory: Farmer should wait for sufficient amount of rainfall (75-100 mm in a week) for sowing of Kharif Rice.

Kharif rice:

Sowing should be started soon after receive of good amount of early monsoon showers (75-100 mm in a week) where direct seeding is practiced. Incorporate FYM or compost @ 2000 Kg/acre. Apply 27 kg DAP and 20 kg MoP per acre during last ploughing. Before sowing seed should be treated with Trichoderma viride @ 10 g or Carbendazim 50% WP @ 1.5 g/kg seed.

Transplanting rice:

After receive of sufficient rainfall, farmers are advised for sowing of dhanicha seeds @ 12 kg/acre prior to the transplanting of paddy so that the dhanicha crop will be 3-4 weeks old before 10-15 days of transplanting of rice. Apply all phosphatic fertilizers of Kharif rice to this green manuring crop in advance. Farmers who are sowing paddy nurseries are advised to treat the seeds with carbendazim @ 3 g/kg seed followed by shade drying for 24 hours before sowing in case of dry nursery and in case of wet nursery seeds should be treated with carbendazim @ 1 g/l of water by soaking for 24 hrs and another 24 hrs for sprouting and the seeds should be sown in wet nurseries. Raise the nursery for transplanted kharif rice. 400 m² area is required for one acre land. Prepare a 3 feet width and 6 inch raised bed with suitable length keeping 1 feet between two beds for irrigation and drainage. 40 baskets of FYM, 12 kg SSP and 2 kg MoP is required for 400 m² nursery area.

Maize: It is time for sowing of maize for green cob purpose under irrigated condition. Grow high yielding varieties like Navjot, Shakti, Decan-107, Decan-109, Use 6 kg seeds/acre. Apply one third of N and all P & K of the recommended dose of 50:24:24 kg. NPK/acre. Sow the seeds in lines with spacing of 45-60X20 cm at 5 cm soil depth.

VEGETABLES AND FRUITS:

- Fruit and shoot borer** in **Brinjal**- Spray Neem based pesticide 1500ppm @ 600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC @ 70ml/acre and Thiodicarb @ 300g/acre at 15 days interval.
- Due to Past rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- After cessation of rain, **downy mildew disease** will appear in **cucurbits**. To control it, go for alternate spray of Metalaxyl+Mancozeb @ 2g/litre and Azoxystrobin+Tebuconazole @ 0.6g/litre of water at 10 days interval.
- Rust** disease is likely to appear in **Cowpea**. Spray Chlorothalonil @ 2g/litre of water.
- Wilting** may observe in **tomato and brinjal** field, to control it drenching with Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water.
- Current weather favours spread of **YMV** in **Ladies finger**. In case of appearance of diseases, rogue out affected plants and spray Thiometoxam 25WG @ 60g/acre.

Fruit:

Banana: For leaf spot/sigatoka as a precautionary measure spray carbendazim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalonil @ 2.0 g / litre of water. Drenching soil with Metiram 75% WG @ 1.5g/4 lit or carbendazim 12%+ mancozeb 63% @ 2g/lit to prevent Panama wilt.

Mango: Harvest the mature mango fruits during morning or evening hours. Keep the harvested fruits in shade to prevent from spongy tissue disease and heat.

Papaya: After rain due to water logging there may be increase in the infestation of **phytophthora crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

Animal Husbandry: Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination. Keep the animal shed clean and dry.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-413
 Date- 30/06/2023

				
MEGHDOOT APP	DAMINI APP	FACEBOOK (Damu Cuttack)	BLOCK SPECIFIC WHAT'S APP GROUPS	@DAMUCUTTACK

- Kindly install and use **“Meghdoot mobile App”** for Cuttack district weather forecast, warnings and weather based agromet advisory services.
<https://play.google.com/store/apps/details?id=com.aas.meghdoot>
- Kindly install and use **“Damini mobile App”** for location specific for lightening forecast and warnings.
<https://play.google.com/store/apps/details?id=com.lightening.live.damini>
- Follow the **block specific whats app groups** and join in your block wise groups for getting weather based agromet advisory services twice in a week (Tuesday and Friday) and other weather warning based farm management advisories.
- Follow the **Official Website links** to get the weather based agromet advisory service bulletins (**English & Odia**) twice in a week (Tuesday and Friday).

1	ICAR-NRRI, Cuttack, Odisha	https://icar-nrri.in/agromate/	
2	Agrimet division, IMD, Pune	https://imdagrmet.gov.in/imdproject/AGIndex.php	

Dr Sujata Sethy
 (Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
 SMS (Agrometeorology)
 Contact No-9861445080/7008453427