




Cuttack District Block Wise Agro-met Advisory Service Bulletin

*Tuesday, 22nd August, 2023
(For the period 22nd to 26th August,
2023)*



*Issued by
District Agro-met Unit,
Krishi Vigyan Kendra Cuttack*






GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-429
Date- 22/08/2023

STANDARD PRECIPITATION INDEX AND NDVI MAP



LEGEND (Medium range block level rainfall forecast)		
	No Rainfall	00 mm
	Very light rainfall	01-05 mm
	Light rainfall	06-10 mm
	Light to moderate rainfall	11-30 mm
	Moderate to heavy rainfall	31-65 mm
	Heavy to very heavy rainfall	66-115 mm
	Extremely heavy rainfall	>116 mm

LEGEND (Past week temperature)		
Color code	Temperature	Departure
	Below Normal	< 2°C from normal
	Normal	-2°C to +2°C from normal
	Above Normal	>2°C from normal

Dr Sujata Sethy
 (Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
 SMS (Agrometeorology)
 Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA INDIA METEOROLOGICAL DEPARTMENT DISTRICT AGROMET UNIT KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)



Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com







Bulletin No.-429

Date- 22/08/2023

Past week (15-21 August, 2023) cumulative rainfall(mm) & its departure% from normal and (16-22 August) average maximum and minimum temperature for Cuttack district



Medium range (Today (8.30IST) to Sunday (8.30IST)) Rainfall (mm) forecast for Cuttack district



Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-429
Date- 22/08/2023

Met Sub division	Extended range outlook for subsequent 1 week (6 th to 12 th days) (27 August-02 September 2023) after 5 days				
Odisha	<i>Parameter</i>	<i>Actual</i>	<i>Normal</i>	<i>Departure (%)</i>	<i>Category</i>
	Rainfall (mm)	11.95	10.08	18.46	Normal
	<i>Parameter</i>	<i>Actual</i>	<i>Normal</i>	<i>Anomaly</i>	<i>Category</i>
	Maximum temperature (°C)	31.20	31.17	0.03	Normal
	Minimum temperature (°C)	24.62	24.17	0.45	Normal

***Agromet Advisory based upon subsequent week extended range forecast-:** As there will be **normal rainfall in the subsequent week (27 August-02 September, 2023) after 5 days in Odisha condition**, so farmers are advised to drain out excess water from paddy field and should maintain 5-10cm during transplanting/reproductive stage. Farmers may go for sowing of kharif vegetables before the subsequent week. In heavy weed infested areas, before the subsequent week, spray Bispyribac sodium 10% SC @ 120 ml/acre in 8 tanks of 16 litre capacity sprayer at 8-10 days after emergence of weeds (or when the weeds are at 2-3 leaf stage). **Farmers should pause irrigation, fertilizer/pesticide application; human labour engagement in subsequent week after 5days based upon rainfall forecast.**

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427













GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-429
 Date- 22/08/2023

Weather (Past and Forecasted) based Disease and pest in Major Crops and vegetables

Sl No	Crop	Stage and condition of the crop	Anticipated Disease and pest	
1	Rice	Tillering		Blast
				Sheath blight
				Stem Borer
				Gall midge
				Sawarming caterpillar
2	Sugarcane	Vegetative to Tillering		Red rot
				Wilt
				Early shoot borer
3	Maize	5 week stage		Stem fly
				Leaf blight

Dr Sujata Sethy
 (Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area














Debasish Jena
 SMS (Agrometeorology)
 Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-429
Date- 22/08/2023

				Stalk rot
3	Jute	30-50days		Stem rot
				Indigo caterpillar
4	Brinjal, tomatoto, chilli	seedling		Damping off
				Stem rot
5	Tomato and brinjal			Bacterial wilt
				White fly
				Jassid
6	okra	Fruting		cercospora leaf spot
7	Banana			Sigatoka
8	Papaya			Yellow mosaic virus

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA INDIA METEOROLOGICAL DEPARTMENT DISTRICT AGROMET UNIT KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-429

Date- 22/08/2023

AGROMET ADVISORY BULLETIN FOR ATHAGAD

District Past week average weather condition and Block past week average Rainfall condition

The block received average 12mm & district received average 10mm rainfall during the last week. The mean daily maximum and minimum temperature was 32.8°C (+1°C from normal) and 25.8°C (+1.7°C from normal), respectively. Mean relative humidity of the district was around 84-90%.

Forecasted period: Today(8.30IST) to 27th August (8.30IST)2023

Weather advisory: ; Light to moderate rainfall is forecasted in between next 5 days in the block. The day temperature will be 30-33°C and night temperature will be 24-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 83-86% and 50-63% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Athagad (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	22/08/2023	23/08/2023	24/08/2023	25/08/2023	26/08/2023
Rainfall (mm)	15	25	22	7	6
T-Max (°C)	33	32	30	31	32
T-Min (°C)	26	26	25	24	25
Cloud Cover (octa)	8	8	8	8	8
RH Max (%)	85	86	83	84	83
RH Min (%)	54	56	63	57	50
Wind Speed (kmph)	12	11	11	11	11
Wind Direction (deg)	204	243	249	283	287

Agromet Advisory

General advisory: Farmers should drain out excess water from vegetable and rice field near As relative humidity was above 80% and rainy day condition, there will be chances of disease and pest infestation mainly sheath blight, blight, blast, foot rot and swarming caterpillar, leaf folder, stem borer in rice. They may pause all agricultural activities based on rainfall forecast especially today to Thursday and resume it after.

Kharif Rice:

- The crop is at tillering stage. Maintain about 2-3 cm standing water during tillering stage. Problem of weeds may aggravate after transplanting, apply Bensulfuron methyl 0.6% + Pretilachlor 6% GR with sand mix at 1:1 ratio @ 4 kg/acre within 3-7 days after transplanting for controlling weeds.
- Intermittent dry and wet weather condition, is favorable for incidence of BLB and Sheath blight in early sown nursery seedlings. Go for foliar spray of Hexaconazole with Plantomycin @ 1.0 g/l to prevent the spread of the disease.
- If leaf blast is observed in nursery, spray Tebuconazole 50% + Trifloxystrobin 25% WG @ 0.4 g or Isoprothiolane 40EC @ 1.5 ml per litre of water. Repeat the spray at 7-10 days interval.
- In Gall midge affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre
- If swarming caterpillar will be found after the rain water recedes, apply Lambda cyhalothrin 5% EC @ 100 ml/acre.
- Temperature below 28deg c and continuous rainfall lead to infestation of stem borer at tillering stage of rice, to control this apply Azadirachtin 0.15% EC @ 800 ml/acre or Cartap hydrochloride 4G @ 10kg/acre.

Maize: Maize crop is highly sensitive to waterlogged conditions during seedling stage. Hence, drain out excess water from the field as early as possible.

- After cessation of rains apply booster dose of fertilizer.
- Weeds may be controlled after rainfall.
- Stem fly in Maize infects the plants from sowing to 5 weeks after sowing, temperature between 25-30 deg. celsius and humidity above 60% favour pest incidence. In vegetative stage to control stem fly spray Methyl demeton @ 1 ml/lit or Dimethoate @ 2ml/lit or apply Carbofuran 3G granules @ 14 kg/acre.
- High humid conditions, temperature between 18-27 Deg. Celsius favours the occurrence of leaf blight in Maize. To control spray Azoxystrobin + Mancozeb 300 g /ac. Due to water logging condition stem rot may be found, then farmers should apply carbendazim @ 2g/ltr water.

VEGETABLES: Due to past rain, Chances of fruit rot infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary, spray second and third round at 15 days interval.

- If incidence of Damping off is noticed in Tomato, Brinjal, Chilli nurseries, drench nursery bed with Copper-Oxy-Chloride @ 3 g per liter of water.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever sucking pests incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. To prevent of soil borne diseases like seedling rot, stem rot and wilt diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenconazole @ 1ml/lit at 15 days interval to control Cercospora leaf spot.

ANIMAL HUSBANDRY: Keep animal shed and shelter house free from water logging by keeping it dry and free from weed. Fresh and immature leaves should not be fed more amount as they may cause diarrhea. If grasses were cut in morning or evening time, it can be sun dried for about 30 minute before feeding.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-429

Date- 22/08/2023

AGROMET ADVISORY BULLETIN FOR BADAMBA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 7mm & district received average 10mm rainfall during the last week. The mean daily maximum and minimum temperature was **32.8°C (+1°C from normal)** and **25.8°C (+1.7°C from normal)**, respectively. Mean relative humidity of the district was around 84-90%.

Forecasted period: Today(8.30IST) to 27th August (8.30IST)2023

Weather advisory: Moderate to isolated heavy rainfall is forecasted for tomorrow & light to moderate rainfall is forecasted in between other 4 days in the block. The day temperature will be 30-33°C and night temperature will be 24-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 82-87% and 50-65% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Badamba (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	22/08/2023	23/08/2023	24/08/2023	25/08/2023	26/08/2023
Rainfall(mm)	18	32	21	7	6
T-Max(°C)	33	32	30	31	33
T-Min(°C)	26	25	25	24	25
Cloud Cover(octa)	8	8	8	8	8
RH Max (%)	85	87	82	82	82
RH Min (%)	54	58	65	58	50
Wind Speed (Kmph)	11	10	11	10	10
Wind Direction (deg)	206	243	249	252	279

Agromet Advisory

General advisory: Farmers should drain out excess water from vegetable and rice field near As relative humidity was above 80% and rainy day condition, there will be chances of disease and pest infestation mainly sheath blight, blight, blast, foot rot and swarming caterpillar, leaf folder, stem borer in rice. They may pause all agricultural activities based on rainfall forecast especially today to Thursday and resume it after.

Kharif Rice:

- The crop is at tillering stage. Maintain about 2-3 cm standing water during tillering stage. Problem of weeds may aggravate after transplanting, apply Bensulfuron methyl 0.6% + Pretilachlor 6% GR with sand mix at 1:1 ratio @ 4 kg/acre within 3-7 days after transplanting for controlling weeds.
- Intermittent dry and wet weather condition, is favorable for incidence of **BLB and Sheath blight** in early sown nursery seedlings. Go for foliar spray of Hexaconazole with Plantomycin @ 1.0 g/l to prevent the spread of the disease.
- If **leaf blast** is observed in nursery, spray Tebuconazole 50% + Trifloxystrobin 25% WG @ 0.4 g or Isoprothiolane 40EC @ 1.5 ml per litre of water. Repeat the spray at 7-10 days interval.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre
- If **swarming caterpillar** will be found after the rain water recedes, apply Lambda cyhalothrin 5% EC @ 100 ml/acre.
- Temperature below 28deg c and continuous rainfall lead to infestation of **stem borer** at tillering stage of rice, to control this apply Azadirachtin 0.15% EC @ 800 ml/acre or Cartap hydrochloride 4G @ 10kg/acre.

Maize: Maize crop is highly sensitive to waterlogged conditions during seedling stage. Hence, drain out excess water from the field as early as possible.

- After cessation of rains apply booster dose of fertilizer.
- Weeds may be controlled after rainfall.
- Stem fly** in Maize infects the plants from sowing to 5 weeks after sowing, temperature between 25-30 deg. celsius and humidity above 60% favour pest incidence. In vegetative stage to control stem fly spray Methyl demeton @ 1 ml/lit or Dimethoate @ 2ml/lit or apply Carbofuran 3G granules @ 14 kg/acre.
- High humid conditions, temperature between 18-27 Deg. Celsius favours the occurrence of **leaf blight** in Maize. To control spray Azoxystrobin + Mancozeb 300 g/ac. Due to water logging condition stem rot may be found, then farmers should apply carbendazim @ 2g/ltr water.

VEGETABLES: Due to past rain, Chances of **fruit rot** infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary, spray second and third round at 15 days interval.

- If incidence of **Damping off** is noticed in Tomato, Brinjal, Chilli nurseries, drench nursery bed with Copper-Oxy-Chloride @ 3 g per liter of water.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever **sucking pests** incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. To prevent of soil borne diseases like **seedling rot, stem rot and wilt** diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenconazole @ 1ml/lit at 15 days interval to control **Cercospora leaf spot**.

ANIMAL HUSBANDRY: Keep animal shed and shelter house free from water logging by keeping it dry and free from weed. Fresh and immature leaves should not to be fed more amount as they may cause diarrhea. If grasses were cut in morning or evening time, it can be sun dried for about 30 minute before feeding.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-429
Date- 22/08/2023

AGROMET ADVISORY BULLETIN FOR BANKI

District Past week average weather condition

The district received average 10mm rainfall during the last week. The mean daily maximum and minimum temperature was **32.8°C (+1°C from normal)** and **25.8°C (+1.7°C from normal)** respectively. Mean relative humidity of the district was around 84-90%.

Forecasted period: Today(8.30IST) to 27th August (8.30IST)2023

Weather advisory: *Moderate to isolated heavy rainfall is forecasted for tomorrow & light to moderate rainfall is forecasted in between other 4 days in the block.* The day temperature will be **30-33°C** and night temperature will be **24-26°C** for upcoming 5 days. Maximum and minimum relative humidity will be 80-85% and 48-60% respectively. **There will be partly cloudy to overcast sky for next 5 days.**

Block	Banki (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	22/08/2023	23/08/2023	24/08/2023	25/08/2023	26/08/2023
Rainfall(mm)	20	35	18	7	6
T-Max(°C)	33	32	30	31	33
T-Min(°C)	26	25	25	24	25
Cloud Cover(octa)	8	8	8	8	8
RH Max (%)	82	85	80	81	81
RH Min (%)	52	53	60	55	48
Wind Speed (Kmph)	13	12	12	11	11
Wind Direction (deg)	204	238	249	252	279

Agromet Advisory

General advisory: *Farmers should drain out excess water from vegetable and rice field near As relative humidity was above 80% and rainy day condition, there will be chances of disease and pest infestation mainly sheath blight, blight, blast, foot rot and swarming caterpillar, leaf folder, stem borer in rice. They may pause all agricultural activities based on rainfall forecast especially today to Thursday and resume it after.*

Kharif Rice:

- The crop is at tillering stage. Maintain about 2-3 cm standing water during tillering stage. Problem of weeds may aggravate after transplanting, apply Bensulfuron methyl 0.6% + Pretilachlor 6% GR with sand mix at 1:1 ratio @ 4 kg/acre within 3-7 days after transplanting for controlling weeds.
- Intermittent dry and wet weather condition, is favorable for incidence of **BLB and Sheath blight** in early sown nursery seedlings. Go for foliar spray of Hexaconazole with Plantomycin @ 1.0 g/l to prevent the spread of the disease.
- If **leaf blast** is observed in nursery, spray Tebuconazole 50% + Trifloxystrobin 25% WG @ 0.4 g or Isoprothiolane 40EC @ 1.5 ml per litre of water. Repeat the spray at 7-10 days interval.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre
- If **swarming caterpillar** will be found after the rain water recedes, apply Lambda cyhalothrin 5% EC @ 100 ml/acre.
- Temperature below 28deg c and continuous rainfall lead to infestation of **stem borer** at tillering stage of rice, to control this apply Azadirachtin 0.15% EC @ 800 ml/acre or Cartap hydrochloride 4G @ 10kg/acre.

Maize: Maize crop is highly sensitive to waterlogged conditions during seedling stage. Hence, drain out excess water from the field as early as possible.

- After cessation of rains apply booster dose of fertilizer.
- Weeds may be controlled after rainfall.
- Stem fly** in Maize infects the plants from sowing to 5 weeks after sowing, temperature between 25-30 deg. celsius and humidity above 60% favour pest incidence. In vegetative stage to control stem fly spray Methyl demeton @ 1 ml/lit or Dimethoate @ 2ml/lit or apply Carbofuran 3G granules @ 14 kg/acre.
- High humid conditions, temperature between 18-27 Deg. Celsius favours the occurrence of **leaf blight** in Maize. To control spray Azoxystrobin + Mancozeb 300 g /ac. **Due to water logging condition stem rot may be found, then farmers should apply carbendazim @ 2g/ltr water.**

VEGETABLES: Due to past rain, Chances of **fruit rot** infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary, spray second and third round at 15 days interval.

- If incidence of **Damping off** is noticed in Tomato, Brinjal, Chilli nurseries, drench nursery bed with Copper-Oxy-Chloride @ 3 g per liter of water.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever **sucking pests** incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. **To prevent of soil borne diseases like seedling rot, stem rot and wilt diseases apply 3g of Copper oxychloride per litre of water around the main stem.**
- Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenconazole @ 1ml/lit at 15 days interval to control **Cercospora leaf spot.**

ANIMAL HUSBANDRY: Keep animal shed and shelter house free from water logging by keeping it dry and free from weed. Fresh and immature leaves should not be fed more amount as they may cause diarrhea. If grasses were cut in morning or evening time, it can be sun dried for about 30 minute before feeding.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA INDIA METEOROLOGICAL DEPARTMENT DISTRICT AGROMET UNIT KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-429

Date- 22/08/2023

AGROMET ADVISORY BULLETIN FOR BANKI-DAMPARA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 27mm & district received average 10mm rainfall during the last week. The mean daily maximum and minimum temperature was 32.8°C (+1°C from normal) and 25.8°C (+1.7°C from normal), respectively. Mean relative humidity of the district was around 84-90%.

Forecasted period: Today(8.30IST) to 27th August (8.30IST)2023

Weather Advisory: Moderate to isolated heavy rainfall is forecasted for tomorrow & light to moderate rainfall is forecasted in between other 4 days in the block. The day temperature will be 30-33°C and night temperature will be 24-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 80-85% and 48-60% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Banki-Dampada (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	22/08/2023	23/08/2023	24/08/2023	25/08/2023	26/08/2023
Rainfall(mm)	18	36	20	7	6
T-Max(°C)	33	32	30	31	33
T-Min(°C)	26	25	25	24	25
Cloud Cover(octa)	8	8	8	8	8
RH Max (%)	83	85	80	82	81
RH Min (%)	52	54	60	55	48
Wind Speed (Kmph)	13	12	11	11	11
Wind Direction (deg)	204	238	249	252	279

Agromet Advisory

General advisory: Farmers should drain out excess water from vegetable and rice field near As relative humidity was above 80% and rainy day condition, there will be chances of disease and pest infestation mainly sheath blight, blight, blast, foot rot and swarming caterpillar, leaf folder, stem borer in rice. They may pause all agricultural activities based on rainfall forecast especially today to Thursday and resume it after.

Kharif Rice:

- The crop is at tillering stage. Maintain about 2-3 cm standing water during tillering stage. Problem of weeds may aggravate after transplanting, apply Bensulfuron methyl 0.6% + Pretilachlor 6% GR with sand mix at 1:1 ratio @ 4 kg/acre within 3-7 days after transplanting for controlling weeds.
- Intermittent dry and wet weather condition, is favorable for incidence of BLB and Sheath blight in early sown nursery seedlings. Go for foliar spray of Hexaconazole with Plantomycin @ 1.0 g/l to prevent the spread of the disease.
- If leaf blast is observed in nursery, spray Tebuconazole 50% + Trifloxystrobin 25% WG @ 0.4 g or Isoprothiolane 40EC @ 1.5 ml per litre of water. Repeat the spray at 7-10 days interval.
- In Gall midge affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre
- If swarming caterpillar will be found after the rain water recedes, apply Lambda cyhalothrin 5% EC @ 100 ml/acre.
- Temperature below 28°C and continuous rainfall lead to infestation of stem borer at tillering stage of rice, to control this apply Azadirachtin 0.15% EC @ 800 ml/acre or Cartap hydrochloride 4G @ 10kg/acre.

Maize: Maize crop is highly sensitive to waterlogged conditions during seedling stage. Hence, drain out excess water from the field as early as possible.

- After cessation of rains apply booster dose of fertilizer.
- Weeds may be controlled after rainfall.
- Stem fly in Maize infects the plants from sowing to 5 weeks after sowing, temperature between 25-30 deg. celsius and humidity above 60% favour pest incidence. In vegetative stage to control stem fly spray Methyl demeton @ 1 ml/lit or Dimethoate @ 2ml/lit or apply Carbofuran 3G granules @ 14 kg/acre.
- High humid conditions, temperature between 18-27 Deg. Celsius favours the occurrence of leaf blight in Maize. To control spray Azoxystrobin + Mancozeb 300 g/acre. Due to water logging condition stem rot may be found, then farmers should apply carbendazim @ 2g/ltr water.

VEGETABLES: Due to past rain, Chances of fruit rot infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary, spray second and third round at 15 days interval.

- If incidence of Damping off is noticed in Tomato, Brinjal, Chilli nurseries, drench nursery bed with Copper-Oxy-Chloride @ 3 g per litre of water.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever sucking pests incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. To prevent of soil borne diseases like seedling rot, stem rot and wilt diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenconazole @ 1ml/lit at 15 days interval to control Cercospora leaf spot.

ANIMAL HUSBANDRY: Keep animal shed and shelter house free from water logging by keeping it dry and free from weed. Fresh and immature leaves should not be fed more amount as they may cause diarrhea. If grasses were cut in morning or evening time, it can be sun dried for about 30 minute before feeding.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-429

Date- 22/08/2023

AGROMET ADVISORY BULLETIN FOR BARANGA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 7mm & district received average 10mm rainfall during the last week. The mean daily maximum and minimum temperature was **32.8°C (+1°C from normal)** and **25.8°C (+1.7°C from normal)**, respectively. Mean relative humidity of the district was around 84-90%.

Forecasted period: Today(8.30IST) to 27th August (8.30IST)2023

Weather advisory: *Moderate to isolated heavy rainfall is forecasted for tomorrow & light to moderate rainfall is forecasted in between other 4 days in the block.* The day temperature will be 31-33°C and night temperature will be 25-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 79-83% and 45-57% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Baranga (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	22/08/2023	23/08/2023	24/08/2023	25/08/2023	26/08/2023
Rainfall(mm)	16	34	22	7	6
T-Max(°C)	33	32	31	31	33
T-Min(°C)	26	26	26	25	25
Cloud Cover	8	8	8	8	8
RH Max (%)	83	80	79	81	81
RH Min (%)	51	52	57	52	45
Wind Speed (Kmph)	16	14	13	12	14
Wind Direction (deg)	204	225	252	257	279

Agromet Advisory

General advisory: Farmers should drain out excess water from vegetable and rice field near As relative humidity was above 80% and rainy day condition, there will be chances of disease and pest infestation mainly sheath blight, blight, blast, foot rot and swarming caterpillar, leaf folder, stem borer in rice. They may pause all agricultural activities based on rainfall forecast especially today to Thursday and resume it after.

Kharif Rice:

- The crop is at tillering stage. Maintain about 2-3 cm standing water during tillering stage. Problem of weeds may aggravate after transplanting, apply Bensulfuron methyl 0.6% + Pretilachlor 6% GR with sand mix at 1:1 ratio @ 4 kg/acre within 3-7 days after transplanting for controlling weeds.
- Intermittent dry and wet weather condition, is favorable for incidence of **BLB and Sheath blight** in early sown nursery seedlings. Go for foliar spray of Hexaconazole with Plantomycin @ 1.0 g/l to prevent the spread of the disease.
- If **leaf blast** is observed in nursery, spray Tebuconazole 50% + Trifloxystrobin 25% WG @ 0.4 g or Isoprothiolane 40EC @ 1.5 ml per litre of water. Repeat the spray at 7-10 days interval.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre
- If **swarming caterpillar** will be found after the rain water recedes, apply Lambda cyhalothrin 5% EC @ 100 ml/acre.
- Temperature below 28°C and continuous rainfall lead to infestation of **stem borer** at tillering stage of rice, to control this apply Azadirachtin 0.15% EC @ 800 ml/acre or Cartap hydrochloride 4G @ 10kg/acre.

Maize: Maize crop is highly sensitive to waterlogged conditions during seedling stage. Hence, drain out excess water from the field as early as possible.

- After cessation of rains apply booster dose of fertilizer.
- Weeds may be controlled after rainfall.
- Stem fly** in Maize infects the plants from sowing to 5 weeks after sowing, temperature between 25-30 deg. celsius and humidity above 60% favour pest incidence. In vegetative stage to control stem fly spray Methyl demeton @ 1 ml/lit or Dimethoate @ 2ml/lit or apply Carbofuran 3G granules @ 14 kg/acre.
- High humid conditions, temperature between 18-27 Deg. Celsius favours the occurrence of **leaf blight** in Maize. To control spray Azoxystrobin + Mancozeb 300 g /ac. Due to water logging condition stem rot may be found, then farmers should apply carbendazim @ 2g/ltr water.

VEGETABLES: Due to past rain, Chances of **fruit rot** infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary, spray second and third round at 15 days interval.

- If incidence of **Damping off** is noticed in Tomato, Brinjal, Chilli nurseries, drench nursery bed with Copper-Oxy-Chloride @ 3 g per liter of water.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever **sucking pests** incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. To prevent of soil borne diseases like **seedling rot, stem rot and wilt** diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenconazole @ 1ml/lit at 15 days interval to control **Cercospora leaf spot**.

ANIMAL HUSBANDRY: Keep animal shed and shelter house free from water logging by keeping it dry and free from weed. Fresh and immature leaves should not to be fed more amount as they may cause diarrhea. If grasses were cut in morning or evening time, it can be sun dried for about 30 minute before feeding.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA INDIA METEOROLOGICAL DEPARTMENT DISTRICT AGROMET UNIT KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

AGROMET ADVISORY BULLETIN FOR CUTTACK SADAR



Bulletin No.-429

Date- 22/08/2023

District Past week average weather condition and Block past week average Rainfall condition

The block received average 5mm & district received average 10mm rainfall during the last week. The mean daily maximum and minimum temperature was 32.8°C (+1°C from normal) and 25.8°C (+1.7°C from normal), respectively. Mean relative humidity of the district was around 84-90%.

Forecasted period: Today(8.30IST) to 27th August (8.30IST)2023

Weather advisory: Light to moderate rainfall is forecasted in between next 5 days in the block. The day temperature will be 30-33°C and night temperature will be 25-27°C for upcoming 5 days. Maximum and minimum relative humidity will be 80-83% and 48-59% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Cuttack sadar (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	22/08/2023	23/08/2023	24/08/2023	25/08/2023	26/08/2023
Rainfall(mm)	9	26	25	7	6
T-Max(°C)	33	32	30	31	32
T-Min(°C)	27	26	26	25	26
Cloud Cover(octa)	8	8	8	8	8
RH Max (%)	83	82	80	82	82
RH Min (%)	52	53	59	54	48
Wind Speed (Kmph)	14	13	13	12	13
Wind Direction (deg)	204	225	249	257	287

Agromet Advisory

General advisory: Farmers should drain out excess water from vegetable and rice field near As relative humidity was above 80% and rainy day condition, there will be chances of disease and pest infestation mainly sheath blight, blight, blast, foot rot and swarming caterpillar, leaf folder, stem borer in rice. They may pause all agricultural activities based on rainfall forecast especially today to Thursday and resume it after.

Kharif Rice:

- The crop is at tillering stage. Maintain about 2-3 cm standing water during tillering stage. Problem of weeds may aggravate after transplanting, apply Bensulfuron methyl 0.6% + Pretilachlor 6% GR with sand mix at 1:1 ratio @ 4 kg/acre within 3-7 days after transplanting for controlling weeds.
- Intermittent dry and wet weather condition, is favorable for incidence of BLB and Sheath blight in early sown nursery seedlings. Go for foliar spray of Hexaconazole with Plantomycin @ 1.0 g/l to prevent the spread of the disease.
- If leaf blast is observed in nursery, spray Tebuconazole 50% + Trifloxystrobin 25% WG @ 0.4 g or Isoprothiolane 40EC @ 1.5 ml per litre of water. Repeat the spray at 7-10 days interval.
- In Gall midge affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre
- If swarming caterpillar will be found after the rain water recedes, apply Lambda cyhalothrin 5% EC @ 100 ml/acre.
- Temperature below 28deg c and continuous rainfall lead to infestation of stem borer at tillering stage of rice, to control this apply Azadirachtin 0.15% EC @ 800 ml/acre or Cartap hydrochloride 4G @ 10kg/acre.

Sugarcane:

- Drain out excess water from the field as early as possible.
- After cessation of rains weeding and earthing up should be done in this time after that Top dressing of Urea @ 30 kg/acre for 30-60 days old sugarcane plant.
- For Red rot and wilt disease, spray the crop with Mancozeb 0.3% or carbendazim 0.1% or Azoxystrobin 18.2% SC / Difenconazole 11.45SC @ 1ml per liter water in dry days.
- For early shoot borer Spray Chlorpyrifos @ 400 ml/ac in 200 lts of water after cessation of rain.

VEGETABLES: Due to past rain, Chances of fruit rot infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary, spray second and third round at 15 days interval.

- If incidence of Damping off is noticed in Tomato, Brinjal, Chilli nurseries, drench nursery bed with Copper-Oxy-Chloride @ 3 g per liter of water.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever sucking pests incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. To prevent of soil borne diseases like seedling rot, stem rot and wilt diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenconazole @ 1ml/lit at 15 days interval to control Cercospora leaf spot.

ANIMAL HUSBANDRY: Keep animal shed and shelter house free from water logging by keeping it dry and free from weed. Fresh and immature leaves should not be fed more amount as they may cause diarrhea. If grasses were cut in morning or evening time, it can be sun dried for about 30 minute before feeding.

Dr Sujata Sathy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)



Bulletin No.-429
Date- 22/08/2023

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

AGROMET ADVISORY BULLETIN FOR KANTAPADA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 8mm & district received average 10mm rainfall during the last week. The mean daily maximum and minimum temperature was **32.8°C (+1°C from normal)** and **25.8°C (+1.7°C from normal)**, respectively. Mean relative humidity of the district was around 84-90%.

Forecasted period: Today(8.30IST) to 27th August (8.30IST)2023

Weather advisory: *Moderate to isolated heavy rainfall is forecasted for tomorrow & light to moderate rainfall is forecasted in between other 4 days in the block.* The day temperature will be 31-33°C and night temperature will be 25-27°C for upcoming 5 days. Maximum and minimum relative humidity will be 79-81% and 45-55% respectively. **There will be partly cloudy to overcast sky for next 5 days.**

Block	Kantapada (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	22/08/2023	23/08/2023	24/08/2023	25/08/2023	26/08/2023
Rainfall(mm)	9	31	27	7	9
T-Max(°C)	33	32	31	31	33
T-Min(°C)	27	26	26	25	26
Cloud Cover(octa)	8	8	8	8	8
RH Max (%)	81	79	79	81	80
RH Min (%)	52	52	55	52	45
Wind Speed (Kmph)	19	15	13	13	14
Wind Direction (deg)	203	225	249	248	259

Agromet Advisory

General advisory: *Farmers should drain out excess water from vegetable and rice field near As relative humidity was above 80% and rainy day condition, there will be chances of disease and pest infestation mainly sheath blight, blight, blast, foot rot and swarming caterpillar, leaf folder, stem borer in rice. They may pause all agricultural activities based on rainfall forecast especially today to Thursday and resume it after.*

Kharif Rice:

- The crop is at tillering stage. Maintain about 2-3 cm standing water during tillering stage..Problem of weeds may aggravate after transplanting, apply Bensulfuron methyl 0.6% + Pretilachlor 6% GR with sand mix at 1:1 ratio @ 4 kg/acre within 3-7 days after transplanting for controlling weeds.
- Intermittent dry and wet weather condition, is favorable for incidence of **BLB and Sheath blight** in early sown nursery seedlings. Go for foliar spray of Hexaconazole with Plantomycin@1.0 g/l to prevent the spread of the disease.
- If **leaf blast** is observed in nursery, spray Tebuconazole 50% + Trifloxystrobin 25% WG@ 0.4 g or Isoprothiolane 40EC @ 1.5 ml per litre of water. Repeat the spray at 7-10 days interval.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre
- If **swarming caterpillar** will be found after the rain water recedes, apply Lambda cyhalothrin 5% EC @ 100 ml/acre.
- Temperature below 28deg c and continuous rainfall lead to infestation of **stem borer** at tillering stage of rice, to control this apply Azadirachtin 0.15% EC @ 800 ml/acre or Cartap hydrochloride 4G @ 10kg/acre.

Sugarcane:

- Drain out excess water from the field as early as possible.
- After cessation of rains weeding and earthing up should be done in this time after that Top dressing of Urea @30 kg/acre for 30-60 days old sugarcane plant.
- For **Red rot and wilt disease**, spray the crop with Mancozeb 0.3% or carbendazim 0.1% or Azoxystrobin 18.2%SC/ Difenconazole 11.45SC @ 1ml per liter water in dry days.
- For early shoot borer Spray Chloropyrifos @ 400 ml/ac in 200 lts of water after cessation of rain.

VEGETABLES: Due to past rain, Chances of **fruit rot** infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary, spray second and third round at 15 days interval.

- If incidence of **Damping off** is noticed in Tomato, Brinjal, Chilli nurseries, drench nursery bed with Copper-Oxy-Chloride @ 3 g per liter of water.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever **sucking pests** incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. To prevent of soil borne diseases like **seedling rot, stem rot and wilt** diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenconazole @ 1ml/lit at 15 days interval to control **Cercospora leaf spot**.

ANIMAL HUSBANDRY: Keep animal shed and shelter house free from water logging by keeping it dry and free from weed. Fresh and immature leaves should not be fed more amount as they may cause diarrhea. If grasses were cut in morning or evening time, it can be sun dried for about 30 minute before feeding.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA INDIA METEOROLOGICAL DEPARTMENT DISTRICT AGROMET UNIT KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

AGROMET ADVISORY BULLETIN FOR MAHANGA



Bulletin No.-429

Date- 22/08/2023

District Past week average weather condition and Block past week average Rainfall condition

The block received average 6mm & district received average 10mm rainfall during the last week. The mean daily maximum and minimum temperature was **32.8°C (+1°C from normal)** and **25.8°C (+1.7°C from normal)**, respectively. Mean relative humidity of the district was around 84-90%.

Forecasted period: Today(8.30IST) to 27th August (8.30IST)2023

Weather advisory: ; Light to moderate rainfall is forecasted in between next 5 days in the block. The day temperature will be and it will be 30-33°C and night temperature will be 25-26°C for upcoming 5 days. Maximum and minimum relative humidity will 80-84% and 48-59% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Mahanga (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	22/08/2023	23/08/2023	24/08/2023	25/08/2023	26/08/2023
Rainfall(mm)	7	24	26	7	9
T-Max(°C)	33	33	31	30	32
T-Min(°C)	26	26	25	25	26
Cloud Cover(octal)	8	8	8	8	8
RH Max (%)	84	80	81	81	81
RH Min (%)	52	53	59	55	48
Wind Speed (Kmph)	16	14	16	14	15
Wind Direction (deg)	203	225	249	249	259

Agro-met Advisory

General advisory: Farmers should drain out excess water from vegetable and rice field near As relative humidity was above 80% and rainy day condition, there will be chances of disease and pest infestation mainly sheath blight, blight, blast, foot rot and swarming caterpillar, leaf folder, stem borer in rice. They may pause all agricultural activities based on rainfall forecast especially today to Thursday and resume it after.

Kharif Rice:

- The crop is at tillering stage. Maintain about 2-3 cm standing water during tillering stage. Problem of weeds may aggravate after transplanting, apply Bensulfuron methyl 0.6% + Pretilachlor 6% GR with sand mix at 1:1 ratio @ 4 kg/acre within 3-7 days after transplanting for controlling weeds.
- Intermittent dry and wet weather condition, is favorable for incidence of **BLB and Sheath blight** in early sown nursery seedlings. Go for foliar spray of Hexaconazole with Plantomycin @ 1.0 g/l to prevent the spread of the disease.
- If **leaf blast** is observed in nursery, spray Tebuconazole 50% + Trifloxystrobin 25% WG @ 0.4 g or Isoprothiolane 40EC @ 1.5 ml per litre of water. Repeat the spray at 7-10 days interval.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre
- If **swarming caterpillar** will be found after the rain water recedes, apply Lambda cyhalothrin 5% EC @ 100 ml/acre.
- Temperature below 28deg c and continuous rainfall lead to infestation of **stem borer** at tillering stage of rice, to control this apply Azadirachtin 0.15% EC @ 800 ml/acre or Cartap hydrochloride 4G @ 10kg/acre.

Sugarcane:

- Drain out excess water from the field as early as possible.
- After cessation of rains weeding and earthing up should be done in this time after that Top dressing of Urea @ 30 kg/acre for 30-60 days old sugarcane plant.
- For **Red rot and wilt disease**, spray the crop with Mancozeb 0.3% or carbendazim 0.1% or Azoxystrobin 18.2% SC / Difenoconazole 11.45SC @ 1ml per liter water in dry days.

For early shoot borer Spray Chloropyriphos @ 400 ml/ac in 200 lts of water after cessation of rain.

Jute: Recent rain has aggravated **weed problem** in early sown crop. To control grass weeds, apply Quisalofof-pethyl @ 2ml/litre of water at 2-3 leaf stage of weeds. After rain in vegetative stage to control **stem rot of jute**, spray copper oxychloride @ 3-4 gram/liter of water.

- To control **indigo caterpillar** apply Spinosad (45% S.C) 0.3 ml/liter of water or Emamectin benzoate (5% SC) 0.5 gram per liter of water.

VEGETABLES:

Due to past rain, Chances of **fruit rot** infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary, spray second and third round at 15 days interval.

- If incidence of **Damping off** is noticed in Tomato, Brinjal, Chilli nurseries, drench nursery bed with Copper-Oxy-Chloride @ 3 g per liter of water.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever **sucking pests** incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. To prevent of soil borne diseases like **seedling rot, stem rot and wilt** diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenoconazole @ 1ml/lit at 15 days interval to control **Cercospora leaf spot**.

Dr Sujata Sathy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA INDIA METEOROLOGICAL DEPARTMENT DISTRICT AGROMET UNIT KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-429

Date- 22/08/2023

AGROMET ADVISORY BULLETIN FOR NARASINGHPUR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 8mm & district received average 10mm rainfall during the last week. The mean daily maximum and minimum temperature was **32.8°C (+1°C from normal)** and **25.8°C (+1.7°C from normal)**, respectively. Mean relative humidity of the district was around 84-90%.

Forecasted period: Today(8.30IST) to 27th August (8.30IST)2023

Weather advisory: Moderate to isolated heavy rainfall is forecasted for tomorrow & light to moderate rainfall is forecasted in between other 4 days in the block. The day temperature will be 30-32°C and night temperature will be 24-25°C for upcoming 5 days. Maximum and minimum relative humidity will be 82-90% and 52-62% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Narasinghpur (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	22/08/2023	23/08/2023	24/08/2023	25/08/2023	26/08/2023
Rainfall(mm)	20	35	18	7	6
T-Max(°C)	32	31	30	30	32
T-Min(°C)	25	24	24	24	24
Cloud Cover(octa)	8	8	8	8	8
RH Max (%)	88	90	84	82	83
RH Min (%)	59	62	62	61	52
Wind Speed (Kmph)	7	8	9	9	9
Wind Direction (deg)	214	243	249	270	279

Agromet Advisory

General advisory: Farmers should drain out excess water from vegetable and rice field near As relative humidity was above 80% and rainy day condition, there will be chances of disease and pest infestation mainly sheath blight, blight, blast, foot rot and swarming caterpillar, leaf folder, stem borer in rice. They may pause all agricultural activities based on rainfall forecast especially today to Thursday and resume it after.

Kharif Rice:

- The crop is at tillering stage. Maintain about 2-3 cm standing water during tillering stage. Problem of weeds may aggravate after transplanting, apply Bensulfuron methyl 0.6% + Pretilachlor 6% GR with sand mix at 1:1 ratio @ 4 kg/acre within 3-7 days after transplanting for controlling weeds.
- Intermittent dry and wet weather condition, is favorable for incidence of **BLB and Sheath blight** in early sown nursery seedlings. Go for foliar spray of Hexaconazole with Plantomycin @ 1.0 g/l to prevent the spread of the disease.
- If **leaf blast** is observed in nursery, spray Tebuconazole 50% + Trifloxystrobin 25% WG @ 0.4 g or Isoprothiolane 40EC @ 1.5 ml per litre of water. Repeat the spray at 7-10 days interval.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre
- If **swarming caterpillar** will be found after the rain water recedes, apply Lambda cyhalothrin 5% EC @ 100 ml/acre.
- Temperature below 28deg c and continuous rainfall lead to infestation of **stem borer** at tillering stage of rice, to control this apply Azadirachtin 0.15% EC @ 800 ml/acre or Cartap hydrochloride 4G @ 10kg/acre.

Maize: Maize crop is highly sensitive to waterlogged conditions during seedling stage. Hence, drain out excess water from the field as early as possible.

- After cessation of rains apply booster dose of fertilizer.
- Weeds may be controlled after rainfall.
- Stem fly** in Maize infects the plants from sowing to 5 weeks after sowing, temperature between 25-30 deg. celsius and humidity above 60% favour pest incidence. In vegetative stage to control stem fly spray Methyl demeton @ 1 ml/lit or Dimethoate @ 2ml/lit or apply Carbofuran 3G granules @ 14 kg/acre.
- High humid conditions, temperature between 18-27 Deg. Celsius favours the occurrence of **leaf blight** in Maize. To control spray Azoxystrobin + Mancozeb 300 g/ac. Due to water logging condition stem rot may be found, then farmers should apply carbendazim @ 2g/ltr water.

VEGETABLES: Due to past rain, Chances of **fruit rot** infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary, spray second and third round at 15 days interval.

- If incidence of **Damping off** is noticed in Tomato, Brinjal, Chilli nurseries, drench nursery bed with Copper-Oxy-Chloride @ 3 g per liter of water.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever **sucking pests** incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. To prevent of soil borne diseases like **seedling rot, stem rot and wilt** diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenconazole @ 1ml/lit at 15 days interval to control **Cercospora leaf spot**.

ANIMAL HUSBANDRY: Keep animal shed and shelter house free from water logging by keeping it dry and free from weed. Fresh and immature leaves should not to be fed more amount as they may cause diarrhea. If grasses were cut in morning or evening time, it can be sun dried for about 30 minute before feeding.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)



Bulletin No.-429
Date- 22/08/2023

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

AGROMET ADVISORY BULLETIN FOR NIALI

District Past week average weather condition and Block past week average Rainfall condition

The block received average 13mm & district received average 10mm rainfall during the last week. The mean daily maximum and minimum temperature was **32.8°C (+1°C from normal)** and **25.8°C (+1.7°C from normal)**, respectively. Mean relative humidity of the district was around 84-90%.

Forecasted period: Today(8.30IST) to 27th August (8.30IST)2023

Weather advisory: *Moderate to isolated heavy rainfall is forecasted for tomorrow & light to moderate rainfall is forecasted in between other 4 days in the block.* The day temperature will be 31-33°C and night temperature will be 25-27°C for upcoming 5 days. Maximum and minimum relative humidity will be 78-80% and 46-54% respectively. **There will be partly cloudy to overcast sky for next 5 days.**

Block	Niali (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	22/08/2023	23/08/2023	24/08/2023	25/08/2023	26/08/2023
Rainfall(mm)	8	32	26	7	6
T-Max(°C)	33	33	31	31	33
T-Min(°C)	27	26	26	25	26
Cloud Cover(octa)	8	8	8	8	8
RH Max (%)	80	79	78	80	80
RH Min (%)	53	53	54	52	46
Wind Speed(Kmph)	21	17	14	14	15
Wind Direction(deg)	203	212	248	248	248

Agromet Advisory

General advisory: Farmers should drain out excess water from vegetable and rice field near As relative humidity was above 80% and rainy day condition, there will be chances of disease and pest infestation mainly sheath blight, blight, blast, foot rot and swarming caterpillar, leaf folder, stem borer in rice. They may pause all agricultural activities based on rainfall forecast especially today to Thursday and resume it after.

Kharif Rice:

- The crop is at tillering stage. Maintain about 2-3 cm standing water during tillering stage. Problem of weeds may aggravate after transplanting, apply Bensulfuron methyl 0.6% + Pretilachlor 6% GR with sand mix at 1:1 ratio @ 4 kg/acre within 3-7 days after transplanting for controlling weeds.
- Intermittent dry and wet weather condition, is favorable for incidence of **BLB and Sheath blight** in early sown nursery seedlings. Go for foliar spray of Hexaconazole with Plantomycin @ 1.0 g/l to prevent the spread of the disease.
- If **leaf blast** is observed in nursery, spray Tebuconazole 50% + Trifloxystrobin 25% WG @ 0.4 g or Isoprothiolane 40EC @ 1.5 ml per litre of water. Repeat the spray at 7-10 days interval.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre
- If **swarming caterpillar** will be found after the rain water recedes, apply Lambda cyhalothrin 5% EC @ 100 ml/acre.
- Temperature below 28deg c and continuous rainfall lead to infestation of **stem borer** at tillering stage of rice, to control this apply Azadirachtin 0.15% EC @ 800 ml/acre or Cartap hydrochloride 4G @ 10kg/acre.

Sugarcane:

- Drain out excess water from the field as early as possible.
- After cessation of rains weeding and earthing up should be done in this time after that Top dressing of Urea @ 30 kg/acre for 30-60 days old sugarcane plant.
- For **Red rot and wilt disease**, spray the crop with Mancozeb 0.3% or carbendazim 0.1% or Azoxystrobin 18.2% SC/ Difenconazole 11.45SC @ 1ml per liter water in dry days.
- For early shoot borer Spray Chloropyrifos @ 400 ml/ac in 200 lts of water after cessation of rain.

VEGETABLES: Due to past rain, Chances of **fruit rot** infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary, spray second and third round at 15 days interval.

- If incidence of **Damping off** is noticed in Tomato, Brinjal, Chilli nurseries, drench nursery bed with Copper-Oxy-Chloride @ 3 g per liter of water.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever **sucking pests** incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. To prevent of soil borne diseases like **seedling rot, stem rot and wilt** diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenconazole @ 1ml/lit at 15 days interval to control **Cercospora leaf spot**.

ANIMAL HUSBANDRY: Keep animal shed and shelter house free from water logging by keeping it dry and free from weed. Fresh and immature leaves should not to be fed more amount as they may cause diarrhea. If grasses were cut in morning or evening time, it can be sun dried for about 30 minute before feeding.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA INDIA METEOROLOGICAL DEPARTMENT DISTRICT AGROMET UNIT KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

AGROMET ADVISORY BULLETIN FOR NISCHINTAKOILI

District Past week average weather condition and Block past week average Rainfall condition

The block received average 5mm & district received average 10mm rainfall during the last week. The mean daily maximum and minimum temperature was 32.8°C (+1°C from normal) and 25.8°C (+1.7°C from normal), respectively. Mean relative humidity of the district was around 84-90%.

Forecasted period: Today(8.30IST) to 27th August (8.30IST)2023

Weather advisory : Light to moderate rainfall is forecasted in between next 5 days in the block. The day temperature will be 30-33°C and night temperature will be 25-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 79-83% and 47-57% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Nischintakoili (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	22/08/2023	23/08/2023	24/08/2023	25/08/2023	26/08/2023
Rainfall(mm)	9	25	26	7	10
T-Max(°C)	33	33	31	30	32
T-Min(°C)	26	26	26	25	26
Cloud Cover(octa)	8	8	8	8	8
RH Max (%)	83	79	81	82	81
RH Min (%)	52	53	57	55	47
Wind Speed (Kmph)	17	15	15	14	15
Wind Direction (deg)	203	225	249	249	259

Agromet Advisory

General advisory: Farmers should drain out excess water from vegetable and rice field near As relative humidity was above 80% and rainy day condition, there will be chances of disease and pest infestation mainly sheath blight, blight, blast, foot rot and swarming caterpillar, leaf folder, stem borer in rice. They may pause all agricultural activities based on rainfall forecast especially today to Thursday and resume it after.

Kharif Rice:

- The crop is at tillering stage. Maintain about 2-3 cm standing water during tillering stage. Problem of weeds may aggravate after transplanting, apply Bensulfuron methyl 0.6% + Pretilachlor 6% GR with sand mix at 1:1 ratio @ 4 kg/acre within 3-7 days after transplanting for controlling weeds.
- Intermittent dry and wet weather condition, is favorable for incidence of **BLB and Sheath blight** in early sown nursery seedlings. Go for foliar spray of Hexaconazole with Plantomycin @ 1.0 g/l to prevent the spread of the disease.
- If **leaf blast** is observed in nursery, spray Tebuconazole 50% + Trifloxystrobin 25% WG @ 0.4 g or Isoprothiolane 40EC @ 1.5 ml per litre of water. Repeat the spray at 7-10 days interval.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre
- If **swarming caterpillar** will be found after the rain water recedes, apply Lambda cyhalothrin 5% EC @ 100 ml/acre.
- Temperature below 28°C and continuous rainfall lead to infestation of **stem borer** at tillering stage of rice, to control this apply Azadirachtin 0.15% EC @ 800 ml/acre or Cartap hydrochloride 4G @ 10kg/acre.

Sugarcane:

- Drain out excess water from the field as early as possible.
- After cessation of rains weeding and earthing up should be done in this time after that Top dressing of Urea @ 30 kg/acre for 30-60 days old sugarcane plant.
- For **Red rot and wilt disease**, spray the crop with Mancozeb 0.3% or carbendazim 0.1% or Azoxystrobin 18.2% SC / Difenoconazole 11.45SC @ 1ml per liter water in dry days.
- For early shoot borer Spray Chloropyrifos @ 400 ml/ac in 200 lts of water after cessation of rain.

VEGETABLES: Due to past rain, Chances of **fruit rot** infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary, spray second and third round at 15 days interval.

- If incidence of **Damping off** is noticed in Tomato, Brinjal, Chilli nurseries, drench nursery bed with Copper-Oxy-Chloride @ 3 g per liter of water.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever **sucking pests** incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. To prevent of soil borne diseases like **seedling rot, stem rot and wilt** diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenoconazole @ 1ml/lit at 15 days interval to control **Cercospora leaf spot**.

ANIMAL HUSBANDRY: Keep animal shed and shelter house free from water logging by keeping it dry and free from weed. Fresh and immature leaves should not be fed more amount as they may cause diarrhea. If grasses were cut in morning or evening time, it can be sun dried for about 30 minute before feeding.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-429
Date- 22/08/2023

AGROMET ADVISORY BULLETIN FOR SALEPUR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 7mm & district received average 10mm rainfall during the last week. The mean daily maximum and minimum temperature was **32.8°C (+1°C from normal)** and **25.8°C (+1.7°C from normal)**, respectively. Mean relative humidity of the district was around 84-90%.

Forecasted period: Today(8.30IST) to 27th August (8.30IST)2023

Weather advisory: ; *Light to moderate rainfall is forecasted in between next 5 days in the block.* The day temperature will be 30-33°C and night temperature will be 25-27°C for upcoming 5 days. Maximum and minimum relative humidity will be 79-83% and 47-58% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Salepur (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	22/08/2023	23/08/2023	24/08/2023	25/08/2023	26/08/2023
Rainfall(mm)	8	24	23	7	9
T-Max(°C)	33	33	31	30	32
T-Min(°C)	27	26	26	25	26
Cloud Cover(octa)	8	8	8	8	8
RH Max (%)	83	79	81	82	81
RH Min (%)	52	53	58	55	47
Wind Speed (Kmph)	16	14	15	14	15
Wind Direction (deg)	203	225	249	249	259

Agromet Advisory

General advisory: Farmers should drain out excess water from vegetable and rice field near As relative humidity was above 80% and rainy day condition, there will be chances of disease and pest infestation mainly sheath blight, blight, blast, foot rot and swarming caterpillar, leaf folder, stem borer in rice. They may pause all agricultural activities based on rainfall forecast especially today to Thursday and resume it after.

Kharif Rice:

- The crop is at tillering stage. Maintain about 2-3 cm standing water during tillering stage.. Problem of weeds may aggravate after transplanting, apply Bensulfuron methyl 0.6% + Pretilachlor 6% GR with sand mix at 1:1 ratio @ 4 kg/acre within 3-7 days after transplanting for controlling weeds.
- Intermittent dry and wet weather condition, is favorable for incidence of **BLB and Sheath blight** in early sown nursery seedlings. Go for foliar spray of Hexaconazole with Plantomycin @ 1.0 g/l to prevent the spread of the disease.
- If **leaf blast** is observed in nursery, spray Tebuconazole 50% + Trifloxystrobin 25% WG @ 0.4 g or Isoprothiolane 40EC @ 1.5 ml per litre of water. Repeat the spray at 7-10 days interval.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre
- If **swarming caterpillar** will be found after the rain water recedes, apply Lambda cyhalothrin 5% EC @ 100 ml/acre.
- Temperature below 28deg c and continuous rainfall lead to infestation of **stem borer** at tillering stage of rice, to control this apply Azadirachtin 0.15% EC @ 800 ml/acre or Cartap hydrochloride 4G @ 10kg/acre.

Sugarcane:

- Drain out excess water from the field as early as possible.
- After cessation of rains weeding and earthing up should be done in this time after that Top dressing of Urea @ 30 kg/acre for 30-60 days old sugarcane plant.
- For **Red rot and wilt disease**, spray the crop with Mancozeb 0.3% or carbendazim 0.1% or Azoxystrobin 18.2% SC/ Difenconazole 11.45SC @ 1ml per liter water in dry days.

For early shoot borer Spray Chloropyrifos @ 400 ml/ac in 200 lts of water after cessation of rain.

Jute: Recent rain has aggravated **weed problem** in early sown crop. To control grass weeds, apply Quisalofof-pethyl @ 2ml/litre of water at 2-3 leaf stage of weeds. After rain in vegetative stage to control **stem rot of jute**, spray copper oxychloride @ 3-4 gram/liter of water.

- To control **indigo caterpillar** apply Spinosad (45% S.C) 0.3 ml/liter of water or Emamectin benzoate (5% SC) 0.5 gram per liter of water.

VEGETABLES:

Due to past rain, Chances of **fruit rot** infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary, spray second and third round at 15 days interval.

- If incidence of **Damping off** is noticed in Tomato, Brinjal, Chilli nurseries, drench nursery bed with Copper-Oxy-Chloride @ 3 g per liter of water.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever **sucking pests** incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. To prevent of soil borne diseases like **seedling rot, stem rot and wilt** diseases apply 3g of Copper oxychloride per litre of water around the main stem.

Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenconazole @ 1ml/lit at 15 days interval to control **Cercospora leaf spot**.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

AGROMET ADVISORY BULLETIN FOR TANGI-CHOUDWAR



Bulletin No.-429

Date- 22/08/2023

District Past week average weather condition and Block past week average Rainfall condition

The block received average 11mm & district received average 10mm rainfall during the last week. The mean daily maximum and minimum temperature was **32.8°C (+1°C from normal)** and **25.8°C (+1.7°C from normal)**, respectively. Mean relative humidity of the district was around 84-90%.

Forecasted period: Today(8.30IST) to 27th August (8.30IST)2023

Weather advisory: : *Light to moderate rainfall is forecasted in between next 5 days in the block.* The day temperature will be 30-33°C and night temperature will be 25-27°C for upcoming 5 days. Maximum and minimum relative humidity will be 79-81% and 50-61% respectively.

There will be partly cloudy to overcast sky for next 5 days.

Block	Tangi-Choudwar (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	22/08/2023	23/08/2023	24/08/2023	25/08/2023	26/08/2023
Rainfall(mm)	8	25	24	7	6
T-Max(°C)	33	33	31	30	32
T-Min(°C)	27	26	26	25	26
Cloud Cover(octa)	8	8	8	8	8
RH Max (%)	80	80	80	81	79
RH Min (%)	61	60	58	57	50
Wind Speed (Kmph)	23	21	16	17	18
Wind Direction (deg)	203	212	248	249	251

Agromet Advisory

General advisory: Farmers should drain out excess water from vegetable and rice field near As relative humidity was above 80% and rainy day condition, there will be chances of disease and pest infestation mainly sheath blight, blight, blast, foot rot and swarming caterpillar, leaf folder, stem borer in rice. They may pause all agricultural activities based on rainfall forecast especially today to Thursday and resume it after.

Kharif Rice:

- The crop is at tillering stage. Maintain about 2-3 cm standing water during tillering stage. Problem of weeds may aggravate after transplanting, apply Bensulfuron methyl 0.6% + Pretilachlor 6% GR with sand mix at 1:1 ratio @ 4 kg/acre within 3-7 days after transplanting for controlling weeds.
- Intermittent dry and wet weather condition, is favorable for incidence of **BLB and Sheath blight** in early sown nursery seedlings. Go for foliar spray of Hexaconazole with Plantomycin @ 1.0 g/l to prevent the spread of the disease.
- If **leaf blast** is observed in nursery, spray Tebuconazole 50% + Trifloxystrobin 25% WG @ 0.4 g or Isoprothiolane 40EC @ 1.5 ml per litre of water. Repeat the spray at 7-10 days interval.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre
- If **swarming caterpillar** will be found after the rain water recedes, apply Lambda cyhalothrin 5% EC @ 100 ml/acre.
- Temperature below 28deg c and continuous rainfall lead to infestation of **stem borer** at tillering stage of rice, to control this apply Azadirachtin 0.15% EC @ 800 ml/acre or Cartap hydrochloride 4G @ 10kg/acre.

Maize: Maize crop is highly sensitive to waterlogged conditions during seedling stage. Hence, drain out excess water from the field as early as possible.

- After cessation of rains apply booster dose of fertilizer.
- Weeds may be controlled after rainfall.
- Stem fly** in Maize infects the plants from sowing to 5 weeks after sowing, temperature between 25-30 deg. celsius and humidity above 60% favour pest incidence. In vegetative stage to control stem fly spray Methyl demeton @ 1 ml/lit or Dimethoate @ 2ml/lit or apply Carbofuran 3G granules @ 14 kg/acre.
- High humid conditions, temperature between 18-27 Deg. Celsius favours the occurrence of **leaf blight** in Maize. To control spray Azoxystrobin + Mancozeb 300 g/ac. Due to water logging condition stem rot may be found, then farmers should apply carbendazim @ 2g/ltr water.

VEGETABLES: Due to past rain, Chances of **fruit rot** infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary, spray second and third round at 15 days interval.

- If incidence of **Damping off** is noticed in Tomato, Brinjal, Chilli nurseries, drench nursery bed with Copper-Oxy-Chloride @ 3 g per liter of water.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever **sucking pests** incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. To prevent of soil borne diseases like **seedling rot, stem rot and wilt** diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenconazole @ 1ml/lit at 15 days interval to control **Cercospora leaf spot**.

ANIMAL HUSBANDRY: Keep animal shed and shelter house free from water logging by keeping it dry and free from weed. Fresh and immature leaves should not be fed more amount as they may cause diarrhea. If grasses were cut in morning or evening time, it can be sun dried for about 30 minute before feeding.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-429

Date- 22/08/2023

AGROMET ADVISORY BULLETIN FOR TIGIRIA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 10mm & district received average 10mm rainfall during the last week. The mean daily maximum and minimum temperature was **32.8°C (+1°C from normal)** and **25.8°C (+1.7°C from normal)**, respectively. Mean relative humidity of the district was around 84-90%.

Forecasted period: Today(8.30IST) to 27th August (8.30IST)2023

Weather advisory: ; *Light to moderate rainfall is forecasted in between next 5 days in the block.* The day temperature will be 30-33°C and night temperature will be 24-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 83-87% and 51-65% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Tigiria (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	22/08/2023	23/08/2023	24/08/2023	25/08/2023	26/08/2023
Rainfall(mm)	16	27	20	7	6
T-Max(°C)	33	32	30	31	32
T-Min(°C)	26	25	25	24	25
Cloud Cover(octa)	8	8	8	8	8
RH Max (%)	85	87	83	84	84
RH Min (%)	55	58	65	58	51
Wind Speed (Kmph)	11	11	11	10	11
Wind Direction (deg)	204	243	249	270	287

Agro met Advisory

General advisory: Farmers should drain out excess water from vegetable and rice field near As relative humidity was above 80% and rainy day condition, there will be chances of disease and pest infestation mainly sheath blight, blight, blast, foot rot and swarming caterpillar, leaf folder, stem borer in rice. They may pause all agricultural activities based on rainfall forecast especially today to Thursday and resume it after.

Kharif Rice:

- The crop is at tillering stage. Maintain about 2-3 cm standing water during tillering stage. Problem of weeds may aggravate after transplanting, apply Bensulfuron methyl 0.6% + Pretilachlor 6% GR with sand mix at 1:1 ratio @ 4 kg/acre within 3-7 days after transplanting for controlling weeds.
- Intermittent dry and wet weather condition, is favorable for incidence of **BLB and Sheath blight** in early sown nursery seedlings. Go for foliar spray of Hexaconazole with Plantomycin @ 1.0 g/l to prevent the spread of the disease.
- If **leaf blast** is observed in nursery, spray Tebuconazole 50% + Trifloxystrobin 25% WG @ 0.4 g or Isoprothiolane 40EC @ 1.5 ml per litre of water. Repeat the spray at 7-10 days interval.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre
- If **swarming caterpillar** will be found after the rain water recedes, apply Lambda cyhalothrin 5% EC @ 100 ml/acre.
- Temperature below 28°C and continuous rainfall lead to infestation of **stem borer** at tillering stage of rice, to control this apply Azadirachtin 0.15% EC @ 800 ml/acre or Cartap hydrochloride 4G @ 10kg/acre.

Maize: Maize crop is highly sensitive to waterlogged conditions during seedling stage. Hence, drain out excess water from the field as early as possible.

- After cessation of rains apply booster dose of fertilizer.
- Weeds may be controlled after rainfall.
- Stem fly** in Maize infects the plants from sowing to 5 weeks after sowing, temperature between 25-30 deg. celsius and humidity above 60% favour pest incidence. In vegetative stage to control stem fly spray Methyl demeton @ 1 ml/lit or Dimethoate @ 2ml/lit or apply Carbofuran 3G granules @ 14 kg/acre.
- High humid conditions, temperature between 18-27 Deg. Celsius favours the occurrence of **leaf blight** in Maize. To control spray Azoxystrobin + Mancozeb 300 g /ac. Due to water logging condition stem rot may be found, then farmers should apply carbendazim @ 2g/ltr water.
- VEGETABLES:** Due to past rain, Chances of **fruit rot** infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary, spray second and third round at 15 days interval.
- If incidence of **Damping off** is noticed in Tomato, Brinjal, Chilli nurseries, drench nursery bed with Copper-Oxy-Chloride @ 3 g per liter of water.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever **sucking pests** incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. To prevent of soil borne diseases like **seedling rot, stem rot and wilt** diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenconazole @ 1ml/lit at 15 days interval to control **Cercospora leaf spot**.

ANIMAL HUSBANDRY: Keep animal shed and shelter house free from water logging by keeping it dry and free from weed. Fresh and immature leaves should not be fed more amount as they may cause diarrhea. If grasses were cut in morning or evening time, it can be sun dried for about 30 minute before feeding.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-429
 Date- 22/08/2023

				
MEGHDOOT APP	DAMINI APP	FACEBOOK (Damu Cuttack)	BLOCK SPECIFIC WHAT'S APP GROUPS	@DAMUCUTTACK



- Kindly install and use **“Meghdoot mobile App”** for Cuttack district weather forecast, warnings and weather based agromet advisory services.

<https://play.google.com/store/apps/details?id=com.aas.meghdoot>

- Kindly install and use **“Damini mobile App”** for location specific for lightening forecast and warnings.

<https://play.google.com/store/apps/details?id=com.lightening.live.damini>

- Follow the **block specific whats app groups** and join in your block wise groups for getting weather based agromet advisory services twice in a week (Tuesday and Friday) and other weather warning based farm management advisories.
- Follow the **Official Website links** to get the weather based agromet advisory service bulletins (**English & Odia**) twice in a week (Tuesday and Friday).

1	ICAR-NRRI, Cuttack, Odisha	https://icar-nrri.in/agromate/	
2	Agromet division, IMD, Pune	https://imdagrmet.gov.in/imdproject/AGIndex.php	

Dr Sujata Sethy
 (Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
 SMS (Agrometeorology)
 Contact No-9861445080/7008453427