




Cuttack District Block Wise Agro-met Advisory Service Bulletin

*Tuesday, 12th September, 2023
(For the period 12th to 16th September,
2023)*



*Issued by
District Agro-met Unit,
Krishi Vigyan Kendra Cuttack*






GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-435
Date- 12/09/2023

STANDARD PRECIPITATION INDEX AND NDVI MAP



LEGEND (Medium range block level rainfall forecast)		
	No Rainfall	00 mm
	Very light rainfall	01-05 mm
	Light rainfall	06-10 mm
	Light to moderate rainfall	11-30 mm
	Moderate to heavy rainfall	31-65 mm
	Heavy to very heavy rainfall	66-115 mm
	Extremely heavy rainfall	>116 mm

LEGEND (Past week temperature)		
Color code	Temperature	Departure
	Below Normal	< 2°C from normal
	Normal	-2°C to +2°C from normal
	Above Normal	>2°C from normal

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA INDIA METEOROLOGICAL DEPARTMENT DISTRICT AGROMET UNIT KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)



Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com







Bulletin No.-435

Date- 12/09/2023

Past week (05-11 September, 2023) cumulative rainfall(mm) & its departure% from normal and (06-12 September) average maximum and minimum temperature for Cuttack district



Medium range (Today (8.30IST) to Sunday (8.30IST)) Rainfall (mm) forecast for Cuttack district



Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-435
 Date- 12/09/2023

Met Sub division	Extended range outlook for subsequent 1 week (6 th to 12 th days) (17-23 September 2023) after 5 days				
Odisha	<i>Parameter</i>	<i>Actual</i>	<i>Normal</i>	<i>Departure (%)</i>	<i>Category</i>
	Rainfall (mm)	13.16	11.26	16.82	Normal
	<i>Parameter</i>	<i>Actual</i>	<i>Normal</i>	<i>Anomaly</i>	<i>Category</i>
	Maximum temperature (°C)	31.20	31.84	-0.64	Normal
	Minimum temperature (°C)	24.45	24.01	0.44	Normal

***Agromet Advisory based upon subsequent week extended range forecast:-** There will be **normal rainfall is forecasted in the subsequent week (17-23 September, 2023) after 5 days in Odisha condition** and also there will be light to moderate with isolated heavy to very heavy rainfall is forecasted in between upcoming 5 days, so there may be water logging condition in rice and other crop and vegetable field. Farmers may go for drainage of excess water from rice, maize, pulses, kharif groundnut and vegetable field in upcoming 5 days/in the subsequent week. Due to likely normal rainfall activity in the subsequent week they may resume sowing of vegetables in the subsequent week. **Farmers should pause fertilizer/pesticide application, irrigation activities in the subsequent week.**



Dr Sujata Sethy
 (Nodal officer)



Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area



Debasish Jena
 SMS (Agrometeorology)
 Contact No-9861445080/7008453427












GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-435
Date- 12/09/2023

Weather (Past and Forecasted) based Disease and pest in Major Crops and vegetables

Sl No	Crop	Stage and condition of the crop	Anticipated Disease and pest	
1	Rice	Tillering to panicle initiation		Sheath blight
				Blast
				Stem/Foot rot
				Brown plant hopper
				Stem borer
				Gall midge
2	Sugarcane	6-7 month old		Red rot
				Rust
3	Maize	Tasseling to silking stage		Fall army worm

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area













Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-435
Date- 12/09/2023

				Banded leaf blight
				Downy mildew
4	Groundnut	Flowering to pegging/Harvesting		Collar rot
				Alternaria leaf spot
				Rust
4	Brinjal, tomato, chilli	seedling		Seedling rot
				Stem rot
5	Tomato and brinjal	Fruting		Bacterial wilt
6	Pointed gourd			Fruit rot
7	Pumpkin(Cucurbits)	Flowering to fruting		Stem and root rot

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-435

Date- 12/09/2023

District Past week average weather condition and Block past week average Rainfall condition

The block received average 12mm& district received average 12mm rainfall during the last week. The mean daily maximum and minimum temperature was **32°C (-0.09°C from normal)** and **26°C (+1.6°C from normal)**, respectively. Mean relative humidity of the district was around 88-92%.

Forecasted period: Today(8.30IST) to 17th September (8.30IST)2023

Weather advisory::Light to moderate with isolated heavy to very heavy rainfall is forecasted in between upcoming 5 days in the block. The day temperature will be **30-33°C** and night temperature will be **25-26°C** for upcoming 5 days. Maximum and minimum relative humidity will be 87-91% and 59-73% respectively. **There will be partly cloudy to overcast sky for next 5days.**

Block	Athagad(VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	12/09/2023	13/09/2023	14/09/2023	15/09/2023	16/09/2023
Rainfall (mm)	26	66	25	8	6
T-Max (°C)	33	31	31	31	30
T-Min (°C)	26	25	25	26	25
Cloud Cover (octa)	8	8	8	8	8
RH Max (%)	88	90	90	87	91
RH Min (%)	59	63	65	67	73
Wind Speed (kmph)	7	7	7	5	6
Wind Direction (deg)	333	203	198	18	203

Agromet Advisory

General advisory: Keep the drainage channels open to drain out excess rain water from crop fields. They should stop irrigation, pesticide and fertilizer application activities on rainy days. They should harvest matured vegetables and fruits before the rainfall activities & cover young seedlings with ploythine cover to protect them from heavy/intense shower.

Kharif Rice

Drain out excess water from the field. Maintain about 5-10cm standing water during panicle initiation stage.

Weather based Disease and pest of rice

- **Forecasted cloudy and humid weather** and the 20-25°C minimum temperature favors incidence of **stem borer and blast disease** in rice if the infestation will happen, spray Chlorantraniliprole 18.5% SC @ 60 ml/acre in 200 litres of water or Flubendiamide 20 WG 50g/acre in 200 litres of water to control **stem borer** and to control **blast** disease spray Tebuconazole 50%+ Trifloxystrobin 25% @ 0.4g/litre or, Carbendazim 50 WP @ 2g/litre of water.
- If **Brown Plant Hopper (BPH) infestation** will observe in rice, go for alternate wetting and drying. Whenever the number of hoppers/hill reaches 5-10, spray Pymetrozine 50% WG @ 120 g or Buprofezin 25% SC @ 330 ml or Imidacloprid 17.8% SL @ 50 ml per acre in the base of the plant. Before spraying remove water from the field.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre or Chlorpyrifos 20EC @ 500ml/acre to manage the pest.
- Due to likely water logging situation in Cuttack district as well as in the blocks **sheath blight** will be found in early transplanted rice then apply carbendazim 2g/ltr or propiconazole 1ml/ltr of water. In case of **Foot rot**, spray the basal portion of the crop with 0.4% mancozeb or 0.2% carbendazim after drying the weather.

Maize:

- **Provide drainage facility.** Rainfall after a dry spell may aggravate infestation of fall army worm in maize, to control fall armyworm in maize erect 4 pheromone traps /acre and spray 5 ml of 1500ppm neem oil /lit water. If more severe incidence observed spray Emamectin benzoate @ 0.5 g / lit of water. High humidity (>90%) and decrease in temperature (21-33°C) and continuous rain in the forecasted days may aggravate **downy mildew disease** in Maize. High humid conditions, temperature between 18-27 Deg. Celsius favors the occurrence of **leaf blight** in Maize. To control both disease spray Azoxystrobin + Mancozeb @ 300 g /ac.

Rabi-Groundnut:

- If groundnut is at harvesting stage then go for harvesting before rainy days. Anticipated humid, rainy weather favors infestation of **collar rot, alternaria leaf spot and rust** in groundnut. To control farmers may spray mancozeb/copper oxychloride 2g/liter water.

VEGETABLES:

- **Drain out excess water from the field.** Due to rain and high humidity, infestation of **blight and sucking pest** in vegetables likely to increase; farmers are advised to regular monitor the field. If observe, spray Mancozeb 75% WP @ 2g/l of water for blight and Imidacloprid 17.8% SL @ 0.25 ml/l of water for sucking pest.
- Due to rain and high humid condition during last week, **wilting** is likely to appear in young plants. Drain out water from the field. Spray mixing Streptocycline@20g with Carboxin+Thiram @400g or Metalaxyl 8%+Mancozeb 64% WP @400g per acre in 200 liters of water. Apply additional dose of potassic fertilizer to the crop.
- **Downy Mildew, vine rot, leaf spot and blight disease** in cucurbit vegetables, spray Carbendazim 12 % + Mancozeb 63 % WP @ 400 g/acre or Chlorothalonil @ 400 g/acre. Due to forecast of rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- **Brinjal, tomato and chilli** seedlings should be planted in the main field. To prevent of soil borne diseases like **seedling rot, stem rot and wilt** diseases apply 3g of Copper oxychloride per litre of water around the main stem.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA INDIA METEOROLOGICAL DEPARTMENT DISTRICT AGROMET UNIT KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-435

Date- 12/09/2023

AGROMET ADVISORY BULLETIN FOR BADAMBA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 16mm& district received average 12mm rainfall during the last week. The mean daily maximum and minimum temperature was 32°C (-0.09°C from normal) and 26°C (+1.6°C from normal), respectively. Mean relative humidity of the district was around 88-92%.

Forecasted period: Today(8.30IST) to 17th September (8.30IST)2023

Weather advisory: Light to moderate with isolated heavy to very heavy rainfall is forecasted in between upcoming 5 days in the block. The day temperature will be 30-33°C and night temperature will be 24-25°C for upcoming 5 days. Maximum and minimum relative humidity will be 87-91% and 60-74% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Badamba (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	12/09/2023	13/09/2023	14/09/2023	15/09/2023	16/09/2023
Rainfall(mm)	19	70	35	15	9
T-Max(°C)	33	31	30	31	30
T-Min(°C)	25	24	24	25	25
Cloud Cover(octa)	8	8	8	8	8
RH Max (%)	87	91	91	88	91
RH Min (%)	60	65	68	68	74
Wind Speed (Kmph)	7	5	6	4	4
Wind Direction (deg)	255	293	207	342	180

Agromet Advisory

General advisory: Keep the drainage channels open to drain out excess rain water from crop fields. They should stop irrigation, pesticide and fertilizer application activities on rainy days. They should harvest matured vegetables and fruits before the rainfall activities & cover young seedlings with polythene cover to protect them from heavy/intense shower.

Kharif Rice

Drain out excess water from the field. Maintain about 5-10cm standing water during panicle initiation stage.

Weather based Disease and pest of rice

- Forecasted cloudy and humid weather and the 20-25°C minimum temperature favors incidence of **stem borer and blast disease** in rice if the infestation will happen, spray Chlorantraniliprole 18.5% SC @ 60 ml/acre in 200 litres of water or Flubendiamide 20 WG 50g/acre in 200 litres of water to control **stem borer** and to control **blast** disease spray Tebuconazole 50%+ Trifloxystrobin 25% @ 0.4g/litre or, Carbendazim 50 WP @ 2g/litre of water.
- If **Brown Plant Hopper (BPH)** infestation will observe in rice, go for alternate wetting and drying. Whenever the number of hoppers/hill reaches 5-10, spray Pymetrozine 50% WG @ 120 g or Buprofezin 25% SC @ 330 ml or Imidacloprid 17.8% SL @ 50 ml per acre in the base of the plant. Before spraying remove water from the field.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre or Chlorpyrifos 20EC @ 500ml/acre to manage the pest.
- Due to likely water logging situation in Cuttack district as well as in the blocks **sheath blight** will be found in early transplanted rice then apply carbendazim 2g/ltr or propiconazole 1ml/ltr of water. In case of **Foot rot** spray the basal portion of the crop with 0.4% mancozeb or 0.2% carbendazim after drying the weather.

Maize:

- Provide drainage facility. Rainfall after a dry spell may aggravate infestation of fall army worm in maize, to control fall army worm in maize erect 4 pheromone traps /acre and spray 5 ml of 1500ppm neem oil /lit water. If more severe incidence observed spray Emamectin benzoate @ 0.5 g / lit of water. High humidity (>90%) and decrease in temperature (21-33°C) and continuous rain in the forecasted days may aggravate **downy mildew disease** in Maize. High humid conditions, temperature between 18-27 Deg. Celsius favors the occurrence of **leaf blight** in Maize. To control both disease spray Azoxystrobin + Mancozeb @ 300 g /ac.

Rabi-Groundnut:

- If groundnut is at harvesting stage then go for harvesting before rainy days. Anticipated humid, rainy weather favors infestation of **collar rot, alternaria leaf spot and rust** in groundnut. To control farmers may spray mancozeb/copper oxychloride 2g/liter water.

VEGETABLES:

- Drain out excess water from the field. Due to rain and high humidity, infestation of **blight and sucking pest** in vegetables likely to increase; farmers are advised to regular monitor the field. If observe, spray Mancozeb 75% WP @ 2g/l of water for blight and Imidacloprid 17.8% SL @ 0.25 ml/l of water for sucking pest.
- Due to rain and high humid condition during last week, **wilting** is likely to appear in young plants. Drain out water from the field. Spray mixing Streptocycline @ 20g with Carboxin+Thiram @ 400g or Metalaxyl 8%+Mancozeb 64% WP @ 400g per acre in 200 liters of water. Apply additional dose of potassic fertilizer to the crop.
- **Downy Mildew, vine rot, leaf spot and blight disease** in cucurbit vegetables, spray Carbendazim 12 % + Mancozeb 63 % WP @ 400 g/acre or Chlorothalonil @ 400 g/acre. Due to forecast of rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.

Brinjal, tomato and chilli seedlings should be planted in the main field. To prevent of soil borne diseases like **seedling rot, stem rot and wilt** diseases apply 3g of Copper oxychloride per litre of water around the main stem.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-435
Date- 12/09/2023

AGROMET ADVISORY BULLETIN FOR BANKI

District Past week average weather condition

The district received average 12mm rainfall during the last week. The mean daily maximum and minimum temperature was **32°C (-0.09°C from normal)** and **26°C (+1.6°C from normal)**, respectively. Mean relative humidity of the district was around 88-92%.

Forecasted period: Today(8.30IST) to 17th September (8.30IST)2023

Weather advisory: Light to moderate with isolated heavy to very heavy rainfall is forecasted in between upcoming 5 days in the block. The day temperature will be 30-32°C and night temperature will be 25-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 86-90% and 58-73% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Banki (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	12/09/2023	13/09/2023	14/09/2023	15/09/2023	16/09/2023
Rainfall(mm)	20	70	40	16	8
T-Max(°C)	32	31	30	31	30
T-Min(°C)	26	25	25	25	25
Cloud Cover(octa)	8	8	8	8	8
RH Max (%)	86	90	89	87	90
RH Min (%)	58	64	67	66	73
Wind Speed (Kmph)	7	6	7	5	5
Wind Direction (deg)	255	293	203	342	203

Agromet Advisory

General advisory: Keep the drainage channels open to drain out excess rain water from crop fields. They should stop irrigation, pesticide and fertilizer application activities on rainy days. They should harvest matured vegetables and fruits before the rainfall activities & cover young seedlings with ploythine cover to protect them from heavy/intense shower.

Kharif Rice

Drain out excess water from the field. Maintain about 5-10cm standing water during panicle initiation stage.

Weather based Disease and pest of rice

- Forecasted cloudy and humid weather and the 20-25°C minimum temperature favors incidence of **stem borer and blast disease** in rice if the infestation will happen, spray Chlorantraniliprole 18.5% SC @ 60 ml/acre in 200 litres of water or Flubendiamide 20 WG 50g/acre in 200 litres of water to control **stem borer** and to control **blast** disease spray Tebuconazole 50%+ Trifloxystrobin 25% @ 0.4g/litre or, Carbendazim 50 WP @ 2g/litre of water.
- If **Brown Plant Hopper (BPH) infestation** will observe in rice, go for alternate wetting and drying. Whenever the number of hoppers/hill reaches 5-10, spray Pymetrozine 50% WG @ 120 g or Buprofezin 25% SC @ 330 ml or Imidacloprid 17.8% SL @ 50 ml per acre in the base of the plant. Before spraying remove water from the field.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre or Chlorpyrifos 20EC @ 500ml/acre to manage the pest.
- Due to likely water logging situation in Cuttack district as well as in the blocks **sheath blight** will be found in early transplanted rice then apply carbendazim 2g/ltr or propiconazole 1ml/ltr of water. In case of **Foot rot**, spray the basal portion of the crop with 0.4% mancozeb or 0.2% carbendazim after drying the weather.
- Maize:**
- Provide drainage facility.** Rainfall after a dry spell may aggravate infestation of fall army worm in maize, to control fall armyworm in maize erect 4 pheromone traps /acre and spray 5 ml of 1500ppm neem oil /lit water. If more severe incidence observed spray Emamectin benzoate @ 0.5 g / lit of water. High humidity (>90%) and decrease in temperature (21-33°C) and continuous rain in the forecasted days may aggravate **downy mildew disease** in Maize. High humid conditions, temperature between 18-27 Deg. Celsius favors the occurrence of **leaf blight** in Maize. To control both disease spray Azoxystrobin + Mancozeb @ 300 g /ac.

Rabi-Groundnut:

- If groundnut is at harvesting stage then go for harvesting before rainy days. Anticipated humid, rainy weather favors infestation of **collar rot, alternaria leaf spot and rust** in groundnut. To control farmers may spray mancozeb/copper oxychloride 2g/liter water.

VEGETABLES:

- Drain out excess water from the field.** Due to rain and high humidity, infestation of **blight and sucking pest** in vegetables likely to increase; farmers are advised to regular monitor the field. If observe, spray Mancozeb 75% WP @ 2g/l of water for blight and Imidacloprid 17.8% SL @ 0.25 ml/l of water for sucking pest.
- Due to rain and high humid condition during last week, **wilting** is likely to appear in young plants. Drain out water from the field. Spray mixing Streptocycline @ 20g with Carboxin+Thiram @ 400g or Metalaxyl 8%+Mancozeb 64% WP @ 400g per acre in 200 liters of water. Apply additional dose of potassic fertilizer to the crop.
- Downy Mildew, vine rot, leaf spot and blight disease** in cucurbit vegetables, spray Carbendazim 12 % + Mancozeb 63 % WP @ 400 g/acre or Chlorothalonil @ 400 g/acre. Due to forecast of rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- Brinjal, tomato and chilli seedlings** should be planted in the main field. To prevent of soil borne diseases like **seedling rot, stem rot and wilt** diseases apply 3g of Copper oxychloride per litre of water around the main stem.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA INDIA METEOROLOGICAL DEPARTMENT DISTRICT AGROMET UNIT KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-435

Date- 12/09/2023

AGROMET ADVISORY BULLETIN FOR BANKI-DAMPARA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 20mm& district received average 12mm rainfall during the last week. The mean daily maximum and minimum temperature was 32°C (-0.09°C from normal) and 26°C (+1.6°C from normal), respectively. Mean relative humidity of the district was around 88-92%.

Forecasted period: Today(8.30IST) to 17th September (8.30IST)2023

Weather Advisory: Light to moderate with isolated heavy to very heavy rainfall is forecasted in between upcoming 5 days in the block. The day temperature will be 30-32°C and night temperature will be 25-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 87-90% and 59-74% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Banki-Dampada (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	12/09/2023	13/09/2023	14/09/2023	15/09/2023	16/09/2023
Rainfall(mm)	23	68	38	17	8
T-Max(°C)	32	31	31	31	30
T-Min(°C)	26	25	25	25	25
Cloud Cover(octa)	8	8	8	8	8
RH Max (%)	87	90	89	87	90
RH Min (%)	59	64	67	67	74
Wind Speed (Kmph)	7	6	7	5	5
Wind Direction (deg)	225	203	203	342	203

Agromet Advisory

General advisory: Keep the drainage channels open to drain out excess rain water from crop fields. They should stop irrigation, pesticide and fertilizer application activities on rainy days. They should harvest matured vegetables and fruits before the rainfall activities & cover young seedlings with polythene cover to protect them from heavy/intense shower.

Kharif Rice

Drain out excess water from the field. Maintain about 5-10cm standing water during panicle initiation stage.

Weather based Disease and pest of rice

- Forecasted cloudy and humid weather and the 20-25°C minimum temperature favors incidence of **stem borer and blast disease** in rice if the infestation will happen, spray Chlorantraniliprole 18.5% SC @ 60 ml/acre in 200 litres of water or Flubendiamide 20 WG 50g/acre in 200 litres of water to control **stem borer** and to control **blast** disease spray Tebuconazole 50%+ Trifloxystrobin 25% @ 0.4g/litre or, Carbendazim 50 WP @ 2g/litre of water.
- If **Brown Plant Hopper (BPH)** infestation will observe in rice, go for alternate wetting and drying. Whenever the number of hoppers/hill reaches 5-10, spray Pymetrozine 50% WG @ 120 g or Buprofezin 25% SC @ 330 ml or Imidacloprid 17.8% SL @ 50 ml per acre in the base of the plant. Before spraying remove water from the field.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre or Chlorpyrifos 20EC @ 500ml/acre to manage the pest.
- Due to likely water logging situation in Cuttack district as well as in the blocks **sheath blight** will be found in early transplanted rice then apply carbendazim 2g/ltr or propiconazole 1ml/ltr of water. In case of **Foot rot**, spray the basal portion of the crop with 0.4% mancozeb or 0.2% carbendazim after drying the weather.

Maize:

- **Provide drainage facility.** Rainfall after a dry spell may aggravate infestation of fall army worm in maize, to control fall armyworm in maize erect 4 pheromone traps /acre and spray 5 ml of 1500ppm neem oil /lit water. If more severe incidence observed spray Emamectin benzoate @ 0.5 g / lit of water. High humidity (>90%) and decrease in temperature (21-33°C) and continuous rain in the forecasted days may aggravate **downy mildew disease** in Maize. High humid conditions, temperature between 18-27 Deg. Celsius favors the occurrence of **leaf blight** in Maize. To control both disease spray Azoxystrobin + Mancozeb @ 300 g /ac.

Rabi-Groundnut:

- If groundnut is at harvesting stage then go for harvesting before rainy days. Anticipated humid, rainy weather favors infestation of **collar rot, alternaria leaf spot and rust** in groundnut. To control farmers may spray mancozeb/copper oxychloride 2g/liter water.

VEGETABLES:

- **Drain out excess water from the field.** Due to rain and high humidity, infestation of **blight and sucking pest** in vegetables likely to increase; farmers are advised to regular monitor the field. If observe, spray Mancozeb 75% WP @ 2g/l of water for blight and Imidacloprid 17.8% SL @ 0.25 ml/l of water for sucking pest.
- Due to rain and high humid condition during last week, **wilting** is likely to appear in young plants. Drain out water from the field. Spray mixing Streptocycline @ 20g with Carboxin+Thiram @ 400g or Metalaxyl 8%+Mancozeb 64% WP @ 400g per acre in 200 liters of water. Apply additional dose of potassic fertilizer to the crop.
- **Downy Mildew, vine rot, leaf spot and blight disease** in cucurbit vegetables, spray Carbendazim 12 % + Mancozeb 63 % WP @ 400 g/acre or Chlorothalonil @ 400 g/acre. Due to forecast of rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.

Brinjal, tomato and chilli seedlings should be planted in the main field. To prevent of soil borne diseases like **seedling rot, stem rot and wilt** diseases apply 3g of Copper oxychloride per litre of water around the main stem.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-435
Date- 12/09/2023

AGROMET ADVISORY BULLETIN FOR BARANGA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 12mm& district received average 12mm rainfall during the last week. The mean daily maximum and minimum temperature was **32°C (-0.09°C from normal)** and **26°C (+1.6°C from normal)**, respectively. Mean relative humidity of the district was around 88-92%.

Forecasted period: Today(8.30IST) to 17th September (8.30IST)2023

Weather advisory: *Light to moderate with isolated heavy to very heavy rainfall is forecasted in between upcoming 5 days in the block.* The day temperature will be **31-32°C** and night temperature will be **25-26°C** for upcoming 5 days. Maximum and minimum relative humidity will be 87-90% and 57-70% respectively. **There will be partly cloudy to overcast sky for next 5 days.**

Block	Baranga (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	12/09/2023	13/09/2023	14/09/2023	15/09/2023	16/09/2023
Rainfall(mm)	29	65	37	14	7
T-Max(°C)	32	31	32	31	31
T-Min(°C)	26	26	25	26	26
Cloud Cover	8	8	8	8	8
RH Max (%)	88	90	89	87	90
RH Min (%)	57	60	64	66	70
Wind Speed (Kmph)	8	8	8	5	6
Wind Direction (deg)	225	203	203	199	203

Agromet Advisory

General advisory: *Keep the drainage channels open to drain out excess rain water from crop fields. They should stop irrigation, pesticide and fertilizer application activities on rainy days. They should harvest matured vegetables and fruits before the rainfall activities & cover young seedlings with ploythine cover to protect them from heavy/intense shower.*

Kharif Rice

Drain out excess water from the field. Maintain about 5-10cm standing water during panicle initiation stage.

Weather based Disease and pest of rice

- **Forecasted cloudy and humid weather** and the 20-25°C minimum temperature favors incidence of **stem borer and blast disease** in rice if the infestation will happen, spray Chlorantraniliprole 18.5% SC @ 60 ml/acre in 200 litres of water or Flubendiamide 20 WG 50g/acre in 200 litres of water to control **stem borer** and to control **blast** disease spray Tebuconazole 50%+ Trifloxystrobin 25% @ 0.4g/litre or, Carbendazim 50 WP @ 2g/litre of water.
- If **Brown Plant Hopper (BPH) infestation** will observe in rice, go for alternate wetting and drying. Whenever the number of hoppers/hill reaches 5-10, spray Pymetrozine 50% WG @ 120 g or Buprofezin 25% SC @ 330 ml or Imidacloprid 17.8% SL @ 50 ml per acre in the base of the plant. Before spraying remove water from the field.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre or Chlorpyrifos 20EC @ 500ml/acre to manage the pest.
- Due to likely water logging situation in Cuttack district as well as in the blocks **sheath blight** will be found in early transplanted rice then apply carbendazim 2g/ltr or propiconazole 1ml/ltr of water. In case of **Foot rot**, spray the basal portion of the crop with 0.4% mancozeb or 0.2% carbendazim after drying the weather.
- **Maize:**
- **Provide drainage facility.** Rainfall after a dry spell may aggravate infestation of **fall army worm in maize**, to control **fall army worm** in maize erect 4 pheromone traps /acre and spray 5 ml of 1500ppm neem oil /lit water. If more severe incidence observed spray Emamectin benzoate @ 0.5 g / lit of water. High humidity (>90%) and decrease in temperature (21-33°C) and continuous rain in the forecasted days may aggravate **downy mildew disease** in Maize. High humid conditions, temperature between 18-27 Deg. Celsius favors the occurrence of **leaf blight** in Maize. To control both disease spray Azoxystrobin + Mancozeb @ 300 g /ac.

Rabi-Groundnut:

- If groundnut is at harvesting stage then go for harvesting before rainy days. Anticipated humid, rainy weather favors infestation of **collar rot, alternaria leaf spot and rust** in groundnut. To control farmers may spray mancozeb/copper oxychloride 2g/liter water.

VEGETABLES:

- **Drain out excess water from the field.** Due to rain and high humidity, infestation of **blight and sucking pest** in vegetables likely to increase; farmers are advised to regular monitor the field. If observe, spray Mancozeb 75% WP @ 2g/l of water for blight and Imidacloprid 17.8% SL @ 0.25 ml/l of water for sucking pest.
- Due to rain and high humid condition during last week, **wilting** is likely to appear in young plants. Drain out water from the field. Spray mixing Streptocycline @ 20g with Carboxin+Thiram @ 400g or Metalaxyl 8%+Mancozeb 64% WP @ 400g per acre in 200 liters of water. Apply additional dose of potassic fertilizer to the crop.
- **Downy Mildew, vine rot, leaf spot and blight disease** in cucurbit vegetables, spray Carbendazim 12 % + Mancozeb 63 % WP @ 400 g/acre or Chlorothalonil @ 400 g/acre. Due to forecast of rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- Brinjal, tomato and chilli seedlings** should be planted in the main field. To prevent of soil borne diseases like **seedling rot, stem rot and wilt** diseases apply 3g of Copper oxychloride per litre of water around the main stem.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA INDIA METEOROLOGICAL DEPARTMENT DISTRICT AGROMET UNIT KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

AGROMET ADVISORY BULLETIN FOR CUTTACK SADAR



Bulletin No.-435

Date- 12/09/2023

District Past week average weather condition and Block past week average Rainfall condition

The block received average 8mm & district received average 12mm rainfall during the last week. The mean daily maximum and minimum temperature was 32°C (-0.09°C from normal) and 26°C (+1.6°C from normal), respectively. Mean relative humidity of the district was around 88-92%.

Forecasted period: Today(8.30IST) to 17th September (8.30IST)2023

Weather advisory: Light to moderate with isolated heavy to very heavy rainfall is forecasted in between upcoming 5 days in the block.

The day temperature will be 31-32°C and night temperature will be 26°C for upcoming 5 days. Maximum and minimum relative humidity will be 86-90% and 56-67% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Cuttack sadar (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	12/09/2023	13/09/2023	14/09/2023	15/09/2023	16/09/2023
Rainfall(mm)	32	63	28	8	6
T-Max(°C)	32	32	32	31	31
T-Min(°C)	26	26	26	26	26
Cloud Cover(octa)	8	8	8	8	8
RH Max (%)	88	89	90	86	90
RH Min (%)	56	58	62	65	67
Wind Speed (Kmph)	8	8	8	6	7
Wind Direction (deg)	225	203	203	191	203

Agromet Advisory

General advisory: Keep the drainage channels open to drain out excess rain water from crop fields. They should stop irrigation, pesticide and fertilizer application activities on rainy days. They should harvest matured vegetables and fruits before the rainfall activities & cover young seedlings with ploythine cover to protect them from heavy/intense shower.

Kharif Rice

Drain out excess water from the field. Maintain about 5-10cm standing water during panicle initiation stage.

Weather based Disease and pest of rice

- Forecasted cloudy and humid weather and the 20-25°C minimum temperature favors incidence of **stem borer and blast disease** in rice if the infestation will happen, spray Chlorantraniliprole 18.5% SC @ 60 ml/acre in 200 litres of water or Flubendiamide 20 WG 50g/acre in 200 litres of water to control **stem borer** and to control **blast** disease spray Tebuconazole 50%+ Trifloxystrobin 25% @ 0.4g/litre or, Carbendazim 50 WP @ 2g/litre of water.
- If **Brown Plant Hopper (BPH) infestation** will observe in rice, go for alternate wetting and drying. Whenever the number of hoppers/hill reaches 5-10, spray Pymetrozine 50% WG @ 120 g or Buprofezin 25% SC @ 330 ml or Imidacloprid 17.8% SL @ 50 ml per acre in the base of the plant. Before spraying remove water from the field.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre or Chlorpyrifos 20EC @ 500ml/acre to manage the pest.
- Due to likely water logging situation in Cuttack district as well as in the blocks **sheath blight** will be found in early transplanted rice then apply carbendazim 2g/ltr or propiconazole 1ml/ltr of water. In case of **Foot rot** spray the basal portion of the crop with 0.4% mancozeb or 0.2% carbendazim after drying the weather.
- Sugarcane:** Wrapping and propping should be done when crop are 4-5 month old in medium and late duration varieties. For **early shoot borer** Spray Chlorpyrifos or Quinalophos @ 400 ml/ac in 200 lts of water after cessation of rain. For **Red rot**, avoid rationing in red rot affected area, spray the crop with Mancozeb 0.3% or carbendazim 0.1% at 15 days interval. Complete hoeing, weeding and apply fertilizer urea 100kg with 34 kg MOP in lines followed by earthing up during non rainy period.

VEGETABLES:

- Drain out excess water from the field.** Due to rain and high humidity, infestation of **blight and sucking pest** in vegetables likely to increase; farmers are advised to regular monitor the field. If observe, spray Mancozeb 75% WP @ 2g/l of water for blight and Imidacloprid 17.8% SL @ 0.25 ml/l of water for sucking pest.
- Due to rain and high humid condition during last week, **wilting** is likely to appear in young plants. Drain out water from the field. Spray mixing Streptocycline @ 20g with Carboxin+Thiram @ 400g or Metalaxyl 8%+Mancozeb 64% WP @ 400g per acre in 200 liters of water. Apply additional dose of potassic fertilizer to the crop.
- Downy Mildew, vine rot, leaf spot and blight disease** in cucurbit vegetables, spray Carbendazim 12 % + Mancozeb 63 % WP @ 400 g/acre or Chlorothalonil @ 400 g/acre. Due to forecast of rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval. Brinjal, tomato and chill seedlings should be planted in the main field. To prevent of soil borne diseases like **seedling rot, stem rot and wilt** diseases apply 3g of Copper oxychloride per litre of water around the main stem.

Fruits:

- In Banana Present weather conditions are favourable for the incidence and spread of sigatoka leaf spot. To curb the spread of this disease, go for spraying of propiconazole @ 0.1% (or) tebuconazole + trifloxystrobin @ 1ml per litre for 2 -3 times in 25 days interval. Go for Drenching of Carbendazim 0.2% or Propiconazole 0.5 ml/ litre once in a month for panama wilt of Banana disease.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

AGROMET ADVISORY BULLETIN FOR KANTAPADA



Bulletin No.-435

Date- 12/09/2023

District Past week average weather condition and Block past week average Rainfall condition

The block received average 15mm& district received average 12mm rainfall during the last week. The mean daily maximum and minimum temperature was **32°C (-0.09°C from normal)** and **26°C (+1.6°C from normal)**, respectively. Mean relative humidity of the district was around 88-92%.

Forecasted period: Today(8.30IST) to 17th September (8.30IST)2023

Weather advisory: *Light to moderate with isolated heavy to very heavy rainfall is forecasted in between upcoming 5 days in the block.* The day temperature will be **31-32°C** and night temperature will be **26°C** for upcoming 5 days. Maximum and minimum relative humidity will be 87-90% and 57-67% respectively. **There will be partly cloudy to overcast sky for next 5days.**

Block	Kantapada (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	12/09/2023	13/09/2023	14/09/2023	15/09/2023	16/09/2023
Rainfall(mm)	31	64	25	8	6
T-Max(°C)	32	32	32	31	31
T-Min(°C)	26	26	26	26	26
Cloud Cover(octa)	7	8	8	8	8
RH Max (%)	88	90	88	87	90
RH Min (%)	57	60	63	67	67
Wind Speed (Kmph)	9	8	8	6	6
Wind Direction (deg)	207	207	206	203	207

Agromet Advisory

General advisory: *Keep the drainage channels open to drain out excess rain water from crop fields. They should stop irrigation, pesticide and fertilizer application activities on rainy days. They should harvest matured vegetables and fruits before the rainfall activities & cover young seedlings with polythene cover to protect them from heavy/intense shower.*

Kharif Rice

Drain out excess water from the field. Maintain about 5-10cm standing water during panicle initiation stage.

Weather based Disease and pest of rice

- **Forecasted cloudy and humid weather** and the 20-25°C minimum temperature favors incidence of **stem borer and blast disease** in rice if the infestation will happen, spray Chlorantraniliprole 18.5% SC @ 60 ml/acre in 200 litres of water or Flubendiamide 20 WG 50g/acre in 200 litres of water to control **stem borer** and to control **blast** disease spray Tebuconazole 50%+ Trifloxystrobin 25% @ 0.4g/litre or, Carbendazim 50 WP @ 2g/litre of water.
- If **Brown Plant Hopper (BPH) infestation** will observe in rice, go for alternate wetting and drying. Whenever the number of hoppers/hill reaches 5-10, spray Pymetrozine 50% WG @ 120 g or Buprofezin 25% SC @ 330 ml or Imidacloprid 17.8% SL @ 50 ml per acre in the base of the plant. Before spraying remove water from the field.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre or Chlorpyrifos 20EC @ 500ml/acre to manage the pest.
- Due to likely water logging situation in Cuttack district as well as in the blocks **sheath blight** will be found in early transplanted rice then apply carbendazim 2g/ltr or propiconazole 1ml/ltr of water. In case of **Foot rot**, spray the basal portion of the crop with 0.4% mancozeb or 0.2% carbendazim after drying the weather.
- **Sugarcane:** Wrapping and propping should be done when crop are 4-5 month old in medium and late duration varieties. For **early shoot borer** Spray Chlorpyrifos or Quinalphos @ 400 ml/ac in 200 lts of water after cessation of rain. For **Red rot**, avoid rationing in red rot affected area, spray the crop with Mancozeb 0.3% or carbendazim 0.1% at 15 days interval. Complete hoeing, weeding and apply fertilizer urea 100kg with 34 kg MOP in lines followed by earthing up during non rainy period.

• **VEGETABLES:**

- **Drain out excess water from the field.** Due to rain and high humidity, infestation of **blight and sucking pest** in vegetables likely to increase; farmers are advised to regular monitor the field. If observe, spray Mancozeb 75% WP @ 2g/l of water for blight and Imidacloprid 17.8% SL @ 0.25 ml/l of water for sucking pest.
- Due to rain and high humid condition during last week, **wilting** is likely to appear in young plants. Drain out water from the field. Spray mixing Streptocycline @ 20g with Carboxin+Thiram @ 400g or Metalaxyl 8%+Mancozeb 64% WP @ 400g per acre in 200 liters of water. Apply additional dose of potassic fertilizer to the crop.
- **Downy Mildew, vine rot, leaf spot and blight disease** in cucurbit vegetables, spray Carbendazim 12 % + Mancozeb 63 % WP @ 400 g/acre or Chlorothalonil @ 400 g/acre. Due to forecast of rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval. Brinjal, tomato and chill seedlings should be planted in the main field. To prevent of soil borne diseases like **seedling rot, stem rot and wilt** diseases apply 3g of Copper oxychloride per litre of water around the main stem.

Fruits:

In Banana Present weather conditions are favourable for the incidence and spread of sigatoka leaf spot. To curb the spread of this disease, go for spraying of propiconazole @ 0.1% (or) tebuconazole + trifloxystrobin @ 1ml per litre for 2 -3 times in 25 days interval. Go for Drenching of Carbendazim 0.2% or Propiconazole 0.5 ml/ litre once in a month for panama wilt of Banana disease.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA INDIA METEOROLOGICAL DEPARTMENT DISTRICT AGROMET UNIT KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

AGROMET ADVISORY BULLETIN FOR MAHANGA



Bulletin No.-435

Date- 12/09/2023

District Past week average weather condition and Block past week average Rainfall condition

The block received average 11mm& district received average 12mm rainfall during the last week. The mean daily maximum and minimum temperature was 32°C (-0.09°C from normal) and 26°C (+1.6°C from normal), respectively. Mean relative humidity of the district was around 88-92%.

Forecasted period: Today(8.30IST) to 17th September (8.30IST)2023

Weather advisory:: Light to moderate with isolated heavy to very heavy rainfall is forecasted in between upcoming 5 days in the block. The day temperature will be and it will be 32-34°C and night temperature will be 26°C for upcoming 5 days. Maximum and minimum relative humidity will 86-90% and 52-63% respectively. **There will be partly cloudy to overcast sky for next 5 days.**

Block	Mahanga (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	12/09/2023	13/09/2023	14/09/2023	15/09/2023	16/09/2023
Rainfall(mm)	36	61	19	7	6
T-Max(°C)	34	33	32	32	33
T-Min(°C)	26	26	26	26	26
Cloud Cover(octal)	7	8	8	8	8
RH Max (%)	88	89	90	86	88
RH Min (%)	52	55	60	63	59
Wind Speed (Kmph)	9	9	8	7	8
Wind Direction (deg)	15	203	199	203	207

Agro-met Advisory

General advisory: Keep the drainage channels open to drain out excess rain water from crop fields. They should stop irrigation, pesticide and fertilizer application activities on rainy days. They should harvest matured vegetables and fruits before the rainfall activities & cover young seedlings with polythene cover to protect them from heavy/intense shower.

Kharif Rice

Drain out excess water from the field. Maintain about 5-10cm standing water during panicle initiation stage.

Weather based Disease and pest of rice

- **Forecasted cloudy and humid weather** and the 20-25°C minimum temperature favors incidence of **stem borer and blast disease** in rice if the infestation will happen, spray Chlorantraniliprole 18.5% SC @ 60 ml/acre in 200 litres of water or Flubendiamide 20 WG 50g/acre in 200 litres of water to control **stem borer** and to control **blast** disease spray Tebuconazole 50%+ Trifloxystrobin 25% @ 0.4g/litre or, Carbendazim 50 WP @ 2g/litre of water.
- If **Brown Plant Hopper (BPH) infestation** will observe in rice, go for alternate wetting and drying. Whenever the number of hoppers/hill reaches 5-10, spray Pymetrozine 50% WG @ 120 g or Buprofezin 25% SC @ 330 ml or Imidacloprid 17.8% SL @ 50 ml per acre in the base of the plant. Before spraying remove water from the field.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre or Chlorpyrifos 20EC @ 500ml/acre to manage the pest.
- Due to likely water logging situation in Cuttack district as well as in the blocks **sheath blight** will be found in early transplanted rice then apply carbendazim 2g/ltr or propiconazole 1ml/ltr of water. In case of **Foot rot**, spray the basal portion of the crop with 0.4% mancozeb or 0.2% carbendazim after drying the weather.
- **Sugarcane:** Wrapping and propping should be done when crop are 4-5 month old in medium and late duration varieties. For **early shoot borer** Spray Chlorpyrifos or Quinalophos @ 400 ml/ac in 200 lts of water after cessation of rain. For **Red rot**, avoid rationing in red rot affected area, spray the crop with Mancozeb 0.3% or carbendazim 0.1% at 15 days interval. Complete hoeing, weeding and apply fertilizer urea 100kg with 34 kg MOP in lines followed by earthing up during non rainy period.

VEGETABLES:

- **Drain out excess water from the field.** Due to rain and high humidity, infestation of **blight and sucking pest** in vegetables likely to increase; farmers are advised to regular monitor the field. If observe, spray Mancozeb 75% WP @ 2g/l of water for blight and Imidacloprid 17.8% SL @ 0.25 ml/l of water for sucking pest.
- Due to rain and high humid condition during last week, **wilting** is likely to appear in young plants. Drain out water from the field. Spray mixing Streptocycline @ 20g with Carboxin+Thiram @ 400g or Metalaxyl 8%+Mancozeb 64% WP @ 400g per acre in 200 liters of water. Apply additional dose of potassic fertilizer to the crop.
- **Downy Mildew, vine rot, leaf spot and blight disease** in cucurbit vegetables, spray Carbendazim 12 % + Mancozeb 63 % WP @ 400 g/acre or Chlorothalonil @ 400 g/acre. Due to forecast of rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval. Brinjal, tomato and chill seedlings should be planted in the main field. To prevent of soil borne diseases like **seedling rot, stem rot and wilt** diseases apply 3g of Copper oxychloride per litre of water around the main stem.

Fruits:

- In Banana Present weather conditions are favourable for the incidence and spread of sigatoka leaf spot. To curb the spread of this disease, go for spraying of propiconazole @ 0.1% (or) tebuconazole + trifloxystrobin @ 1ml per litre for 2 -3 times in 25 days interval. Go for Drenching of Carbendazim 0.2% or Propiconazole 0.5 ml/ litre once in a month for panama wilt of Banana disease.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)



Bulletin No.-435
Date- 12/09/2023

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

AGROMET ADVISORY BULLETIN FOR NARASINGHPUR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 8mm& district received average 12mm rainfall during the last week. The mean daily maximum and minimum temperature was **32°C (-0.09°C from normal)** and **26°C (+1.6°C from normal)**, respectively. Mean relative humidity of the district was around 88-92%.

Forecasted period: Today(8.30IST) to 17th September (8.30IST)2023

Weather advisory: *Light to moderate with isolated heavy to very heavy rainfall is forecasted in between upcoming 5 days in the block.* The day temperature will be **30-32°C** and night temperature will be **24°C** for upcoming 5 days. Maximum and minimum relative humidity will be 90-92% and 62-76% respectively. **There will be partly cloudy to overcast sky for next 5 days.**

Block	Narasinghpur (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	12/09/2023	13/09/2023	14/09/2023	15/09/2023	16/09/2023
Rainfall(mm)	18	70	31	14	7
T-Max(°C)	32	30	30	31	30
T-Min(°C)	24	24	24	24	24
Cloud Cover(octa)	8	8	8	8	8
RH Max (%)	90	92	92	91	92
RH Min (%)	62	69	67	66	76
Wind Speed (Kmph)	6	5	4	3	3
Wind Direction (deg)	270	293	281	203	161

Agromet Advisory

General advisory: *Keep the drainage channels open to drain out excess rain water from crop fields. They should stop irrigation, pesticide and fertilizer application activities on rainy days. They should harvest matured vegetables and fruits before the rainfall activities & cover young seedlings with polythene cover to protect them from heavy/intense shower.*

Kharif Rice

Drain out excess water from the field. Maintain about 5-10cm standing water during panicle initiation stage.

Weather based Disease and pest of rice

- **Forecasted cloudy and humid weather** and the 20-25°C minimum temperature favors incidence of **stem borer and blast disease** in rice if the infestation will happen, spray Chlorantraniliprole 18.5% SC @ 60 ml/acre in 200 litres of water or Flubendiamide 20 WG 50g/acre in 200 litres of water to control **stem borer** and to control **blast** disease spray Tebuconazole 50%+ Trifloxystrobin 25% @ 0.4g/litre or, Carbendazim 50 WP @ 2g/litre of water.
- If **Brown Plant Hopper (BPH)** infestation will observe in rice, go for alternate wetting and drying. Whenever the number of hoppers/hill reaches 5-10, spray Pymetrozine 50% WG @ 120 g or Buprofezin 25% SC @ 330 ml or Imidacloprid 17.8% SL @ 50 ml per acre in the base of the plant. Before spraying remove water from the field.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre or Chlorpyrifos 20EC @ 500ml/acre to manage the pest.
- Due to likely water logging situation in Cuttack district as well as in the blocks **sheath blight** will be found in early transplanted rice then apply carbendazim 2g/ltr or propiconazole 1ml/ltr of water. In case of **Foot rot** spray the basal portion of the crop with 0.4% mancozeb or 0.2% carbendazim after drying the weather.

Maize:

- **Provide drainage facility.** Rainfall after a dry spell may aggravate infestation of fall army worm in maize, to control fall armyworm in maize erect 4 pheromone traps /acre and spray 5 ml of 1500ppm neem oil /lit water. If more severe incidence observed spray Emamectin benzoate @ 0.5 g / lit of water. High humidity (>90%) and decrease in temperature (21-33°C) and continuous rain in the forecasted days may aggravate **downy mildew disease** in Maize. High humid conditions, temperature between 18-27 Deg. Celsius favors the occurrence of **leaf blight** in Maize. To control both disease spray Azoxystrobin + Mancozeb @ 300 g /ac.

Rabi-Groundnut:

- If groundnut is at harvesting stage then go for harvesting before rainy days. Anticipated humid, rainy weather favors infestation of **collar rot, alternaria leaf spot and rust** in groundnut. To control farmers may spray mancozeb/copper oxychloride 2g/liter water.

VEGETABLES:

- **Drain out excess water from the field.** Due to rain and high humidity, infestation of **blight and sucking pest** in vegetables likely to increase; farmers are advised to regular monitor the field. If observe, spray Mancozeb 75% WP @ 2g/l of water for blight and Imidacloprid 17.8% SL @ 0.25 ml/l of water for sucking pest.
- Due to rain and high humid condition during last week, **wilting** is likely to appear in young plants. Drain out water from the field. Spray mixing Streptocycline @ 20g with Carboxin+Thiram @ 400g or Metalaxyl 8%+Mancozeb 64% WP @ 400g per acre in 200 liters of water. Apply additional dose of potassic fertilizer to the crop.
- **Downy Mildew, vine rot, leaf spot and blight disease** in cucurbit vegetables, spray Carbendazim 12 % + Mancozeb 63 % WP @ 400 g/acre or Chlorothalonil @ 400 g/acre. Due to forecast of rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- Brinjal, tomato and chilli seedlings should be planted in the main field. To prevent of soil borne diseases like **seedling rot, stem rot and wilt** diseases apply 3g of Copper oxychloride per litre of water around the main stem.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-435
Date- 12/09/2023

AGROMET ADVISORY BULLETIN FOR NIALI

District Past week average weather condition and Block past week average Rainfall condition

The block received average 15mm& district received average 12mm rainfall during the last week. The mean daily maximum and minimum temperature was **32°C (-0.09°C from normal)** and **26°C (+1.6°C from normal)**, respectively. Mean relative humidity of the district was around 88-92%.

Forecasted period: Today(8.30IST) to 17th September (8.30IST)2023

Weather advisory: *Light to moderate with isolated heavy to very heavy rainfall is forecasted in between upcoming 5 days in the block.* The day temperature will be **31-33°C** and night temperature will be **25-26°C** for upcoming 5 days. Maximum and minimum relative humidity will be 87-90% and 59-68% respectively. **There will be partly cloudy to overcast sky for next 5days.**

Block	Niali (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	12/09/2023	13/09/2023	14/09/2023	15/09/2023	16/09/2023
Rainfall(mm)	31	62	24	7	6
T-Max(°C)	33	32	32	31	32
T-Min(°C)	26	25	26	26	26
Cloud Cover(octa)	7	8	8	8	8
RH Max (%)	87	90	87	87	89
RH Min (%)	59	62	64	68	67
Wind Speed(Kmph)	9	8	9	6	7
Wind Direction(deg)	225	225	206	205	210

Agromet Advisory

General advisory: *Keep the drainage channels open to drain out excess rain water from crop fields. They should stop irrigation, pesticide and fertilizer application activities on rainy days. They should harvest matured vegetables and fruits before the rainfall activities & cover young seedlings with ploythine cover to protect them from heavy/intense shower.*

Kharif Rice

Drain out excess water from the field. Maintain about 5-10cm standing water during panicle initiation stage.

Weather based Disease and pest of rice

- **Forecasted cloudy and humid weather** and the 20-25°C minimum temperature favors incidence of **stem borer and blast disease** in rice if the infestation will happen, spray Chlorantraniliprole 18.5% SC @ 60 ml/acre in 200 litres of water or Flubendiamide 20 WG 50g/acre in 200 litres of water to control **stem borer** and to control **blast** disease spray Tebuconazole 50%+ Trifloxystrobin 25% @ 0.4g/litre or, Carbendazim 50 WP @ 2g/litre of water.
- If **Brown Plant Hopper (BPH)** infestation will observe in rice, go for alternate wetting and drying. Whenever the number of hoppers/hill reaches 5-10, spray Pymetrozine 50% WG @ 120 g or Buprofezin 25% SC @ 330 ml or Imidacloprid 17.8% SL @ 50 ml per acre in the base of the plant. Before spraying remove water from the field.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre or Chlorpyrifos 20EC @ 500ml/acre to manage the pest.
- Due to likely water logging situation in Cuttack district as well as in the blocks **sheath blight** will be found in early transplanted rice then apply carbendazim 2g/ltr or propiconazole 1ml/ltr of water. In case of **Foot rot**, spray the basal portion of the crop with 0.4% mancozeb or 0.2% carbendazim after drying the weather.
- **Sugarcane:** Wrapping and propping should be done when crop are 4-5 month old in medium and late duration varieties. For **early shoot borer** Spray Chlorpyrifos or Quinalophos @ 400 ml/ac in 200 lts of water after cessation of rain. For **Red rot**, avoid rationing in red rot affected area, spray the crop with Mancozeb 0.3% or carbendazim 0.1% at 15 days interval. Complete hoeing, weeding and apply fertilizer urea 100kg with 34 kg MOP in lines followed by earthing up during non rainy period.

VEGETABLES:

- **Drain out excess water from the field.** Due to rain and high humidity, infestation of **blight and sucking pest** in vegetables likely to increase; farmers are advised to regular monitor the field. If observe, spray Mancozeb 75% WP @ 2g/l of water for blight and Imidacloprid 17.8% SL @ 0.25 ml/l of water for sucking pest.
- Due to rain and high humid condition during last week, **wilting** is likely to appear in young plants. Drain out water from the field. Spray mixing Streptocycline @ 20g with Carboxin+Thiram @ 400g or Metalaxyl 8%+Mancozeb 64% WP @ 400g per acre in 200 liters of water. Apply additional dose of potassic fertilizer to the crop.
- **Downy Mildew, vine rot, leaf spot and blight disease** in cucurbit vegetables, spray Carbendazim 12 % + Mancozeb 63 % WP @ 400 g/acre or Chlorothalonil @ 400 g/acre. Due to forecast of rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval. Brinjal, tomato and chill seedlings should be planted in the main field. To prevent of soil borne diseases like **seedling rot, stem rot and wilt** diseases apply 3g of Copper oxychloride per litre of water around the main stem.

Fruits:

- In Banana Present weather conditions are favourable for the incidence and spread of sigatoka leaf spot. To curb the spread of this disease, go for spraying of propiconazole @ 0.1% (or) tebuconazole + trifloxystrobin @ 1ml per litre for 2 -3 times in 25 days interval. Go for Drenching of Carbendizim 0.2% or Propiconazole 0.5 ml/ litre once in a month for panama wilt of Banana disease.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK



(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-435

Date- 12/09/2023

AGROMET ADVISORY BULLETIN FOR NISCHINTAKOILI

District Past week average weather condition and Block past week average Rainfall condition

The block received average 13mm& district received average 12mm rainfall during the last week. The mean daily maximum and minimum temperature was 32°C (-0.09°C from normal) and 26°C (+1.6°C from normal), respectively. Mean relative humidity of the district was around 88-92%.

Forecasted period: Today(8.30IST) to 17th September (8.30IST)2023

Weather advisory: Light to moderate with isolated heavy to very heavy rainfall is forecasted in between upcoming 5 days in the block. The day temperature will be 32-33°C and night temperature will be 26-27°C for upcoming 5 days. Maximum and minimum relative humidity will be 87-89% and 53-64% respectively. **There will be partly cloudy to overcast sky for next 5 days.**

Block	Nischintakoili (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	12/09/2023	13/09/2023	14/09/2023	15/09/2023	16/09/2023
Rainfall(mm)	41	63	18	7	6
T-Max(°C)	33	32	32	32	33
T-Min(°C)	26	26	26	27	26
Cloud Cover(octa)	7	8	8	8	8
RH Max (%)	88	89	89	87	88
RH Min (%)	53	56	60	64	59
Wind Speed (Kmph)	9	9	8	7	8
Wind Direction (deg)	207	198	199	203	207

Agromet Advisory

General advisory: Keep the drainage channels open to drain out excess rain water from crop fields. They should stop irrigation, pesticide and fertilizer application activities on rainy days. They should harvest matured vegetables and fruits before the rainfall activities & cover young seedlings with polythene cover to protect them from heavy/intense shower.

Kharif Rice

Drain out excess water from the field. Maintain about 5-10cm standing water during panicle initiation stage.

Weather based Disease and pest of rice

- **Forecasted cloudy and humid weather** and the 20-25°C minimum temperature favors incidence of **stem borer and blast disease** in rice if the infestation will happen, spray Chlorantraniliprole 18.5% SC @ 60 ml/acre in 200 litres of water or Flubendiamide 20 WG 50g/acre in 200 litres of water to control **stem borer** and to control **blast** disease spray Tebuconazole 50%+ Trifloxystrobin 25% @ 0.4g/litre or, Carbendazim 50 WP @ 2g/litre of water.
- If **Brown Plant Hopper (BPH) infestation** will observe in rice, go for alternate wetting and drying. Whenever the number of hoppers/hill reaches 5-10, spray Pymetrozine 50% WG @ 120 g or Buprofezin 25% SC @ 330 ml or Imidacloprid 17.8% SL @ 50 ml per acre in the base of the plant. Before spraying remove water from the field.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre or Chlorpyrifos 20EC @ 500ml/acre to manage the pest.
- Due to likely water logging situation in Cuttack district as well as in the blocks **sheath blight** will be found in early transplanted rice then apply carbendazim 2g/ltr or propiconazole 1ml/ltr of water. In case of **Foot rot**, spray the basal portion of the crop with 0.4% mancozeb or 0.2% carbendazim after drying the weather.
- **Sugarcane:** Wrapping and propping should be done when crop are 4-5 month old in medium and late duration varieties. For **early shoot borer** Spray Chlorpyrifos or Quinalophos @ 400 ml/ac in 200 lts of water after cessation of rain. For **Red rot**, avoid rationing in red rot affected area, spray the crop with Mancozeb 0.3% or carbendazim 0.1% at 15 days interval. Complete hoeing, weeding and apply fertilizer urea 100kg with 34 kg MOP in lines followed by earthing up during non rainy period.

• VEGETABLES:

- **Drain out excess water from the field.** Due to rain and high humidity, infestation of **blight and sucking pest** in vegetables likely to increase; farmers are advised to regular monitor the field. If observe, spray Mancozeb 75% WP @ 2g/l of water for blight and Imidacloprid 17.8% SL @ 0.25 ml/l of water for sucking pest.
- Due to rain and high humid condition during last week, **wilting** is likely to appear in young plants. Drain out water from the field. Spray mixing Streptocycline @ 20g with Carboxin+Thiram @ 400g or Metalaxyl 8%+Mancozeb 64% WP @ 400g per acre in 200 liters of water. Apply additional dose of potassic fertilizer to the crop.
- **Downy Mildew, vine rot, leaf spot and blight disease** in cucurbit vegetables, spray Carbendazim 12 % + Mancozeb 63 % WP @ 400 g/acre or Chlorothalonil @ 400 g/acre. Due to forecast of rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval. Brinjal, tomato and chill seedlings should be planted in the main field. To prevent of soil borne diseases like **seedling rot, stem rot and wilt** diseases apply 3g of Copper oxychloride per litre of water around the main stem.

Fruits:

- In Banana Present weather conditions are favourable for the incidence and spread of sigatoka leaf spot. To curb the spread of this disease, go for spraying of propiconazole @ 0.1% (or) tebuconazole + trifloxystrobin @ 1ml per litre for 2 -3 times in 25 days interval. Go for Drenching of Carbendazim 0.2% or Propiconazole 0.5 ml/ litre once in a month for panama wilt of Banana disease.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-435
Date- 12/09/2023

AGROMET ADVISORY BULLETIN FOR SALEPUR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 9mm & district received average 12mm rainfall during the last week. The mean daily maximum and minimum temperature was 32°C (-0.09°C from normal) and 26°C (+1.6°C from normal), respectively. Mean relative humidity of the district was around 88-92%.

Forecasted period: Today(8.30IST) to 17th September (8.30IST)2023

Weather advisory:: Light to moderate with isolated heavy to very heavy rainfall is forecasted in between upcoming 5 days in the block. The day temperature will be 32-33°C and night temperature will be 26-27°C for upcoming 5 days. Maximum and minimum relative humidity will be 87-89% and 53-65% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Salepur (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	12/09/2023	13/09/2023	14/09/2023	15/09/2023	16/09/2023
Rainfall(mm)	39	62	17	7	6
T-Max(°C)	33	32	32	32	33
T-Min(°C)	26	26	26	27	26
Cloud Cover(octa)	8	8	8	8	8
RH Max (%)	89	89	89	87	89
RH Min (%)	53	56	61	65	59
Wind Speed (Kmph)	9	9	8	7	8
Wind Direction (deg)	207	198	199	199	207

Agromet Advisory

General advisory: Keep the drainage channels open to drain out excess rain water from crop fields. They should stop irrigation, pesticide and fertilizer application activities on rainy days. They should harvest matured vegetables and fruits before the rainfall activities & cover young seedlings with polythene cover to protect them from heavy/intense shower.

Kharif Rice

Drain out excess water from the field. Maintain about 5-10cm standing water during panicle initiation stage.

Weather based Disease and pest of rice

- **Forecasted cloudy and humid weather** and the 20-25°C minimum temperature favors incidence of **stem borer and blast disease** in rice if the infestation will happen, spray Chlorantraniliprole 18.5% SC @ 60 ml/acre in 200 litres of water or Flubendiamide 20 WG 50g/acre in 200 litres of water to control **stem borer** and to control **blast** disease spray Tebuconazole 50%+ Trifloxystrobin 25% @ 0.4g/litre or, Carbendazim 50 WP @ 2g/litre of water.
- If **Brown Plant Hopper (BPH)** infestation will observe in rice, go for alternate wetting and drying. Whenever the number of hoppers/hill reaches 5-10, spray Pymetrozine 50% WG @ 120 g or Buprofezin 25% SC @ 330 ml or Imidacloprid 17.8% SL @ 50 ml per acre in the base of the plant. Before spraying remove water from the field.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre or Chlorpyrifos 20EC @ 500ml/acre to manage the pest.
- Due to likely water logging situation in Cuttack district as well as in the blocks **sheath blight** will be found in early transplanted rice then apply carbendazim 2g/ltr or propiconazole 1ml/ltr of water. In case of **Foot rot**, spray the basal portion of the crop with 0.4% mancozeb or 0.2% carbendazim after drying the weather.
- **Sugarcane:** Wrapping and propping should be done when crop are 4-5 month old in medium and late duration varieties. For **early shoot borer** Spray Chlorpyrifos or Quinalphos @ 400 ml/ac in 200 lts of water after cessation of rain. For **Red rot**, avoid rationing in red rot affected area, spray the crop with Mancozeb 0.3% or carbendazim 0.1% at 15 days interval. Complete hoeing, weeding and apply fertilizer urea 100kg with 34 kg MOP in lines followed by earthing up during non rainy period.

• VEGETABLES:

- **Drain out excess water from the field.** Due to rain and high humidity, infestation of **blight and sucking pest** in vegetables likely to increase; farmers are advised to regular monitor the field. If observe, spray Mancozeb 75% WP @ 2g/l of water for blight and Imidacloprid 17.8% SL @ 0.25 ml/l of water for sucking pest.
- Due to rain and high humid condition during last week, **wilting** is likely to appear in young plants. Drain out water from the field. Spray mixing Streptocycline @ 20g with Carboxin+Thiram @ 400g or Metalaxyl 8%+Mancozeb 64% WP @ 400g per acre in 200 liters of water. Apply additional dose of potassic fertilizer to the crop.
- **Downy Mildew, vine rot, leaf spot and blight disease** in cucurbit vegetables, spray Carbendazim 12 % + Mancozeb 63 % WP @ 400 g/acre or Chlorothalonil @ 400 g/acre. Due to forecast of rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval. Brinjal, tomato and chill seedlings should be planted in the main field. To prevent of soil borne diseases like **seedling rot, stem rot and wilt** diseases apply 3g of Copper oxychloride per litre of water around the main stem.

Fruits:

- In Banana Present weather conditions are favourable for the incidence and spread of sigatoka leaf spot. To curb the spread of this disease, go for spraying of propiconazole @ 0.1% (or) tebuconazole + trifloxystrobin @ 1ml per litre for 2 -3 times in 25 days interval. Go for Drenching of Carbendazim 0.2% or Propiconazole 0.5 ml/ litre once in a month for panama wilt of Banana disease.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA INDIA METEOROLOGICAL DEPARTMENT DISTRICT AGROMET UNIT KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-435

Date- 12/09/2023

AGROMET ADVISORY BULLETIN FOR TANGI-CHOUDWAR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 7mm & district received average 12mm rainfall during the last week. The mean daily maximum and minimum temperature was 32°C (-0.09°C from normal) and 26°C (+1.6°C from normal), respectively. Mean relative humidity of the district was around 88-92%.

Forecasted period: Today (8.30 IST) to 17th September (8.30 IST) 2023

Weather advisory: Light to moderate with isolated heavy to very heavy rainfall is forecasted in between upcoming 5 days in the block. The day temperature will be 32-33°C and night temperature will be 26-27°C for upcoming 5 days. Maximum and minimum relative humidity will be 84-86% and 57-68% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Tangi-Choudwar (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	12/09/2023	13/09/2023	14/09/2023	15/09/2023	16/09/2023
Rainfall (mm)	30	64	20	7	6
T-Max (°C)	33	32	32	32	33
T-Min (°C)	26	26	26	27	26
Cloud Cover (octa)	7	8	8	8	8
RH Max (%)	85	86	84	84	86
RH Min (%)	59	57	62	68	63
Wind Speed (Kmph)	12	11	11	9	10
Wind Direction (deg)	243	248	203	198	198

Agromet Advisory

General advisory: Keep the drainage channels open to drain out excess rain water from crop fields. They should stop irrigation, pesticide and fertilizer application activities on rainy days. They should harvest matured vegetables and fruits before the rainfall activities & cover young seedlings with polythene cover to protect them from heavy/intense shower.

Kharif Rice

Drain out excess water from the field. Maintain about 5-10cm standing water during panicle initiation stage.

Weather based Disease and pest of rice

- Forecasted cloudy and humid weather and the 20-25°C minimum temperature favors incidence of **stem borer and blast disease** in rice if the infestation will happen, spray Chlorantraniliprole 18.5% SC @ 60 ml/acre in 200 litres of water or Flubendiamide 20 WG 50g/acre in 200 litres of water to control **stem borer** and to control **blast** disease spray Tebuconazole 50%+ Trifloxystrobin 25% @ 0.4g/litre or, Carbendazim 50 WP @ 2g/litre of water.
- If **Brown Plant Hopper (BPH)** infestation will observe in rice, go for alternate wetting and drying. Whenever the number of hoppers/hill reaches 5-10, spray Pymetrozine 50% WG @ 120 g or Buprofezin 25% SC @ 330 ml or Imidacloprid 17.8% SL @ 50 ml per acre in the base of the plant. Before spraying remove water from the field.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre or Chlorpyrifos 20EC @ 500ml/acre to manage the pest.
- Due to likely water logging situation in Cuttack district as well as in the blocks **sheath blight** will be found in early transplanted rice then apply carbendazim 2g/ltr or propiconazole 1ml/ltr of water. In case of **Foot rot**, spray the basal portion of the crop with 0.4% mancozeb or 0.2% carbendazim after drying the weather.

Maize:

- Provide drainage facility. Rainfall after a dry spell may aggravate infestation of fall army worm in maize, to control fall armyworm in maize erect 4 pheromone traps /acre and spray 5 ml of 1500ppm neem oil /lit water. If more severe incidence observed spray Emamectin benzoate @ 0.5 g / lit of water. High humidity (>90%) and decrease in temperature (21-33°C) and continuous rain in the forecasted days may aggravate **downy mildew disease** in Maize. High humid conditions, temperature between 18-27 Deg. Celsius favors the occurrence of **leaf blight** in Maize. To control both disease spray Azoxystrobin + Mancozeb @ 300 g /ac.

Rabi-Groundnut:

- If groundnut is at harvesting stage then go for harvesting before rainy days. Anticipated humid, rainy weather favors infestation of **collar rot, alternaria leaf spot and rust** in groundnut. To control farmers may spray mancozeb/copper oxychloride 2g/liter water.

VEGETABLES:

- Drain out excess water from the field. Due to rain and high humidity, infestation of **blight and sucking pest** in vegetables likely to increase; farmers are advised to regular monitor the field. If observe, spray Mancozeb 75% WP @ 2g/l of water for blight and Imidacloprid 17.8% SL @ 0.25 ml/l of water for sucking pest.
- Due to rain and high humid condition during last week, **wilting** is likely to appear in young plants. Drain out water from the field. Spray mixing Streptocycline @ 20g with Carboxin+Thiram @ 400g or Metalaxyl 8%+Mancozeb 64% WP @ 400g per acre in 200 liters of water. Apply additional dose of potassic fertilizer to the crop.
- **Downy Mildew, vine rot, leaf spot and blight disease** in cucurbit vegetables, spray Carbendazim 12 % + Mancozeb 63 % WP @ 400 g/acre or Chlorothalonil @ 400 g/acre. Due to forecast of rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- Brinjal, tomato and chilli seedlings should be planted in the main field. To prevent of soil borne diseases like **seedling rot, stem rot and wilt** diseases apply 3g of Copper oxychloride per litre of water around the main stem.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427



GRAMIN KRISHIMAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)



Bulletin No.-435
Date- 12/09/2023

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

AGROMET ADVISORY BULLETIN FOR TIGIRIA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 12mm& district received average 12mm rainfall during the last week. The mean daily maximum and minimum temperature was **32°C (-0.09°C from normal)** and **26°C (+1.6°C from normal)**, respectively. Mean relative humidity of the district was around 88-92%.

Forecasted period: Today(8.30IST) to 17th September (8.30IST)2023

Weather advisory: *Light to moderate with isolated heavy to very heavy rainfall is forecasted in between upcoming 5 days in the block.* The day temperature will be **30-33°C** and night temperature will be **25-26°C** for upcoming 5 days. Maximum and minimum relative humidity will be 88-91% and 59-75% respectively. **There will be partly cloudy to overcast sky for next 5 days.**

Block	Tigiria (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	12/09/2023	13/09/2023	14/09/2023	15/09/2023	16/09/2023
Rainfall(mm)	25	67	28	10	6
T-Max(°C)	33	31	31	31	30
T-Min(°C)	26	25	25	25	25
Cloud Cover(octa)	8	8	8	8	8
RH Max (%)	88	91	91	88	91
RH Min (%)	59	64	67	68	75
Wind Speed (Kmph)	7	6	7	5	5
Wind Direction (deg)	259	315	203	342	198

Agro met Advisory

General advisory: *Keep the drainage channels open to drain out excess rain water from crop fields. They should stop irrigation, pesticide and fertilizer application activities on rainy days. They should harvest matured vegetables and fruits before the rainfall activities & cover young seedlings with polythene cover to protect them from heavy/intense shower.*

Kharif Rice

Drain out excess water from the field. Maintain about 5-10cm standing water during panicle initiation stage.

Weather based Disease and pest of rice

- Forecasted cloudy and humid weather and the 20-25°C minimum temperature favors incidence of **stem borer and blast disease** in rice if the infestation will happen, spray Chlorantraniliprole 18.5% SC @ 60 ml/acre in 200 litres of water or Flubendiamide 20 WG 50g/acre in 200 litres of water to control **stem borer** and to control **blast** disease spray Tebuconazole 50%+ Trifloxystrobin 25% @ 0.4g/litre or, Carbendazim 50 WP @ 2g/litre of water.
- If **Brown Plant Hopper (BPH) infestation** will observe in rice, go for alternate wetting and drying. Whenever the number of hoppers/hill reaches 5-10, spray Pymetrozine 50% WG @ 120 g or Buprofezin 25% SC @ 330 ml or Imidacloprid 17.8% SL @ 50 ml per acre in the base of the plant. Before spraying remove water from the field.
- In **Gall midge** affected due to continuous cloudy and rainy weather in late transplanted rice apply Thiamethoxam 25% WG @ 40g/acre or Fipronil 5% SC @ 400ml/acre or Chlorpyrifos 20EC @ 500ml/acre to manage the pest.
- Due to likely water logging situation in Cuttack district as well as in the blocks **sheath blight** will be found in early transplanted rice then apply carbendazim 2g/ltr or propiconazole 1ml/ltr of water. In case of **Foot rot**, spray the basal portion of the crop with 0.4% mancozeb or 0.2% carbendazim after drying the weather.
- **Maize:**
- **Provide drainage facility.** Rainfall after a dry spell may aggravate infestation of fall army worm in maize, to control fall armyworm in maize erect 4 pheromone traps /acre and spray 5 ml of 1500ppm neem oil /lit water. If more severe incidence observed spray Emamectin benzoate @ 0.5 g / lit of water. High humidity (>90%) and decrease in temperature (21-33°C) and continuous rain in the forecasted days may aggravate **downy mildew disease** in Maize. High humid conditions, temperature between 18-27 Deg. Celsius favors the occurrence of **leaf blight** in Maize. To control both disease spray Azoxystrobin + Mancozeb @ 300 g /ac.

Rabi-Groundnut:

- If groundnut is at harvesting stage then go for harvesting before rainy days. Anticipated humid, rainy weather favors infestation of **collar rot, alternaria leaf spot and rust** in groundnut. To control farmers may spray mancozeb/copper oxychloride 2g/liter water.

• VEGETABLES:

- **Drain out excess water from the field.** Due to rain and high humidity, infestation of **blight and sucking pest** in vegetables likely to increase; farmers are advised to regular monitor the field. If observe, spray Mancozeb 75% WP @ 2g/l of water for blight and Imidacloprid 17.8% SL @ 0.25 ml/l of water for sucking pest.
- Due to rain and high humid condition during last week, **wilting** is likely to appear in young plants. Drain out water from the field. Spray mixing Streptocycline @ 20g with Carboxin+Thiram @ 400g or Metalaxyl 8%+Mancozeb 64% WP @ 400g per acre in 200 liters of water. Apply additional dose of potassic fertilizer to the crop.
- **Downy Mildew, vine rot, leaf spot and blight disease** in cucurbit vegetables, spray Carbendazim 12 % + Mancozeb 63 % WP @ 400 g/acre or Chlorothalonil @ 400 g/acre. Due to forecast of rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- Brinjal, tomato and chilli seedlings should be planted in the main field. To prevent of soil borne diseases like **seedling rot, stem rot and wilt** diseases apply 3g of Copper oxychloride per litre of water around the main stem.

Dr Sujata Sathy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-435
 Date- 12/09/2023

				
MEGHDOOT APP	DAMINI APP	FACEBOOK (Damu Cuttack)	BLOCK SPECIFIC WHAT'S APP GROUPS	@DAMUCUTTACK



- Kindly install and use **“Meghdoot mobile App”** for Cuttack district weather forecast, warnings and weather based agromet advisory services.

<https://play.google.com/store/apps/details?id=com.aas.meghdoot>

- Kindly install and use **“Damini mobile App”** for location specific for lightening forecast and warnings.

<https://play.google.com/store/apps/details?id=com.lightening.live.damini>

- Follow the **block specific whats app groups** and join in your block wise groups for getting weather based agromet advisory services twice in a week (Tuesday and Friday) and other weather warning based farm management advisories.
- Follow the **Official Website links** to get the weather based agromet advisory service bulletins (**English & Odia**) twice in a week (Tuesday and Friday).

1	ICAR-NRRI, Cuttack, Odisha	https://icar-nrri.in/agromate/	
2	Agrimet division, IMD, Pune	https://imdagrimet.gov.in/imdproject/AGIndex.php	

Dr Sujata Sethy
 (Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
 SMS (Agrometeorology)
 Contact No-9861445080/7008453427