



GRAMIN KRISHI MAUSAM SEWA

INDIA METEOROLOGICAL DEPARTMENT

DISTRICT AGROMET UNIT

KRISHI VIGYAN KENDRA CUTTACK

(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com




Bulletin No.-485

Date- 12/04/2024

Cuttack District Block Wise Agro-met Advisory Service Bulletin

*Friday, 12th April, 2024
(For the period 12th to 16th April, 2024)*



*Issued by
District Agromet Unit,
KrishiVigyan Kendra Cuttack*

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



STANDARD PRECIPITATION INDEX AND NDVI MAP

Standard Precipitation Index (SPI)	Normalized difference vegetation Index (NDVI)
<p align="center">Standardised Precipitation Index for the period 28th September to 25th October 2023</p>	<p align="center">NOAA/VIIRS/BLENDED NDVI Composite ending on Week no 13 (25.03.2024 to 31.03.2024) over Agricultural region</p> <p align="center">Odisha</p> <p align="center">Agriculture vigour is good at isolated places over bargarh district of the state.</p>
<p><i>As per SPI value (cumulative 4weeks -28 September to 25 October 2023), mildly dry condition was there in Cuttack district.</i></p>	<p><i>According to NDVI map (25.03.2024 to 31.03.2024) the agriculture vigor was very lighttolightover maximum western blocksof Cuttack district and light to moderate over isolated places of Mahanga, Nischintakoili, Tangichoudwar, Salepur, Kantapada and Banki-damapada .</i></p>

LEGEND (Medium range block level rainfall forecast)

	No Rainfall	00 mm
	Very light rainfall	01-05 mm
	Light rainfall	06-10 mm
	Light to moderate rainfall	11-30 mm
	Moderate to heavy rainfall	31-65 mm
	Heavy to very heavy rainfall	66-115 mm
	Extremely heavy rainfall	>116 mm

LEGEND (Past week temperature)

Color code	Temperature	Departure
	Below Normal	< 2°C from normal
	Normal	-2°C to +2°C from normal
	Above Normal	>2°C from normal

Dr Sujata Sethy
 (Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
 SMS (Agrometeorology)
 Contact No-9861445080/7008453427














GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-485
Date- 12/04/2024

Weather (Past and Forecasted) based Disease and pest in Major Crops and vegetables

Sl No	Crop	Stage and condition of the crop	Anticipated Disease and pest	
1	Pulses (Green gram and black gram)	Flowering to pod development		Rust
				Wilting
2	Maize	Vegetative		False army worm
				Red hairy catterpillar
3	Brinjal	Fructing		Fruit and shoot borer
				Phomopsis Blight
				Epilachna Beetle/Hadda beetle
				Cercospera leaf spot
				Helicoverpa catterpillar
4	Tomato	Fruting		Bacterial wilt
5	Okra	Vegetative to fructing		Yellow vein mosaiv(YMV)

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area












Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-485
Date- 12/04/2024

6	Cucurbits	Flowering to Fruiting stage		Red Pumpkin beetle
				Leaf spot
				Late blight
				Stem Rot
				Downy Mildew
				Fruit rot
7	Mango	Fruiting stage		Fruit borer
				Fruit drop
				Bacterial Canker

Dr Sujata Sethy
 (Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
 Contact No-9861445080/7008453427



INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-485
Date- 12/04/2024

AGROMET ADVISORY BULLETIN FOR ATHAGAD

District Past week average weather condition and Block past week average Rainfall condition

The block received average 2 mm and district received average 2mm rainfall during the last week. The mean daily maximum and minimum temperature was 35.47°C (-1.95 from normal) and 23.37°C (-0.3 from normal), respectively. The mean relative humidity was 61-67%.

Forecasted period: Today(8.30IST) to 17th April (8.30IST)2024

Weather advisory: Dry weather is forecasted for today to Tuesday in the block. The day temperature will be 32-37°C and night temperature will be 24-27°C for upcoming 5 days. Maximum and minimum relative humidity will be 48-81% and 19-21% respectively. There will be clear sky for next 5days.

Block	Athagad (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	12/04/2024	13/04/2024	14/04/2024	15/04/2024	16/04/2024
Rainfall (mm)	0	0	0	0	0
T-Max (°C)	32	33	36	37	37
T-Min (°C)	24	27	26	25	25
Cloud Cover (octa)	4	2	2	2	2
RH Max (%)	48	49	56	72	81
RH Min (%)	19	20	21	20	20
Wind Speed (kmph)	10	12	10	12	13
Wind Direction (deg)	270	289	200	191	191

Crop specific Agro-met Advisory

General advisory: Farmers should harvest the matured crops and vegetables in dry days. Farmers may go for intercultural operations, fertilizer and pesticide application in the upcoming dry days. They should start sowing of summer vegetables by utilizing past rain water. They should provide mulching material or polythene cover to recently planted vegetables seedling to protect from increase in day temperature.

Pulses:

Harvest matured pulses in dry days. Rust infestation likely to occur in pulses, if observe spray Mancozeb 35.0%SC 3gram/ltr of water. If **wilting** will be observed after rainfall, then spray Carbendazim 2g/ltr of water to control.

Summer Maize:

There may be increase the infestation of **Red hairy caterpillar** in maize after rainfall, to minimize the infestation farmers may spray Neem oil 5ml/ltr of water or Chlorpyrifos 20 % EC @ 2ml/litre of water. Prolonged dry spell followed by rain may increase the infestation of **fall army worm in maize**. To control spray emamectin benzoate (5% S.G.) @ 0.4g per litre of water.

Vegetables:

Okra: To control **Yellow Vain Mosaic Virus (YVMV)** in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease.

Brinjal:

There are chances of infestation of **Epilachna Beetle/Haddabeetle** in Brinjal and cucurbits crop; to manage Epilachna Beetle in cucurbits, spray neem-based pesticide (Azadirachtin) 1500 PPM @ 3ml/litre of water at early stage of infestation; to manage Epilachna Beetle chemically, spray Chlorpyrifos 20 % EC @ 2ml/litre of water or Profenophos 50 % EC @ 2ml/litre of water.

FRUIT AND SHOOT BORER IN BRINJAL-Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@250g/acre at 15 days interval. If **phomopsis blight** will occur in brinjal, then spray zineb75%WP 2gram/litre of water. **Cercospora leaf spot** may be found after rainfall, if observe, then spray Carbendazim or zineb75%WP 2gram/litre of water. If **helicoverpa infestation** will be found, spray Flunendiamide or indoxacarb 2ml/litre of water.

Cucurbits:

To manage **Red Pumpkin Beetle** in Cucurbit crops spray Cyantraniliprole 10.26% OD @ 400-ml/acre or Imidacloprid 17.8% SL @ 60-ml/ acre or Chlorpyrifos 20% EC @ 400-ml/acre. **Late Blight** may happen due to rising day temperature. There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Don't go for weeding. The grasses and weeds will help in maintaining the humidity and it is safe for creeping of the vines and does not allow the fruit to come in contact with soil and avoid fruit rot. High temperature leads to fungal wilting in pointed gourd, spray 1.5 ml Carbendazim in 1 litre water. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxchloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Mango:

Water stress immediately after fruit set increases fruit dropping in Mango. Hence, protective irrigation is essential during the fruit development stage. To control **fruit drop**, spray Borax @2gm per lit. of water and apply Planofix @2ml per 5 lit. of water. For control of mango **fruit borer** spray Neem oil 3ml + chlorpyrifos 1 ml per litre of water along with stickers at marble stage of the fruit. The sprayings should be preferably carried out during evening hours. To control **bacterial canker**, three sprays of streptocycline (100 ppm) or Agrimycin-100 (100 ppm) after first visual symptom at 10- days intervals is recommended.

Animal husbandry:

Make necessary arrangement for clean and sufficient drinking water for the animals.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-485
Date- 12/04/2024

AGROMET ADVISORY BULLETIN FOR BADAMBA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 3 mm and district received average 2mm rainfall during the last week. The mean daily maximum and minimum temperature was 35.47°C (-1.95 from normal) and 23.37°C (-0.3 from normal), respectively. The mean relative humidity was 61-67%.

Forecasted period: Today(8.30IST) to 17th April (8.30IST)2024

Weather advisory: Dry weather is forecasted for today to Tuesday in the block. The day temperature will be 32-38°C and night temperature will be 23-25°C for upcoming 5 days. Maximum and minimum relative humidity will be 47-71% and 19-21% respectively. There will be clear sky for next 5 days.

Block	Badamba (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	12/04/2024	13/04/2024	14/04/2024	15/04/2024	16/04/2024
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	32	34	36	38	38
T-Min(°C)	23	25	25	24	25
Cloud Cover(octa)	4	3	2	3	4
RH Max (%)	47	47	54	65	71
RH Min (%)	20	20	21	20	19
Wind Speed (Kmph)	9	11	9	11	10
Wind Direction (deg)	270	207	195	199	171

Crop specific Agro-met Advisory

General advisory: Farmers should harvest the matured crops and vegetables in dry days. Farmers may go for intercultural operations, fertilizer and pesticide application in the upcoming dry days. They should start sowing of summer vegetables by utilizing past rain water. They should provide mulching material or polythene cover to recently planted vegetables seedling to protect from increase in day temperature.

Pulses:

Harvest matured pulses in dry days. Rust infestation likely to occur in pulses, if observe spray Mancozeb 35.0%SC 3gram/ltr of water. If wilting will be observed after rainfall, then spray Carbendazim 2g/ltr of water to control.

Summer Maize:

There may be increase the infestation of Red hairy caterpillar in maize after rainfall, to minimize the infestation farmers may spray Neem oil 5ml/ltr of water or Chlorpyrifos 20 % EC @ 2ml/litre of water. Prolonged dry spell followed by rain may increase the infestation of fall army worm in maize. To control spray emamectin benzoate (5% S.G.) @ 0.4g per litre of water.

Vegetables:

Okra: To control Yellow Vain Mosaic Virus (YVMV) in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease.

Brinjal:

There are chances of infestation of Epilachna Beetle/Haddabeetle in Brinjal and cucurbits crop; to manage Epilachna Beetle in cucurbits, spray neem-based pesticide (Azadirachtin) 1500 PPM @ 3ml/litre of water at early stage of infestation; to manage Epilachna Beetle chemically, spray Chlorpyrifos 20 % EC @ 2ml/litre of water or Profenophos 50 % EC @ 2ml/litre of water.

FRUIT AND SHOOT BORER IN BRINJAL: Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@250g/acre at 15 days interval. If phomopsis blight will occur in brinjal, then spray zineb75%WP 2gram/litr of water. Cercopspera leaf spot may be found after rainfall, if observe, then spray Carbendazim or zineb75%WP 2gram/litr of water. If helicoverpa infestation will be found, spray Flunendiamide or indoxacarb 2ml/litr of water.

Cucurbits:

To manage Red Pumpkin Beetle in Cucurbit crops spray Cyantraniliprole 10.26% OD @ 400-ml/acre or Imidacloprid 17.8% SL @ 60-ml/ acre or Chlorpyrifos 20% EC @ 400-ml/acre. Late Blight may happen due to rising day temperature. There are chances of infestation of Downy Mildew in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Don't go for weeding. The grasses and weeds will help in maintaining the humidity and it is safe for creeping of the vines and does not allow the fruit to come in contact with soil and avoid fruit rot. High temperature leads to fungal wilting in pointed gourd, spray 1.5 ml Carbendazim in 1 litre water. Present weather may aggravate fruit and vine rot (symptoms: leaf spot, leaf blight, stem rot, fruit rot) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxychloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Mango:

Water stress immediately after fruit set increases fruit dropping in Mango. Hence, protective irrigation is essential during the fruit development stage. To control fruit drop, spray Borax @2gm per lit. of water and apply Planofix @2ml per 5 lit. of water. For control of mango fruit borer spray Neem oil 3ml + chlorpyrifos 1 ml per litre of water along with stickers at marble stage of the fruit. The sprayings should be preferably carried out during evening hours. To control bacterial canker, three sprays of streptomycin (100 ppm) or Agrimycin-100 (100 ppm) after first visual symptom at 10- days intervals is recommended.

Animal husbandry:

Make necessary arrangement for clean and sufficient drinking water for the animals.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-485
Date- 12/04/2024

AGROMET ADVISORY BULLETIN FOR BANKI

District Past week average weather condition

The district received average 2mm rainfall during the last week. The mean daily maximum and minimum temperature was 35.47°C (-1.95 from normal) and 23.37°C (-0.3 from normal), respectively. The mean relative humidity was 61-67%.

Forecasted period: Today(8.30IST) to 17thApril (8.30IST)2024

Weather advisory; Dry weather is forecasted for today to Tuesday in the block. The day temperature will be 33-37°C and night temperature will be 23-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 49-77% and 20-21% respectively. There will be clear sky for next 5days.

Block	Banki(VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	12/04/2024	13/04/2024	14/04/2024	15/04/2024	16/04/2024
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	33	33	36	36	37
T-Min(°C)	23	26	26	25	25
Cloud Cover(octa)	4	2	2	3	3
RH Max (%)	49	49	55	70	77
RH Min (%)	20	20	21	21	21
Wind Speed (Kmph)	10	12	11	13	13
Wind Direction (deg)	270	214	207	203	191

Crop specific Agro-met Advisory

General advisory: Farmers should harvest the matured crops and vegetables in dry days. Farmers may go for intercultural operations, fertilizer and pesticide application in the upcoming dry days. They should start sowing of summer vegetables by utilizing past rain water. They should provide mulching material or polythene cover to recently planted vegetables seedling to protect from increase in day temperature.

Pulses:

Harvest matured pulses in dry days. Rust infestation likely to occur in pulses, if observe spray Mancozeb 35.0%SC 3gram/ltr of water. If **wilting** will be observed after rainfall, then spray Carbendazim 2g/ltr of water to control.

Summer Maize:

There may be increase the infestation of **Red hairy caterpillar** in maize after rainfall, to minimize the infestation farmers may spray Neem oil 5ml/ltr of water or Chlorpyrifos 20 % EC @ 2ml/litre of water. Prolonged dry spell followed by rain may increase the infestation of **fall army worm in maize**. To control spray emamectin benzoate (5% S.G.) @ 0.4g per litre of water.

Vegetables:

Okra: To control **Yellow Vain Mosaic Virus (YVMV)** in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease.

Brinjal:

There are chances of infestation of **Epilachna Beetle/Haddabeetle** in Brinjal and cucurbits crop; to manage Epilachna Beetle in cucurbits, spray neem-based pesticide (Azadirachtin) 1500 PPM @ 3ml/litre of water at early stage of infestation; to manage Epilachna Beetle chemically, spray Chlorpyrifos 20 % EC @ 2ml/litre of water or Profenophos 50 % EC @ 2ml/litre of water.

FRUIT AND SHOOT BORER IN BRINJAL-Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@250g/acre at 15 days interval. If **phomopsis blight** will occur in brinjal, then spray zineb75%WP 2gram/litr of water.**Cercopspera leaf spot** may be found after rainfall, if observe, then spray Carbendazim or zineb75%WP 2gram/litr of water.If **helicoverpa infestation** will be found, spray Flunendiamide or indoxacarb 2ml/litr of water.

Cucurbits:

To manage **Red Pumpkin Beetle** in Cucurbit crops spray Cyantraniliprole 10.26% OD @ 400-ml/acre or Imidacloprid 17.8% SL @ 60-ml/ acre or Chlorpyrifos 20% EC @ 400-ml/acre. **Late Blight** may happen due to rising day temperature. There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Don't go for weeding .The grasses and weeds will help in maintaining the humidity and it is safe for creeping of the vines and does not allow the fruit to come in contact with soil and avoid fruit rot. High temperature leads to fungal wilting in pointed gourd, spray 1.5 ml Carbendazim in 1 litre water. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxchloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Mango:

Water stress immediately after fruit set increases fruit dropping in Mango. Hence, protective irrigation is essential during the fruit development stage. To control **fruit drop**, spray Borax @2gm per lit. of water and apply Planofix @2ml per 5 lit. of water.For control of mango **fruit borer** spray Neem oil 3ml + chlorpyrifos 1 ml perlitre of water along with stickers at marble stage of the fruit. The sprayingshould be preferably carried out during evening hours. To control **bacterialcanker**, three sprays of streptomycine (100 ppm) or Agrimycin-100 (100 ppm)after first visual symptom at 10- days intervals is recommended.

Animal husbandry:

Make necessary arrangement for clean and sufficient drinking water for the animals.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-485
Date- 12/04/2024

AGROMET ADVISORY BULLETIN FOR BANKI-DAMPADA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 1 mm and district received average 2mm rainfall during the last week. The mean daily maximum and minimum temperature was 35.47°C (-1.95 from normal) and 23.37°C (-0.3 from normal), respectively. The mean relative humidity was 61-67%.

Forecasted period: Today(8.30IST) to 17thApril (8.30IST)2024

Weather Advisory: Dry weather is forecasted for today to Tuesday in the block. The day temperature will be 33-37°C and night temperature will be 24-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 49-78% and 20-21% respectively. There will be clear sky for next 5days.

Block	Banki-Dampada (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	12/04/2024	13/04/2024	14/04/2024	15/04/2024	16/04/2024
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	33	33	36	36	37
T-Min(°C)	24	26	26	25	25
Cloud Cover(octa)	4	2	2	2	3
RH Max (%)	49	50	56	70	78
RH Min (%)	20	20	21	21	21
Wind Speed (Kmph)	10	12	11	14	13
Wind Direction (deg)	270	214	207	203	199

Crop specific Agro-met Advisory

General advisory: Farmers should harvest the matured crops and vegetables in dry days. Farmers may go for intercultural operations, fertilizer and pesticide application in the upcoming dry days. They should start sowing of summer vegetables by utilizing past rain water. They should provide mulching material or polythene cover to recently planted vegetables seedling to protect from increase in day temperature.

Pulses:

Harvest matured pulses in dry days. Rust infestation likely to occur in pulses, if observe spray Mancozeb 35.0%SC 3gram/ltr of water. If wilting will be observed after rainfall, then spray Carbendazim 2g/ltr of water to control.

Summer Maize:

There may be increase the infestation of Red hairy caterpillar in maize after rainfall, to minimize the infestation farmers may spray Neem oil 5ml/ltr of water or Chlorpyrifos 20 % EC @ 2ml/litre of water. Prolonged dry spell followed by rain may increase the infestation of fall army worm in maize. To control spray emamectin benzoate (5% S.G.) @ 0.4g per litre of water.

Vegetables:

Okra: To control Yellow Vain Mosaic Virus (YVMV) in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease.

Brinjal:

There are chances of infestation of Epilachna Beetle/Haddabeetle in Brinjal and cucurbits crop; to manage Epilachna Beetle in cucurbits, spray neem-based pesticide (Azadirachtin) 1500 PPM @ 3ml/litre of water at early stage of infestation; to manage Epilachna Beetle chemically, spray Chlorpyrifos 20 % EC @ 2ml/litre of water or Profenophos 50 % EC @ 2ml/litre of water.

FRUIT AND SHOOT BORER IN BRINJAL-Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@250g/acre at 15 days interval. If phomopsis blight will occur in brinjal, then spray zineb75%WP 2gram/litr of water. Cercopspera leaf spot may be found after rainfall, if observe, then spray Carbendazim or zineb75%WP 2gram/litr of water. If helicoverpa infestation will be found, spray Flunendiamide or indoxacarb 2ml/litr of water.

Cucurbits:

To manage Red Pumpkin Beetle in Cucurbit crops spray Cyantraniliprole 10.26% OD @ 400-ml/acre or Imidacloprid 17.8% SL @ 60-ml/ acre or Chlorpyrifos 20% EC @ 400-ml/acre. Late Blight may happen due to rising day temperature. There are chances of infestation of Downy Mildew in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Don't go for weeding .The grasses and weeds will help in maintaining the humidity and it is safe for creeping of the vines and does not allow the fruit to come in contact with soil and avoid fruit rot. High temperature leads to fungal wilting in pointed gourd, spray 1.5 ml Carbendazim in 1 litre water. Present weather may aggravate fruit and vine rot (symptoms: leaf spot, leaf blight, stem rot, fruit rot) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxychloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Mango:

Water stress immediately after fruit set increases fruit dropping in Mango. Hence, protective irrigation is essential during the fruit development stage. To control fruit drop, spray Borax @2gm per lit. of water and apply Planofix @2ml per 5 lit. of water. For control of mango fruit borer spray Neem oil 3ml + chlorpyrifos 1 ml per litre of water along with stickers at marble stage of the fruit. The sprayings should be preferably carried out during evening hours. To control bacterial canker, three sprays of streptomycin (100 ppm) or Agrimycin-100 (100 ppm) after first visual symptom at 10- days intervals is recommended.

Animal husbandry:

Make necessary arrangement for clean and sufficient drinking water for the animals.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-485
Date- 12/04/2024

AGROMET ADVISORY BULLETIN FOR BARANGA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 1 mm and district received average 2mm rainfall during the last week. The mean daily maximum and minimum temperature was 35.47°C (-1.95 from normal) and 23.37°C (-0.3 from normal), respectively. The mean relative humidity was 61-67%.

Forecasted period: Today(8.30IST) to 17thApril (8.30IST)2024

Weatheradvisory: *Dry weather is forecasted for today to Tuesday in the block.* The day temperature will be 33-36°C and night temperature will be 24-27°C for upcoming 5 days. Maximum and minimum relative humidity will be 53-83% and 18-22% respectively. There will be clear sky for next 5days.

Block	Baranga (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	12/04/2024	13/04/2024	14/04/2024	15/04/2024	16/04/2024
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	33	33	36	36	36
T-Min(°C)	24	27	27	25	26
Cloud Cover	3	2	2	2	2
RH Max (%)	53	57	60	77	83
RH Min (%)	18	20	20	21	22
Wind Speed (Kmph)	12	14	13	16	19
Wind Direction (deg)	270	210	208	203	203

Crop specific Agro-met Advisory

General advisory: *Farmers should harvest the matured crops and vegetables in dry days. Farmers may go for intercultural operations, fertilizer and pesticide application in the upcoming dry days. They should start sowing of summer vegetables by utilizing past rain water. They should provide mulching material or polythene cover to recently planted vegetables seedling to protect from increase in day temperature.*

Pulses:

Harvest matured pulses in dry days. Rust infestation likely to occur in pulses, if observe spray Mancozeb 35.0%SC 3gram/ltr of water. If **wilting** will be observed after rainfall, then spray Carbendazim 2g/ltr of water to control.

Summer Maize:

There may be increase the infestation of **Red hairy caterpillar** in maize after rainfall, to minimize the infestation farmers may spray Neem oil 5ml/ltr of water or Chlorpyrifos 20 % EC @ 2ml/litre of water. Prolonged dry spell followed by rain may increase the infestation of **fall army worm in maize**. To control spray emamectin benzoate (5% S.G.) @ 0.4g per litre of water.

Vegetables:

Okra: To control **Yellow Vain Mosaic Virus (YVMV)** in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease.

Brinjal:

There are chances of infestation of **Epilachna Beetle/Haddabeetle** in Brinjal and cucurbits crop; to manage Epilachna Beetle in cucurbits, spray neem-based pesticide (Azadirachtin) 1500 PPM @ 3ml/litre of water at early stage of infestation; to manage Epilachna Beetle chemically, spray Chlorpyrifos 20 % EC @ 2ml/litre of water or Profenophos 50 % EC @ 2ml/litre of water.

FRUIT AND SHOOT BORER IN BRINJAL-Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@250g/acre at 15 days interval. If **phomopsis blight** will occur in brinjal, then spray zineb75%WP 2gram/litr of water. **Cercopspera leaf spot** may be found after rainfall, if observe, then spray Carbendazim or zineb75%WP 2gram/litr of water. If **helicoverpa infestation** will be found, spray Flunendiamide or indoxacarb 2ml/litr of water.

Cucurbits:

To manage **Red Pumpkin Beetle** in Cucurbit crops spray Cyantraniliprole 10.26% OD @ 400-ml/acre or Imidacloprid 17.8% SL @ 60-ml/ acre or Chlorpyrifos 20% EC @ 400-ml/acre. **Late Blight** may happen due to rising day temperature. There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Don't go for weeding .The grasses and weeds will help in maintaining the humidity and it is safe for creeping of the vines and does not allow the fruit to come in contact with soil and avoid fruit rot. High temperature leads to fungal wilting in pointed gourd, spray 1.5 ml Carbendazim in 1 litre water. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxychloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Mango:

Water stress immediately after fruit set increases fruit dropping in Mango. Hence, protective irrigation is essential during the fruit development stage. To control **fruit drop**, spray Borax @2gm per lit. of water and apply Planofix @2ml per 5 lit. of water. For control of mango **fruit borer** spray Neem oil 3ml + chlorpyrifos 1 ml per litre of water along with stickers at marble stage of the fruit. The sprayings should be preferably carried out during evening hours. To control **bacterialcanker**, three sprays of streptocycline (100 ppm) or Agrimycin-100 (100 ppm) after first visual symptom at 10- days intervals is recommended.

Animal husbandry:

Make necessary arrangement for clean and sufficient drinking water for the animals.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-485
Date- 12/04/2024

AGROMET ADVISORY BULLETIN FOR CUTTACK SADAR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 2 mm and district received average 2mm rainfall during the last week. The mean daily maximum and minimum temperature was 35.47°C (-1.95 from normal) and 23.37°C (-0.3 from normal), respectively. The mean relative humidity was 61-67%.

Forecasted period: Today(8.30IST) to 17thApril (8.30IST)2024

Weather advisory: *Dry weather is forecasted for today to Tuesday in the block.* The day temperature will be 33-37°C and night temperature will be 24-27°C for upcoming 5 days. Maximum and minimum relative humidity will be 50-84% and 18-21% respectively. There will be clear sky for next 5days.

Block	Cuttack sadar (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	12/04/2024	13/04/2024	14/04/2024	15/04/2024	16/04/2024
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	33	33	36	37	36
T-Min(°C)	24	27	27	25	26
Cloud Cover(octa)	4	2	2	2	2
RH Max (%)	50	54	59	77	84
RH Min (%)	18	19	20	20	21
Wind Speed (Kmph)	11	13	12	15	16
Wind Direction (deg)	270	210	205	204	199

Crop specific Agro-met Advisory

General advisory: *Farmers should harvest the matured crops and vegetables in dry days. Farmers may go for intercultural operations, fertilizer and pesticide application in the upcoming dry days. They should start sowing of summer vegetables by utilizing past rain water. They should provide mulching material or polythene cover to recently planted vegetables seedling to protect from increase in day temperature.*

Pulses:

Harvest matured pulses in dry days. Rust infestation likely to occur in pulses, if observe spray Mancozeb 35.0%SC 3gram/ltr of water. If **wilting** will be observed after rainfall, then spray Carbendazim 2g/ltr of water to control.

Summer Maize:

There may be increase the infestation of **Red hairy caterpillar** in maize after rainfall, to minimize the infestation farmers may spray Neem oil 5ml/ltr of water or Chlorpyrifos 20 % EC @ 2ml/litre of water. Prolonged dry spell followed by rain may increase the infestation of **fall army worm in maize**. To control spray emamectin benzoate (5% S.G.) @ 0.4g per litre of water.

Vegetables:

Okra: To control **Yellow Vain Mosaic Virus (YVMV)** in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease.

Brinjal:

There are chances of infestation of **Epilachna Beetle/Haddabeetle** in Brinjal and cucurbits crop; to manage Epilachna Beetle in cucurbits, spray neem-based pesticide (Azadirachtin) 1500 PPM @ 3ml/litre of water at early stage of infestation; to manage Epilachna Beetle chemically, spray Chlorpyrifos 20 % EC @ 2ml/litre of water or Profenophos 50 % EC @ 2ml/litre of water.

FRUIT AND SHOOT BORER IN BRINJAL-Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@250g/acre at 15 days interval. If **phomopsis blight** will occur in brinjal, then spray zineb75%WP 2gram/litr of water. **Cercopspera leaf spot** may be found after rainfall, if observe, then spray Carbendazim or zineb75%WP 2gram/litr of water. If **helicoverpa infestation** will be found, spray Flunendiamide or indoxacarb 2ml/litr of water.

Cucurbits:

To manage **Red Pumpkin Beetle** in Cucurbit crops spray Cyantraniliprole 10.26% OD @ 400-ml/acre or Imidacloprid 17.8% SL @ 60-ml/ acre or Chlorpyrifos 20% EC @ 400-ml/acre. **Late Blight** may happen due to rising day temperature. There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Don't go for weeding .The grasses and weeds will help in maintaining the humidity and it is safe for creeping of the vines and does not allow the fruit to come in contact with soil and avoid fruit rot. High temperature leads to fungal wilting in pointed gourd, spray 1.5 ml Carbendazim in 1 litre water. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxychloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Mango:

Water stress immediately after fruit set increases fruit dropping in Mango. Hence, protective irrigation is essential during the fruit development stage. To control **fruit drop**, spray Borax @2gm per lit. of water and apply Planofix @2ml per 5 lit. of water. For control of mango **fruit borer** spray Neem oil 3ml + chlorpyrifos 1 ml per litre of water along with stickers at marble stage of the fruit. The sprayings should be preferably carried out during evening hours. To control **bacterialcanker**, three sprays of streptocycline (100 ppm) or Agrimycin-100 (100 ppm) after first visual symptom at 10- days intervals is recommended.

Animal husbandry:

Make necessary arrangement for clean and sufficient drinking water for the animals.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SAKSHI
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-485
Date- 12/04/2024

AGROMET ADVISORY BULLETIN FOR KANTAPADA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 3 mm and district received average 2mm rainfall during the last week. The mean daily maximum and minimum temperature was 35.47°C (-1.95 from normal) and 23.37°C (-0.3 from normal), respectively. The mean relative humidity was 61-67%.

Forecasted period: Today(8.30IST) to 17th April (8.30IST)2024

Weather advisory: Dry weather is forecasted for today to Tuesday in the block. The day temperature will be 34-37°C and night temperature will be 24-27°C for upcoming 5 days. Maximum and minimum relative humidity will be 58-83% and 18-25% respectively. There will be clear sky for next 5 days.

Block	Kantapada (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	12/04/2024	13/04/2024	14/04/2024	15/04/2024	16/04/2024
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	34	34	37	37	36
T-Min(°C)	24	26	27	26	27
Cloud Cover(octa)	3	2	1	2	1
RH Max (%)	58	64	64	78	83
RH Min (%)	18	21	21	22	25
Wind Speed (Kmph)	12	15	15	18	22
Wind Direction (deg)	248	208	205	203	203

Crop specific Agro-met Advisory

General advisory: Farmers should harvest the matured crops and vegetables in dry days. Farmers may go for intercultural operations, fertilizer and pesticide application in the upcoming dry days. They should start sowing of summer vegetables by utilizing past rain water. They should provide mulching material or polythene cover to recently planted vegetables seedling to protect from increase in day temperature.

Pulses:

Harvest matured pulses in dry days. Rust infestation likely to occur in pulses, if observe spray Mancozeb 35.0%SC 3gram/ltr of water. If **wilting** will be observed after rainfall, then spray Carbendazim 2g/ltr of water to control.

Summer Maize:

There may be increase the infestation of **Red hairy caterpillar** in maize after rainfall, to minimize the infestation farmers may spray Neem oil 5ml/ltr of water or Chlorpyrifos 20 % EC @ 2ml/litre of water. Prolonged dry spell followed by rain may increase the infestation of **fall army worm in maize**. To control spray emamectin benzoate (5% S.G.) @ 0.4g per litre of water.

Vegetables:

Okra: To control **Yellow Vain Mosaic Virus (YVMV)** in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease.

Brinjal:

There are chances of infestation of **Epilachna Beetle/Haddabeetle** in Brinjal and cucurbits crop; to manage Epilachna Beetle in cucurbits, spray neem-based pesticide (Azadirachtin) 1500 PPM @ 3ml/litre of water at early stage of infestation; to manage Epilachna Beetle chemically, spray Chlorpyrifos 20 % EC @ 2ml/litre of water or Profenophos 50 % EC @ 2ml/litre of water.

FRUIT AND SHOOT BORER IN BRINJAL-Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@250g/acre at 15 days interval. If **phomopsis blight** will occur in brinjal, then spray zineb75%WP 2gram/litr of water. **Cercopspera leaf spot** may be found after rainfall, if observe, then spray Carbendazim or zineb75%WP 2gram/litr of water. If **helicopterpa infestation** will be found, spray Flunendiamide or indoxacarb 2ml/litr of water.

Cucurbits:

To manage **Red Pumpkin Beetle** in Cucurbit crops spray Cyantraniliprole 10.26% OD @ 400-ml/acre or Imidacloprid 17.8% SL @ 60-ml/ acre or Chlorpyrifos 20% EC @ 400-ml/acre. **Late Blight** may happen due to rising day temperature. There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Don't go for weeding. The grasses and weeds will help in maintaining the humidity and it is safe for creeping of the vines and does not allow the fruit to come in contact with soil and avoid fruit rot. High temperature leads to fungal wilting in pointed gourd, spray 1.5 ml Carbendazim in 1 litre water. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxychloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Mango:

Water stress immediately after fruit set increases fruit dropping in Mango. Hence, protective irrigation is essential during the fruit development stage. To control **fruit drop**, spray Borax @2gm per lit. of water and apply Planofix @2ml per 5 lit. of water. For control of mango **fruit borer** spray Neem oil 3ml + chlorpyrifos 1 ml per litre of water along with stickers at marble stage of the fruit. The spraying should be preferably carried out during evening hours. To control **bacterial canker**, three sprays of streptomycin (100 ppm) or Agrimycin-100 (100 ppm) after first visual symptom at 10- days intervals is recommended.

Animal husbandry:

Make necessary arrangement for clean and sufficient drinking water for the animals.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-485
Date- 12/04/2024

AGROMET ADVISORY BULLETIN FOR MAHANGA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 4 mm and district received average 2mm rainfall during the last week. The mean daily maximum and minimum temperature was 35.47°C (-1.95 from normal) and 23.37°C (-0.3 from normal), respectively. The mean relative humidity was 61-67%.

Forecasted period: Today(8.30IST) to 17th April (8.30IST)2024

Weather advisory: Dry weather is forecasted for today to Tuesday in the block. The day temperature will be and it will be 34-37°C and night temperature will be 24-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 50-83% and 19-23% respectively. There will be clear sky for next 5days.

Block	Mahanga (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	12/04/2024	13/04/2024	14/04/2024	15/04/2024	16/04/2024
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	34	34	37	37	37
T-Min(°C)	24	26	26	25	26
Cloud Cover(octal)	4	3	2	3	2
RH Max (%)	50	59	61	76	83
RH Min (%)	19	19	21	21	23
Wind Speed (Kmph)	13	14	14	16	19
Wind Direction (deg)	270	210	210	204	203

Crop specific Agro-met Advisory

General advisory: Farmers should harvest the matured crops and vegetables in dry days. Farmers may go for intercultural operations, fertilizer and pesticide application in the upcoming dry days. They should start sowing of summer vegetables by utilizing past rain water. They should provide mulching material or polythene cover to recently planted vegetables seedling to protect from increase in day temperature.

Pulses:

Harvest matured pulses in dry days. Rust infestation likely to occur in pulses, if observe spray Mancozeb 35.0%SC 3gram/ltr of water. If **wilting** will be observed after rainfall, then spray Carbendazim 2g/ltr of water to control.

Summer Maize:

There may be increase the infestation of **Red hairy caterpillar** in maize after rainfall, to minimize the infestation farmers may spray Neem oil 5ml/ltr of water or Chlorpyrifos 20 % EC @ 2ml/litre of water. Prolonged dry spell followed by rain may increase the infestation of **fall army worm in maize**. To control spray emamectin benzoate (5% S.G.) @ 0.4g per litre of water.

Vegetables:

Okra: To control **Yellow Vain Mosaic Virus (YVMV)** in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease.

Brinjal:

There are chances of infestation of **Epilachna Beetle/Haddabeetle** in Brinjal and cucurbits crop; to manage Epilachna Beetle in cucurbits, spray neem-based pesticide (Azadirachtin) 1500 PPM @ 3ml/litre of water at early stage of infestation; to manage Epilachna Beetle chemically, spray Chlorpyrifos 20 % EC @ 2ml/litre of water or Profenophos 50 % EC @ 2ml/litre of water.

FRUIT AND SHOOT BORER IN BRINJAL-Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@250g/acre at 15 days interval. If **phomopsis blight** will occur in brinjal, then spray zineb75%WP 2gram/litr of water. **Cercopspera leaf spot** may be found after rainfall, if observe, then spray Carbendazim or zineb75%WP 2gram/litr of water. If **helicoverpa infestation** will be found, spray Flunendiamide or indoxacarb 2ml/litr of water.

Cucurbits:

To manage **Red Pumpkin Beetle** in Cucurbit crops spray Cyantraniliprole 10.26% OD @ 400-ml/acre or Imidacloprid 17.8% SL @ 60-ml/ acre or Chlorpyrifos 20% EC @ 400-ml/acre. **Late Blight** may happen due to rising day temperature. There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Don't go for weeding. The grasses and weeds will help in maintaining the humidity and it is safe for creeping of the vines and does not allow the fruit to come in contact with soil and avoid fruit rot. High temperature leads to fungal wilting in pointed gourd, spray 1.5 ml Carbendazim in 1 litre water. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxychloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Mango:

Water stress immediately after fruit set increases fruit dropping in Mango. Hence, protective irrigation is essential during the fruit development stage. To control **fruit drop**, spray Borax @2gm per lit. of water and apply Planofix @2ml per 5 lit. of water. For control of mango **fruit borer** spray Neem oil 3ml + chlorpyrifos 1 ml per litre of water along with stickers at marble stage of the fruit. The spraying should be preferably carried out during evening hours. To control **bacterial canker**, three sprays of streptomycin (100 ppm) or Agrimycin-100 (100 ppm) after first visual symptom at 10- days intervals is recommended.

Animal husbandry:

Make necessary arrangement for clean and sufficient drinking water for the animals.

Dr Sujata Sethy
 (Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
 Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SAKSHI
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-485
Date- 12/04/2024

AGROMET ADVISORY BULLETIN FOR NARASINGHPUR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 3 mm and district received average 2mm rainfall during the last week. The mean daily maximum and minimum temperature was 35.47°C (-1.95 from normal) and 23.37°C (-0.3 from normal), respectively. The mean relative humidity was 61-67%.

Forecasted period: Today(8.30IST) to 17thApril (8.30IST)2024

Weather advisory: Dry weather is forecasted for today to Tuesday in the block. The day temperature will be 32-42°C and night temperature will be 21-24°C for upcoming 5 days. Maximum and minimum relative humidity will be 46-61% and 17-21% respectively. There will be clear sky for next 5 days.

Block	Narasinghpur (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	12/04/2024	13/04/2024	14/04/2024	15/04/2024	16/04/2024
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	32	35	38	40	42
T-Min(°C)	21	23	24	24	24
Cloud Cover(octa)	4	3	2	4	4
RH Max (%)	46	47	53	58	61
RH Min (%)	21	19	21	19	17
Wind Speed (Kmph)	7	8	6	7	7
Wind Direction (deg)	225	296	205	199	180

Crop specific Agro-met Advisory

General advisory: Farmers should harvest the matured crops and vegetables in dry days. Farmers may go for intercultural operations, fertilizer and pesticide application in the upcoming dry days. They should start sowing of summer vegetables by utilizing past rain water. They should provide mulching material or polythene cover to recently planted vegetables seedling to protect from increase in day temperature.

Pulses:

Harvest matured pulses in dry days. Rust infestation likely to occur in pulses, if observe spray Mancozeb 35.0%SC 3gram/ltr of water. If **wilting** will be observed after rainfall, then spray Carbendazim 2g/ltr of water to control.

Summer Maize:

There may be increase the infestation of **Red hairy caterpillar** in maize after rainfall, to minimize the infestation farmers may spray Neem oil 5ml/ltr of water or Chlorpyrifos 20 % EC @ 2ml/litre of water. Prolonged dry spell followed by rain may increase the infestation of **fall army worm in maize**. To control spray emamectin benzoate (5% S.G.) @ 0.4g per litre of water.

Vegetables:

Okra: To control **Yellow Vain Mosaic Virus (YVMV)** in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease.

Brinjal:

There are chances of infestation of **Epilachna Beetle/Haddabeetle** in Brinjal and cucurbits crop; to manage Epilachna Beetle in cucurbits, spray neem-based pesticide (Azadirachtin) 1500 PPM @ 3ml/litre of water at early stage of infestation; to manage Epilachna Beetle chemically, spray Chlorpyrifos 20 % EC @ 2ml/litre of water or Profenophos 50 % EC @ 2ml/litre of water.

FRUIT AND SHOOT BORER IN BRINJAL-Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@250g/acre at 15 days interval. If **phomopsis blight** will occur in brinjal, then spray zineb75%WP 2gram/litr of water. **Cercopspera leaf spot** may be found after rainfall, if observe, then spray Carbendazim or zineb75%WP 2gram/litr of water. If **helicoverta infestation** will be found, spray Flunendiamide or indoxacarb 2ml/litr of water.

Cucurbits:

To manage **Red Pumpkin Beetle** in Cucurbit crops spray Cyantraniliprole 10.26% OD @ 400-ml/acre or Imidacloprid 17.8% SL @ 60-ml/ acre or Chlorpyrifos 20% EC @ 400-ml/acre. **Late Blight** may happen due to rising day temperature. There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Don't go for weeding .The grasses and weeds will help in maintaining the humidity and it is safe for creeping of the vines and does not allow the fruit to come in contact with soil and avoid fruit rot. High temperature leads to fungal wilting in pointed gourd, spray 1.5 ml Carbendazim in 1 litre water. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxychloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Mango:

Water stress immediately after fruit set increases fruit dropping in Mango. Hence, protective irrigation is essential during the fruit development stage. To control **fruit drop**, spray Borax @2gm per lit. of water and apply Planofix @2ml per 5 lit. of water. For control of mango **fruit borer** spray Neem oil 3ml + chlorpyrifos 1 ml per litre of water along with stickers at marble stage of the fruit. The sprayings should be preferably carried out during evening hours. To control **bacterialcanker**, three sprays of streptocycline (100 ppm) or Agrimycin-100 (100 ppm) after first visual symptom at 10- days intervals is recommended.

Animal husbandry:

Make necessary arrangement for clean and sufficient drinking water for the animals.

Dr Sujata Sethy
 (Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-485
Date- 12/04/2024

AGROMET ADVISORY BULLETIN FOR NIALI

District Past week average weather condition and Block past week average Rainfall condition

The block received average 3 mm and district received average 2mm rainfall during the last week. The mean daily maximum and minimum temperature was 35.47°C (-1.95 from normal) and 23.37°C (-0.3 from normal), respectively. The mean relative humidity was 61-67%.

Forecasted period: Today(8.30IST) to 17thApril (8.30IST)2024

Weather advisory: *Dry weather is forecasted for today to Tuesday in the block.* The day temperature will be 35-37°C and night temperature will be 24-27°C for upcoming 5 days. Maximum and minimum relative humidity will be 62-83% and 19-29% respectively. **There will be clear sky for next 5 days.**

Block	Niali (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	12/04/2024	13/04/2024	14/04/2024	15/04/2024	16/04/2024
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	35	35	37	37	36
T-Min(°C)	24	26	27	26	27
Cloud Cover(octa)	2	3	1	1	1
RH Max (%)	62	72	68	78	83
RH Min (%)	19	22	22	24	29
Wind Speed(Kmph)	10	15	16	20	23
Wind Direction(deg)	244	208	205	203	203

Crop specific Agro-met Advisory

General advisory: *Farmers should harvest the matured crops and vegetables in dry days. Farmers may go for inter-cultural operations, fertilizer and pesticide application in the upcoming dry days. They should start sowing of summer vegetables by utilizing past rain water. They should provide mulching material or polythene cover to recently planted vegetables seedling to protect from increase in day temperature.*

Pulses:

Harvest matured pulses in dry days. Rust infestation likely to occur in pulses, if observe spray Mancozeb 35.0%SC 3gram/ltr of water. If **wilting** will be observed after rainfall, then spray Carbendazim 2g/ltr of water to control.

Summer Maize:

There may be increase the infestation of **Red hairy caterpillar** in maize after rainfall, to minimize the infestation farmers may spray Neem oil 5ml/ltr of water or Chlorpyrifos 20 % EC @ 2ml/litre of water. Prolonged dry spell followed by rain may increase the infestation of **fall army worm in maize**. **To control spray emamectin benzoate (5% S.G.) @ 0.4g per litre of water.**

Vegetables:

Okra: To control **Yellow Vain Mosaic Virus (YVMV)** in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease.

Brinjal:

There are chances of infestation of **Epilachna Beetle/Haddabeetle** in Brinjal and cucurbits crop; to manage Epilachna Beetle in cucurbits, spray neem-based pesticide (Azadirachtin) 1500 PPM @ 3ml/litre of water at early stage of infestation; to manage Epilachna Beetle chemically, spray Chlorpyrifos 20 % EC @ 2ml/litre of water or Profenophos 50 % EC @ 2ml/litre of water.

FRUIT AND SHOOT BORER IN BRINJAL-Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@250g/acre at 15 days interval. If **phomopsis blight** will occur in brinjal, then spray zineb75%WP 2gram/ltr of water. **Cercospora leaf spot** may be found after rainfall, if observe, then spray Carbendazim or zineb75%WP 2gram/ltr of water. If **helicoverpa infestation** will be found, spray Flunendiamide or indoxacarb 2ml/ltr of water.

Cucurbits:

To manage **Red Pumpkin Beetle** in Cucurbit crops spray Cyantraniliprole 10.26% OD @ 400-ml/acre or Imidacloprid 17.8% SL @ 60-ml/ acre or Chlorpyrifos 20% EC @ 400-ml/acre. **Late Blight** may happen due to rising day temperature. There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Don't go for weeding. The grasses and weeds will help in maintaining the humidity and it is safe for creeping of the vines and does not allow the fruit to come in contact with soil and avoid fruit rot. High temperature leads to fungal wilting in pointed gourd, spray 1.5 ml Carbendazim in 1 litre water. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxchloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Mango:

Water stress immediately after fruit set increases fruit dropping in Mango. Hence, protective irrigation is essential during the fruit development stage. To control **fruit drop**, spray Borax @2gm per lit. of water and apply Planofix @2ml per 5 lit. of water. For control of mango **fruit borer** spray Neem oil 3ml + chlorpyrifos 1 ml per litre of water along with stickers at marble stage of the fruit. The sprayings should be preferably carried out during evening hours. To control **bacterial canker**, three sprays of streptomycin (100 ppm) or Agrimycin-100 (100 ppm) after first visual symptom at 10- days intervals is recommended.

Animal husbandry: Make necessary arrangement for clean and sufficient drinking water for the animals.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-485
Date- 12/04/2024

AGROMET ADVISORY BULLETIN FOR NISCHINTAKOILI
District Past week average weather condition and Block past week average Rainfall condition

The block received average 1 mm and district received average 2mm rainfall during the last week. The mean daily maximum and minimum temperature was 35.47°C (-1.95 from normal) and 23.37°C (-0.3 from normal), respectively. The mean relative humidity was 61-67%.

Forecasted period: Today(8.30IST) to 17thApril (8.30IST)2024

Weatheradvisory: *Dry weather is forecasted for today to Tuesday in the block.* The day temperature will be 34-37°C and night temperature will be 24-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 52-83% and 18-23% respectively. There will be clear sky for next 5days.

Block	Nischintakoili (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	12/04/2024	13/04/2024	14/04/2024	15/04/2024	16/04/2024
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	34	34	37	37	37
T-Min(°C)	24	26	26	25	26
Cloud Cover(octa)	4	2	2	3	2
RH Max (%)	52	61	62	77	83
RH Min (%)	18	20	21	21	23
Wind Speed (Kmph)	13	14	14	17	20
Wind Direction (deg)	270	210	207	203	203

Crop specific Agro-met Advisory

General advisory: *Farmers should harvest the matured crops and vegetables in dry days. Farmers may go for intercultural operations, fertilizer and pesticide application in the upcoming dry days. They should start sowing of summer vegetables by utilizing past rain water. They should provide mulching material or polythene cover to recently planted vegetables seedling to protect from increase in day temperature.*

Pulses:

Harvest matured pulses in dry days. Rust infestation likely to occur in pulses, if observe spray Mancozeb 35.0%SC 3gram/ltr of water. If **wilting** will be observed after rainfall, then spray Carbendazim 2g/ltr of water to control.

Summer Maize:

There may be increase the infestation of **Red hairy caterpillar** in maize after rainfall, to minimize the infestation farmers may spray Neem oil 5ml/ltr of water or Chlorpyrifos 20 % EC @ 2ml/litre of water. Prolonged dry spell followed by rain may increase the infestation of **fall army worm in maize**. **To control spray emamectin benzoate (5% S.G.) @ 0.4g per litre of water.**

Vegetables:

Okra: To control **Yellow Vain Mosaic Virus (YVMV)** in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease.

Brinjal:

There are chances of infestation of **Epilachna Beetle/Haddabeetle** in Brinjal and cucurbits crop; to manage Epilachna Beetle in cucurbits, spray neem-based pesticide (Azadirachtin) 1500 PPM @ 3ml/litre of water at early stage of infestation; to manage Epilachna Beetle chemically, spray Chlorpyrifos 20 % EC @ 2ml/litre of water or Profenophos 50 % EC @ 2ml/litre of water.

FRUIT AND SHOOT BORER IN BRINJAL-Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@250g/acre at 15 days interval. If **phomopsis blight** will occur in brinjal, then spray zineb75%WP 2gram/litr of water. **Cercopspera leaf spot** may be found after rainfall, if observe, then spray Carbendazim or zineb75%WP 2gram/litr of water. If **helicoverpa infestation** will be found, spray Flunendiamide or indoxacarb 2ml/litr of water.

Cucurbits:

To manage **Red Pumpkin Beetle** in Cucurbit crops spray Cyantraniliprole 10.26% OD @ 400-ml/acre or Imidacloprid 17.8% SL @ 60-ml/ acre or Chlorpyrifos 20% EC @ 400-ml/acre. **Late Blight** may happen due to rising day temperature. There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Don't go for weeding .The grasses and weeds will help in maintaining the humidity and it is safe for creeping of the vines and does not allow the fruit to come in contact with soil and avoid fruit rot. High temperature leads to fungal wilting in pointed gourd, spray 1.5 ml Carbendazim in 1 litre water. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxchloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Mango:

Water stress immediately after fruit set increases fruit dropping in Mango. Hence, protective irrigation is essential during the fruit development stage. To control **fruit drop**, spray Borax @2gm per lit. of water and apply Planofix @2ml per 5 lit. of water. For control of mango **fruit borer** spray Neem oil 3ml + chlorpyrifos 1 ml per litre of water along with stickers at marble stage of the fruit. The sprayings should be preferably carried out during evening hours. To control **bacterialcanker**, three sprays of streptomycin (100 ppm) or Agrimycin-100 (100 ppm) after first visual symptom at 10- days intervals is recommended.

Animal husbandry:

Make necessary arrangement for clean and sufficient drinking water for the animals.

Dr Sujata Sethy
 (Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-485
Date- 12/04/2024

AGROMET ADVISORY BULLETIN FOR SALEPUR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0.3 mm and district received average 2mm rainfall during the last week. The mean daily maximum and minimum temperature was 35.47°C (-1.95 from normal) and 23.37°C (-0.3 from normal), respectively. The mean relative humidity was 61-67%.

Forecasted period: Today(8.30IST) to 17th April (8.30IST)2024

Weather advisory: Dry weather is forecasted for today to Tuesday in the block. The day temperature will be 34-37°C and night temperature will be 24-27°C for upcoming 5 days. Maximum and minimum relative humidity will be 52-83% and 18-23% respectively. There will be clear sky for next 5days.

Block	Salepur (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	12/04/2024	13/04/2024	14/04/2024	15/04/2024	16/04/2024
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	34	34	37	37	37
T-Min(°C)	24	27	26	25	26
Cloud Cover(octa)	4	2	2	3	2
RH Max (%)	52	60	61	77	83
RH Min (%)	18	20	21	21	23
Wind Speed (Kmph)	13	14	14	17	20
Wind Direction (deg)	270	210	225	203	203

Crop specific Agro-met Advisory

General advisory: Farmers should harvest the matured crops and vegetables in dry days. Farmers may go for intercultural operations, fertilizer and pesticide application in the upcoming dry days. They should start sowing of summer vegetables by utilizing past rain water. They should provide mulching material or polythene cover to recently planted vegetables seedling to protect from increase in day temperature.

Pulses:

Harvest matured pulses in dry days. Rust infestation likely to occur in pulses, if observe spray Mancozeb 35.0%SC 3gram/ltr of water. If **wilting** will be observed after rainfall, then spray Carbendazim 2g/ltr of water to control.

Summer Maize:

There may be increase the infestation of **Red hairy caterpillar** in maize after rainfall, to minimize the infestation farmers may spray Neem oil 5ml/ltr of water or Chlorpyrifos 20 % EC @ 2ml/litre of water. Prolonged dry spell followed by rain may increase the infestation of **fall army worm in maize**. To control spray emamectin benzoate (5% S.G.) @ 0.4g per litre of water.

Vegetables:

Okra: To control **Yellow Vain Mosaic Virus (YVMV)** in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease.

Brinjal:

There are chances of infestation of **Epilachna Beetle/Haddabeetle** in Brinjal and cucurbits crop; to manage Epilachna Beetle in cucurbits, spray neem-based pesticide (Azadirachtin) 1500 PPM @ 3ml/litre of water at early stage of infestation; to manage Epilachna Beetle chemically, spray Chlorpyrifos 20 % EC @ 2ml/litre of water or Profenophos 50 % EC @ 2ml/litre of water.

FRUIT AND SHOOT BORER IN BRINJAL-Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@250g/acre at 15 days interval. If **phomopsis blight** will occur in brinjal, then spray zineb75%WP 2gram/litr of water. **Cercopspera leaf spot** may be found after rainfall, if observe, then spray Carbendazim or zineb75%WP 2gram/litr of water. If **helicoverpa infestation** will be found, spray Flunendiamide or indoxacarb 2ml/litr of water.

Cucurbits:

To manage **Red Pumpkin Beetle** in Cucurbit crops spray Cyantraniliprole 10.26% OD @ 400-ml/acre or Imidacloprid 17.8% SL @ 60-ml/ acre or Chlorpyrifos 20% EC @ 400-ml/acre. **Late Blight** may happen due to rising day temperature. There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Don't go for weeding .The grasses and weeds will help in maintaining the humidity and it is safe for creeping of the vines and does not allow the fruit to come in contact with soil and avoid fruit rot. High temperature leads to fungal wilting in pointed gourd, spray 1.5 ml Carbendazim in 1 litre water. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxychloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Mango:

Water stress immediately after fruit set increases fruit dropping in Mango. Hence, protective irrigation is essential during the fruit development stage. To control **fruit drop**, spray Borax @2gm per lit. of water and apply Planofix @2ml per 5 lit. of water. For control of mango **fruit borer** spray Neem oil 3ml + chlorpyrifos 1 ml per litre of water along with stickers at marble stage of the fruit. The sprayings should be preferably carried out during evening hours. To control **bacterialcanker**, three sprays of streptocycline (100 ppm) or Agrimycin-100 (100 ppm) after first visual symptom at 10- days intervals is recommended.

Animal husbandry:

Make necessary arrangement for clean and sufficient drinking water for the animals.

Dr Sujata Sethy
 (Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-485
Date- 12/04/2024

AGROMET ADVISORY BULLETIN FOR TANGI-CHOUDWAR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0.3 mm and district received average 2mm rainfall during the last week. The mean daily maximum and minimum temperature was 35.47°C (-1.95 from normal) and 23.37°C (-0.3 from normal), respectively. The mean relative humidity was 61-67%.

Forecasted period: Today(8.30IST) to 17th April (8.30IST)2024

Weather advisory: Dry weather is forecasted for today to Tuesday in the block. The day temperature will be 39-40°C and night temperature will be 27-29°C for upcoming 5 days. Maximum and minimum relative humidity will be 61-83% and 23-36% respectively. There will be clear sky for next 5days.

Block	Tangi-Choudwar (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	12/04/2024	13/04/2024	14/04/2024	15/04/2024	16/04/2024
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	39	39	40	40	39
T-Min(°C)	27	28	28	28	29
Cloud Cover(octa)	3	3	1	2	1
RH Max (%)	61	75	71	79	83
RH Min (%)	23	27	26	30	36
Wind Speed (Kmph)	11	18	18	22	25
Wind Direction (deg)	162	225	238	203	203

Crop specific Agro-met Advisory

General advisory: Farmers should harvest the matured crops and vegetables in dry days. Farmers may go for intercultural operations, fertilizer and pesticide application in the upcoming dry days. They should start sowing of summer vegetables by utilizing past rain water. They should provide mulching material or polythene cover to recently planted vegetables seedling to protect from increase in day temperature.

Pulses:

Harvest matured pulses in dry days. Rust infestation likely to occur in pulses, if observe spray Mancozeb 35.0%SC 3gram/ltr of water. If **wilting** will be observed after rainfall, then spray Carbendazim 2g/ltr of water to control.

Summer Maize:

There may be increase the infestation of **Red hairy caterpillar** in maize after rainfall, to minimize the infestation farmers may spray Neem oil 5ml/ltr of water or Chlorpyrifos 20 % EC @ 2ml/litre of water. Prolonged dry spell followed by rain may increase the infestation of **fall army worm in maize**. To control spray emamectin benzoate (5% S.G.) @ 0.4g per litre of water.

Vegetables:

Okra: To control **Yellow Vain Mosaic Virus (YVMV)** in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease.

Brinjal:

There are chances of infestation of **Epilachna Beetle/Haddabeetle** in Brinjal and cucurbits crop; to manage Epilachna Beetle in cucurbits, spray neem-based pesticide (Azadirachtin) 1500 PPM @ 3ml/litre of water at early stage of infestation; to manage Epilachna Beetle chemically, spray Chlorpyrifos 20 % EC @ 2ml/litre of water or Profenophos 50 % EC @ 2ml/litre of water.

FRUIT AND SHOOT BORER IN BRINJAL-Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@250g/acre at 15 days interval. If **phomopsis blight** will occur in brinjal, then spray zineb75%WP 2gram/litr of water. **Cercopspera leaf spot** may be found after rainfall, if observe, then spray Carbendazim or zineb75%WP 2gram/litr of water. If **helicopterpa infestation** will be found, spray Flunendiamide or indoxacarb 2ml/litr of water.

Cucurbits:

To manage **Red Pumpkin Beetle** in Cucurbit crops spray Cyantraniliprole 10.26% OD @ 400-ml/acre or Imidacloprid 17.8% SL @ 60-ml/ acre or Chlorpyrifos 20% EC @ 400-ml/acre. **Late Blight** may happen due to rising day temperature. There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Don't go for weeding .The grasses and weeds will help in maintaining the humidity and it is safe for creeping of the vines and does not allow the fruit to come in contact with soil and avoid fruit rot. High temperature leads to fungal wilting in pointed gourd, spray 1.5 ml Carbendazim in 1 litre water. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxychloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Mango:

Water stress immediately after fruit set increases fruit dropping in Mango. Hence, protective irrigation is essential during the fruit development stage. To control **fruit drop**, spray Borax @2gm per lit. of water and apply Planofix @2ml per 5 lit. of water. For control of mango **fruit borer** spray Neem oil 3ml + chlorpyrifos 1 ml per litre of water along with stickers at marble stage of the fruit. The sprayings should be preferably carried out during evening hours. To control **bacterialcanker**, three sprays of streptocycline (100 ppm) or Agrimycin-100 (100 ppm) after first visual symptom at 10- days intervals is recommended.

Animal husbandry:

Make necessary arrangement for clean and sufficient drinking water for the animals.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-485
Date- 12/04/2024

AGROMET ADVISORY BULLETIN FOR TIGIRIA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0 mm and district received average 2mm rainfall during the last week. The mean daily maximum and minimum temperature was 35.47°C (-1.95 from normal) and 23.37°C (-0.3 from normal), respectively. The mean relative humidity was 61-67%.

Forecasted period: Today(8.30IST) to 17thApril (8.30IST)2024

Weatheradvisory: *Dry weather is forecasted for today to Tuesday in the block.* The day temperature will be 32-37°C and night temperature will be 23-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 47-77% and 20-21% respectively. There will be clear sky for next 5days.

Block	Tigiria (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	12/04/2024	13/04/2024	14/04/2024	15/04/2024	16/04/2024
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	32	33	36	37	37
T-Min(°C)	23	26	26	25	25
Cloud Cover(octa)	4	3	2	3	3
RH Max (%)	47	48	55	69	77
RH Min (%)	20	20	21	20	20
Wind Speed (Kmph)	9	11	9	11	11
Wind Direction (deg)	248	293	195	171	191

Crop specific Agro-met Advisory

General advisory: *Farmers should harvest the matured crops and vegetables in dry days. Farmers may go for intercultural operations, fertilizer and pesticide application in the upcoming dry days. They should start sowing of summer vegetables by utilizing past rain water. They should provide mulching material or polythene cover to recently planted vegetables seedling to protect from increase in day temperature.*

Pulses:

Harvest matured pulses in dry days. Rust infestation likely to occur in pulses, if observe spray Mancozeb 35.0%SC 3gram/ltr of water. If **wilting** will be observed after rainfall, then spray Carbendazim 2g/ltr of water to control.

Summer Maize:

There may be increase the infestation of **Red hairy caterpillar** in maize after rainfall, to minimize the infestation farmers may spray Neem oil 5ml/ltr of water or Chlorpyrifos 20 % EC @ 2ml/litre of water. Prolonged dry spell followed by rain may increase the infestation of **fall army worm in maize**. **To control spray emamectin benzoate (5% S.G.) @ 0.4g per litre of water.**

Vegetables:

Okra: To control **Yellow Vain Mosaic Virus (YVMV)** in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease.

Brinjal:

There are chances of infestation of **Epilachna Beetle/Haddabeetle** in Brinjal and cucurbits crop; to manage Epilachna Beetle in cucurbits, spray neem-based pesticide (Azadirachtin) 1500 PPM @ 3ml/litre of water at early stage of infestation; to manage Epilachna Beetle chemically, spray Chlorpyrifos 20 % EC @ 2ml/litre of water or Profenophos 50 % EC @ 2ml/litre of water.

FRUIT AND SHOOT BORER IN BRINJAL-Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@250g/acre at 15 days interval. If **phomopsis blight** will occur in brinjal, then spray zineb75%WP 2gram/litr of water.**Cercopspera leaf spot** may be found after rainfall, if observe, then spray Carbendazim or zineb75%WP 2gram/litr of water.If **helicoverpa infestation** will be found, spray Flunendiamide or indoxacarb 2ml/litr of water.

Cucurbits:

To manage **Red Pumpkin Beetle** in Cucurbit crops spray Cyantraniliprole 10.26% OD @ 400-ml/acre or Imidacloprid 17.8% SL @ 60-ml/ acre or Chlorpyrifos 20% EC @ 400-ml/acre. **Late Blight** may happen due to rising day temperature. There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Don't go for weeding .The grasses and weeds will help in maintaining the humidity and it is safe for creeping of the vines and does not allow the fruit to come in contact with soil and avoid fruit rot. High temperature leads to fungal wilting in pointed gourd, spray 1.5 ml Carbendazim in 1 litre water. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxchloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Mango:

Water stress immediately after fruit set increases fruit dropping in Mango. Hence, protective irrigation is essential during the fruit development stage. To control **fruit drop**, spray Borax @2gm per lit. of water and apply Planofix @2ml per 5 lit. of water.For control of mango **fruit borer** spray Neem oil 3ml + chlorpyrifos 1 ml perlitre of water along with stickers at marble stage of the fruit. The sprayingshould be preferably carried out during evening hours. To control **bacterialcanker**, three sprays of streptomycine (100 ppm) or Agrimycin-100 (100 ppm)after first visual symptom at 10- days intervals is recommended.

Animal husbandry:

Make necessary arrangement for clean and sufficient drinking water for the animals.

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area




Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-485
Date- 12/04/2024

				
MEGHDOOT APP	DAMINI APP	FACEBOOK (Damu Cuttack)	BLOCK SPECIFIC WHAT'S APP GROUPS	@DAMUCUTTACK

- Kindly install and use **“Meghdoot mobile App”** for Cuttack district weather forecast, warnings and weather based agromet advisory services.
<https://play.google.com/store/apps/details?id=com.aas.meghdoot>
- Kindly install and use **“Damini mobile App”** for location specific for lightening forecast and warnings.
<https://play.google.com/store/apps/details?id=com.lightening.live.damini>
- Follow the **block specific whats app groups** and join in your block wise groups for getting weather based agromet advisory services twice in a week (Tuesday and Friday) and other weather warning based farm management advisories.
- Follow the **Official Website links** to get the weather based agromet advisory service bulletins (**English & Odia**) twice in a week (Tuesday and Friday).

1	ICAR-NRRI, Cuttack, Odisha	https://icar-nrri.in/agromate/	
2	Agrimet division, IMD, Pune	https://imdagrmet.gov.in/imdproject/AGIndex.php	

Dr Sujata Sethy
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427