




Cuttack District Block Wise Agro-met Advisory Service Bulletin

*Tuesday, 21st May, 2024
 (For the period 21st to 25th May, 2024)*



*Issued by
 District Agro-met Unit,
 KrishiVigyan Kendra Cuttack*





STANDARD PRECIPITATION INDEX AND NDVI MAP

Standard Precipitation Index (SPI)	Normalized difference vegetation Index (NDVI)
<div style="background-color: #90EE90; padding: 5px; text-align: center; margin-bottom: 10px;"> Standardised Precipitation Index for the period 28th September to 25th October 2023 </div> <div style="text-align: center;"> <p>MINISTRY OF EARTH SCIENCES INDIA METEOROLOGICAL DEPARTMENT HYDROMET SECTION, CRS PUNE</p> <p>STANDARDIZED PRECIPITATION INDEX FOR THE PERIOD 28 SEPTEMBER TO 25 OCTOBER 2023</p> <p>SPI VALUES</p> <ul style="list-style-type: none"> 0.0 to 0.5 0.5 to 1.0 1.0 to 1.5 1.5 to 2.0 2.0 to 2.5 2.5 to 3.0 3.0 to 3.5 3.5 to 4.0 4.0 to 4.5 4.5 to 5.0 5.0 to 5.5 5.5 to 6.0 6.0 to 6.5 6.5 to 7.0 7.0 to 7.5 7.5 to 8.0 8.0 to 8.5 8.5 to 9.0 9.0 to 9.5 9.5 to 10.0 </div>	<div style="text-align: center;"> <p>NOAA/VIIRS/BLENDED NDVI Composite ending on Week no 19 (06.05.2024 to 12.05.2024) over Agricultural region</p> <p>Odisha</p> <p style="font-size: small;"> -9999 - 0 0 - 0.05 0.05 - 0.1 0.1 - 0.15 0.15 - 0.2 0.2 - 0.25 0.25 - 0.3 0.3 - 0.35 0.35 - 0.4 0.4 - 0.45 0.45 - 0.5 0.5 - 0.55 0.55 - 0.6 > 0.6 </p> <p style="font-size: small;"> Bare soil, rocks clouds, rain, snow, etc. Moderate Good Very good </p> <p>Agriculture vigour is moderate at few places of the state.</p> </div>
<p><i>As per SPI value (cumulative 4weeks -28 September to 25 October 2023), mildly dry condition was there in Cuttack district.</i></p>	<p><i>According to NDVI map (06.05.2024 to 12.05.2024) the agriculture vigor was very light to light over maximum blocks of Cuttack district.</i></p>



LEGEND (Medium range block level rainfall forecast)		
	No Rainfall	00 mm
	Very light rainfall	01-05 mm
	Light rainfall	06-10 mm
	Light to moderate rainfall	11-30 mm
	Moderate to heavy rainfall	31-65 mm
	Heavy to very heavy rainfall	66-115 mm
	Extremely heavy rainfall	>116 mm

LEGEND (Past week temperature)		
Color code	Temperature	Departure
	Below Normal	< 2°C from normal
	Normal	-2°C to +2°C from normal
	Above Normal	>2°C from normal









Past week (14-20 May, 2024) cumulative rainfall (mm) and its departure from normal weekly rainfall and (14-20 May, 2024) average maximum temperature and its departure from normal for Cuttack district



Medium range (Today (8.30IST) to Sunday (8.30IST)) Rainfall (mm) forecast for Cuttack district















GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-496
Date- 21/05/2024

Weather (Past and Forecasted) based Disease and pest in Major Crops and vegetables

Sl No	Crop	Stage and condition of the crop	Anticipated Disease and pest
1	Tomato	Fruting	 Bacterial wilt
2	Okra	Vegetative to fruiting	 Yellow vein mosaiv(YMV)
			 fusarium and phytophthora
			 Cercospera leaf spot
3	Cucurbits	Flowering to Fruiting stage	 Late blight
			 Downy Mildew
			 Leaf spot
			 Stem Rot
			 Fruit rot
4	Mango	Fruiting stage	 Fruit borer
			 Anthracnose





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK



(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-496
Date- 21/05/2024

AGROMET ADVISORY BULLETIN FOR ATHAGAD

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0.6mm and district received average 01 mm rainfall during the last week. The mean daily maximum and minimum temperature was 37.6°C (+0.45 from normal) and 26.1°C (+1.14 from normal), respectively. The mean relative humidity was 43-97%.

Forecasted period: Today(8.30IST) to 26th May(8.30IST)2024

Weather advisory: Very Light to light rainfall/thundershower is forecasted for today to, Thursday and Tuesday and dry weather is forecasted for other 1 day in the block. The day temperature will be 37-39°C and night temperature will be 26-30°C for upcoming 5 days. Maximum and minimum relative humidity will be 58-80% and 23-29% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Athagad (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	21/05/2024	22/05/2024	23/05/2024	24/05/2024	25/05/2024
Rainfall (mm)	3	2	6	0	5
T-Max (°C)	38	39	37	39	37
T-Min (°C)	27	26	26	28	30
Cloud Cover (octa)	3	3	2	6	8
RH Max (%)	79	79	80	71	58
RH Min (%)	27	29	29	23	28
Wind Speed (kmph)	17	15	13	10	10
Wind Direction (deg)	193	159	158	198	198

Agromet Advisory

General advisory:

Farmers are advised to go for summer ploughing in the dry days as sufficient amount of rainfall was realized in the past and prepare the field for kharif sowing. As increase in day temperature condition likely to prevail in next 5 days, so farmers should cover the soil with natural/straw mulch and advised to strengthen the field bund to conserve present soil moisture.

SUMMER PLOUGHING:

In dry sunny days, go for summer ploughing at places receiving soaking rainfall to reduce weed infestation, to destroy eggs and larvae of hibernated soil insects and to increase soil profile recharge.

GREEN MANURING:

Optimum time for sowing of green manuring crops like Sesbania and sunhemp in the areas receiving convectional rain. Green manuring crops are sown in medium and low land condition at 6-8 weeks before transplanting of rice

Summer Maize:

Where Maize is a standing crop now, at their critical growth stage of soil moisture/irrigation i.e tasseling to silking stage to grain formation which stage demands high water/soil moisture content due to high LAI. Water deficit at that stage will negatively affect the fertilization, grain number per cob, and as a result, the final yield of maize is reduced, So Farmers are advised to maintain proper moisture in the maize crop by giving light irrigation as per requirement.

Vegetables:

To prevent from wilting or partial wilting in brinjal, Chilli and Okra apply Neem or Karanjaa cake and spray (basal + foliar) Tebuconazole fungicide @ 1ml/liter.

Okra: To control Yellow Vain Mosaic Virus (YVMV) in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease. Spray Metalaxyl+Mancozeb/Carboxin+Thiram @ 2g/litre to control rot or soil borne fungal disease. Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenoconazole @ 1ml/lit at 15 days interval to control Cercospora leaf spot.

Cucurbits: Late Blight may happen due to rising day temperature. There are chances of infestation of Downy Mildew in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.


Pointed gourd is at flowering and fruiting stage. Present weather may aggravate fruit and vine rot (symptoms: leaf spot, leaf blight, stem rot, fruit rot) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxchloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Mango: For control of mango fruit borer spray Neem oil 3ml + chloripyriphos 1 ml per litre of water along with stickers at marble stage of the fruit. The sprayings should be preferably carried out during evening hours. Spray the foliage with Tebuconazole 50% + Trifloxystrobin 25% WG @ 1g/lit to control Anthracnose.

Animal husbandry:

As THI likely to be above 72, to escape dairy animals from thermal stress, make necessary arrangement for clean and sufficient drinking water with sufficient fodder crops for the animals. Keep farm animals indoor during the day time period.





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK



(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-496
Date- 21/05/2024

AGROMET ADVISORY BULLETIN FOR BADAMBA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0.6mm and district received average 01 mm rainfall during the last week. **The mean daily maximum and minimum temperature was 37.6°C (+0.45 from normal) and 26.1°C (+1.14 from normal), respectively.** The mean relative humidity was 43-97%.

Forecasted period: Today(8.30IST) to 26th May(8.30IST)2024

Weather advisory: Very Light to light rainfall/thundershower is forecasted for today to, Thursday and Tuesday and dry weather is forecasted for other 1 day in the block. The day temperature will be 37-40°C and night temperature will be 25-29°C for upcoming 5 days. Maximum and minimum relative humidity will be 54-79% and 26-29% respectively. **There will be clear to partly cloudy sky for next 5 days.**

Block	Badamba (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	21/05/2024	22/05/2024	23/05/2024	24/05/2024	25/05/2024
Rainfall(mm)	4	2	6	0	4
T-Max(°C)	39	40	38	39	37
T-Min(°C)	26	25	25	27	29
Cloud Cover(octa)	3	3	3	5	8
RH Max (%)	79	74	78	70	54
RH Min (%)	26	27	29	26	28
Wind Speed (Kmph)	16	14	13	9	9
Wind Direction (deg)	180	158	162	180	199

Agromet Advisory

General advisory:

Farmers are advised to go for summer ploughing in the dry days as sufficient amount of rainfall was realized in the past and prepare the field for kharif sowing. As increase in day temperature condition likely to prevail in next 5 days, so farmers should cover the soil with natural/straw mulch and advised to strengthen the field bund to conserve present soil moisture.

SUMMER PLOUGHING:

In dry sunny days, go for summer ploughing at places receiving soaking rainfall to reduce weed infestation, to destroy eggs and larvae of hibernated soil insects and to increase soil profile recharge.

GREEN MANURING:

Optimum time for sowing of green manuring crops like Sesbania and sunhemp in the areas receiving convectional rain. Green manuring crops are sown in medium and low land condition at 6-8 weeks before transplanting of rice

Summer Maize:

Where Maize is a standing crop now, at their critical growth stage of soil moisture/irrigation i.e tasseling to silking stage to grain formation which stage demands high water/soil moisture content due to high LAI. Water deficit at that stage will negatively affect the fertilization, grain number per cob, and as a result, the final yield of maize is reduced, So Farmers are advised to maintain proper moisture in the maize crop by giving light irrigation as per requirement.

Vegetables:

To prevent from wilting or partial wilting in brinjal, Chilli and Okra apply Neem or Karanjaa cake and spray (basal + foliar) Tebuconazole fungicide @ 1ml/liter.

Okra: To control **Yellow Vain Mosaic Virus (YVMV)** in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease. Spray Metalaxyl+Mancozeb/Carboxin+Thiram @ 2g/litre to control **rot or soil borne fungal disease.** Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenconazole @ 1ml/lit at 15 days interval to control **Cercospora leaf spot.**

Cucurbits: **Late Blight** may happen due to rising day temperature. There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxchloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Mango: For control of mango **fruit borer** spray Neem oil 3ml + chloripyriphos 1 ml per litre of water along with stickers at marble stage of the fruit. The sprayings should be preferably carried out during evening hours. Spray the foliage with Tebuconazole 50% + Trifloxystrobin 25% WG @ 1g/lit to control

Anthracnose.

Animal husbandry:

As THI likely to be above 72, to escape dairy animals from thermal stress, make necessary arrangement for clean and sufficient drinking water with sufficient fodder crops for the animals. Keep farm animals indoor during the day time period.





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK



(A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-496
 Date- 21/05/2024

AGROMET ADVISORY BULLETIN FOR BANKI

District Past week average weather condition and Block past week average Rainfall condition

The district received average 01 mm rainfall during the last week. The mean daily maximum and minimum temperature was 37.6°C (+0.45 from normal) and 26.1°C (+1.14 from normal), respectively. The mean relative humidity was 43-97%.

Forecasted period: Today(8.30IST) to 26th May(8.30IST)2024

Weather advisory: Very Light to light rainfall/thundershower is forecasted for today to, Thursday and Tuesday and dry weather is forecasted for other 1 day in the block. The day temperature will be 37-39°C and night temperature will be 25-30°C for upcoming 5 days. Maximum and minimum relative humidity will be 58-80% and 25-31% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Banki (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	21/05/2024	22/05/2024	23/05/2024	24/05/2024	25/05/2024
Rainfall(mm)	3	2	6	0	4
T-Max(°C)	38	39	37	39	37
T-Min(°C)	26	25	26	27	30
Cloud Cover(octa)	2	3	3	6	8
RH Max (%)	80	77	79	71	58
RH Min (%)	29	29	31	25	29
Wind Speed (Kmph)	18	16	14	11	11
Wind Direction (deg)	193	162	159	191	198

Agromet Advisory

General advisory:

Farmers are advised to go for summer ploughing in the dry days as sufficient amount of rainfall was realized in the past and prepare the field for kharif sowing. As increase in day temperature condition likely to prevail in next 5 days, so farmers should cover the soil with natural/straw mulch and advised to strengthen the field bund to conserve present soil moisture.

SUMMER PLOUGHING:

In dry sunny days, go for summer ploughing at places receiving soaking rainfall to reduce weed infestation, to destroy eggs and larvae of hibernated soil insects and to increase soil profile recharge.

GREEN MANURING:

Optimum time for sowing of green manuring crops like Sesbania and sunhemp in the areas receiving convectional rain. Green manuring crops are sown in medium and low land condition at 6-8 weeks before transplanting of rice

Summer Maize:

Where Maize is a standing crop now, at their critical growth stage of soil moisture/irrigation i.e tasseling to silking stage to grain formation which stage demands high water/soil moisture content due to high LAI. Water deficit at that stage will negatively affect the fertilization, grain number per cob, and as a result, the final yield of maize is reduced. So Farmers are advised to maintain proper moisture in the maize crop by giving light irrigation as per requirement.

Vegetables:

To prevent from wilting or partial wilting in brinjal, Chilli and Okra apply Neem or Karanjaa cake and spray (basal + foliar) Tebuconazole fungicide @ 1ml/liter.

Okra: To control Yellow Vain Mosaic Virus (YVMV) in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease. Spray Metalaxyl+Mancozeb/Carboxin+Thiram @ 2g/litre to control rot or soil borne fungal disease. Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenconazole @ 1ml/lit at 15 days interval to control Cercospora leaf spot.

Cucurbits: Late Blight may happen due to rising day temperature. There are chances of infestation of Downy Mildew in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Present weather may aggravate fruit and vine rot (symptoms: leaf spot, leaf blight, stem rot, fruit rot) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxchloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Mango: For control of mango fruit borer spray Neem oil 3ml + chloripyriphos 1 ml per litre of water along with stickers at marble stage of the fruit. The sprayings should be preferably carried out during evening hours. Spray the foliage with Tebuconazole 50% + Trifloxystrobin 25% WG @ 1g/lit to control Anthracnose.

Animal husbandry:

As THI likely to be above 72, to escape dairy animals from thermal stress, make necessary arrangement for clean and sufficient drinking water with sufficient fodder crops for the animals. Keep farm animals indoor during the day time period.





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK



(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-496
Date- 21/05/2024

AGROMET ADVISORY BULLETIN FOR BANKI-DAMPARA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0.3mm and district received average 01 mm rainfall during the last week. The mean daily maximum and minimum temperature was 37.6°C (+0.45 from normal) and 26.1°C (+1.14 from normal), respectively. The mean relative humidity was 43-97%.

Forecasted period: Today(8.30IST) to 26thMay(8.30IST)2024

Weather Advisory: Very Light to light rainfall/thundershower is forecasted for today to, Thursday and Tuesday and dry weather is forecasted for other 1 day in the block. The day temperature will be 37-39°C and night temperature will be 26-30°C for upcoming 5 days. Maximum and minimum relative humidity will be 59-80% and 11-18% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Banki-Dampara (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	21/05/2024	22/05/2024	23/05/2024	24/05/2024	25/05/2024
Rainfall(mm)	4	2	6	0	4
T-Max(°C)	38	39	37	39	37
T-Min(°C)	26	26	26	27	30
Cloud Cover(octa)	2	3	2	6	8
RH Max (%)	80	78	79	71	59
RH Min (%)	29	29	31	25	29
Wind Speed (Kmph)	18	16	14	11	11
Wind Direction (deg)	193	162	159	199	198

Agromet Advisory

General advisory:

Farmers are advised to go for summer ploughing in the dry days as sufficient amount of rainfall was realized in the past and prepare the field for kharif sowing. As increase in day temperature condition likely to prevail in next 5 days, so farmers should cover the soil with natural/straw mulch and advised to strengthen the field bund to conserve present soil moisture.

SUMMER PLOUGHING:

In dry sunny days, go for summer ploughing at places receiving soaking rainfall to reduce weed infestation, to destroy eggs and larvae of hibernated soil insects and to increase soil profile recharge.

GREEN MANURING:

Optimum time for sowing of green manuring crops like Sesbania and sunhemp in the areas receiving convectional rain. Green manuring crops are sown in medium and low land condition at 6-8 weeks before transplanting of rice

Summer Maize:

Where Maize is a standing crop now, at their critical growth stage of soil moisture/irrigation i.e tasseling to silking stage to grain formation which stage demands high water/soil moisture content due to high LAI. Water deficit at that stage will negatively affect the fertilization, grain number per cob, and as a result, the final yield of maize is reduced. So Farmers are advised to maintain proper moisture in the maize crop by giving light irrigation as per requirement.

Vegetables:

To prevent from wilting or partial wilting in brinjal, Chilli and Okra apply Neem or Karanjaa cake and spray (basal + foliar) Tebuconazole fungicide @ 1ml/liter.

Okra: To control Yellow Vain Mosaic Virus (YVMV) in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease. Spray Metalaxyl+Mancozeb/Carboxin+Thiram @ 2g/litre to control rot or soil borne fungal disease. Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenoconazole @ 1ml/lit at 15 days interval to control Cercospora leaf spot.

Cucurbits: Late Blight may happen due to rising day temperature. There are chances of infestation of Downy Mildew in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Present weather may aggravate fruit and vine rot (symptoms: leaf spot, leaf blight, stem rot, fruit rot) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxchloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Mango: For control of mango fruit borer spray Neem oil 3ml + chloripyriphos 1 ml per litre of water along with stickers at marble stage of the fruit. The spraying should be preferably carried out during evening hours. Spray the foliage with Tebuconazole 50% + Trifloxystrobin 25% WG @ 1g/lit to control Anthracnose.

Animal husbandry:

As THI likely to be above 72, to escape dairy animals from thermal stress, make necessary arrangement for clean and sufficient drinking water with sufficient fodder crops for the animals. Keep farm animals indoor during the day time period.





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK



(A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-496
 Date- 21/05/2024

AGROMET ADVISORY BULLETIN FOR BARANGA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0.6mm and district received average 01 mm rainfall during the last week. **The mean daily maximum and minimum temperature was 37.6°C (+0.45 from normal) and 26.1°C (+1.14 from normal), respectively.** The mean relative humidity was 43-97%.

Forecasted period: Today(8.30IST) to 26thMay(8.30IST)2024

Weather advisory: *Very Light to light rainfall/thundershower is forecasted for today to, Thursday and Tuesday and dry weather is forecasted for other 1 day in the block.* The day temperature will be 36-37°C and night temperature will be 27-30°C for upcoming 5 days. Maximum and minimum relative humidity will be 68-82% and 26-32% respectively. **There will be clear to partly cloudy sky for next 5 days.**

Block	Baranga (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	21/05/2024	22/05/2024	23/05/2024	24/05/2024	25/05/2024
Rainfall(mm)	4	2	6	0	5
T-Max(°C)	37	37	36	37	36
T-Min(°C)	28	27	27	28	30
Cloud Cover	2	2	2	6	8
RH Max (%)	79	81	82	73	68
RH Min (%)	32	31	32	26	32
Wind Speed (Kmph)	19	16	14	13	13
Wind Direction (deg)	198	180	162	203	203

Agromet Advisory

General advisory:

Farmers are advised to go for summer ploughing in the dry days as sufficient amount of rainfall was realized in the past and prepare the field for kharif sowing. As increase in day temperature condition likely to prevail in next 5 days, so farmers should cover the soil with natural/straw mulch and advised to strengthen the field bund to conserve present soil moisture.

SUMMER PLOUGHING:

In dry sunny days, go for summer ploughing at places receiving soaking rainfall to reduce weed infestation, to destroy eggs and larvae of hibernated soil insects and to increase soil profile recharge.

GREEN MANURING:

Optimum time for sowing of green manuring crops like Sesbania and sunhemp in the areas receiving convectional rain. Green manuring crops are sown in medium and low land condition at 6-8 weeks before transplanting of rice

Summer Maize:

Where Maize is a standing crop now, at their critical growth stage of soil moisture/irrigation i.e tasseling to silking stage to grain formation which stage demands high water/soil moisture content due to high LAI. Water deficit at that stage will negatively affect the fertilization, grain number per cob, and as a result, the final yield of maize is reduced. So Farmers are advised to maintain proper moisture in the maize crop by giving light irrigation as per requirement.

Vegetables:

To prevent from wilting or partial wilting in brinjal, Chilli and Okra apply Neem or Karanjaa cake and spray (basal + foliar) Tebuconazole fungicide @ 1ml/liter.

Okra: To control **Yellow Vain Mosaic Virus (YVMV)** in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease. Spray Metalaxyl+Mancozeb/Carboxin+Thiram @ 2g/litre to control **rot or soil borne fungal disease**. Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenconazole @ 1ml/lit at 15 days interval to control **Cercospora leaf spot**.

Cucurbits: **Late Blight** may happen due to rising day temperature. There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxchloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Mango: For control of mango **fruit borer** spray Neem oil 3ml + chloripyriphos 1 ml per litre of water along with stickers at marble stage of the fruit. The spraying should be preferably carried out during evening hours. Spray the foliage with Tebuconazole 50% + Trifloxystrobin 25% WG @ 1g/lit to control **Anthraco**se.

Animal husbandry:

As THI likely to be above 72, to escape dairy animals from thermal stress, make necessary arrangement for clean and sufficient drinking water with sufficient fodder crops for the animals. Keep farm animals indoor during the day time period.





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK



(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-496
Date- 21/05/2024

AGROMET ADVISORY BULLETIN FOR CUTTACK SADAR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 1.3mm and district received average 01 mm rainfall during the last week. **The mean daily maximum and minimum temperature was 37.6°C (+0.45 from normal) and 26.1°C (+1.14 from normal), respectively.** The mean relative humidity was 43-97%.

Forecasted period: Today(8.30IST) to 26thMay(8.30IST)2024

Weather advisory: *Very Light to light rainfall/thundershower is forecasted for today to, Thursday and Tuesday and dry weather is forecasted for other 1 day in the block.* The day temperature will be 36-37°C and night temperature will be 27-30°C for upcoming 5 days. Maximum and minimum relative humidity will be 64-81% and 24-31% respectively. **There will be clear to partly cloudy sky for next 5 days.**

Block	Cuttack sadar (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	21/05/2024	22/05/2024	23/05/2024	24/05/2024	25/05/2024
Rainfall(mm)	3	2	6	0	6
T-Max(°C)	37	37	36	37	36
T-Min(°C)	28	27	27	29	30
Cloud Cover(octa)	2	3	2	6	8
RH Max (%)	79	80	81	72	64
RH Min (%)	28	30	31	24	29
Wind Speed (Kmph)	17	16	13	12	11
Wind Direction (deg)	193	162	158	207	201

Agromet Advisory

General advisory:

Farmers are advised to go for summer ploughing in the dry days as sufficient amount of rainfall was realized in the past and prepare the field for kharif sowing. As increase in day temperature condition likely to prevail in next 5 days, so farmers should cover the soil with natural/straw mulch and advised to strengthen the field bund to conserve present soil moisture.

SUMMER PLOUGHING:

In dry sunny days, go for summer ploughing at places receiving soaking rainfall to reduce weed infestation, to destroy eggs and larvae of hibernated soil insects and to increase soil profile recharge.

GREEN MANURING:

Optimum time for sowing of green manuring crops like Sesbania and sunhemp in the areas receiving convectional rain. Green manuring crops are sown in medium and low land condition at 6-8 weeks before transplanting of rice

Summer Maize:

Where Maize is a standing crop now, at their critical growth stage of soil moisture/irrigation i.e tasseling to silking stage to grain formation which stage demands high water/soil moisture content due to high LAI. Water deficit at that stage will negatively affect the fertilization, grain number per cob, and as a result, the final yield of maize is reduced. So Farmers are advised to maintain proper moisture in the maize crop by giving light irrigation as per requirement.

Vegetables:

To prevent from wilting or partial wilting in brinjal, Chilli and Okra apply Neem or Karanjaa cake and spray (basal + foliar) Tebuconazole fungicide @ 1ml/liter.

Okra: To control **Yellow Vain Mosaic Virus (YVMV)** in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease. Spray Metalaxyl+Mancozeb/Carboxin+Thiram @ 2g/litre to control **rot or soil borne fungal disease**. Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenoconazole @ 1ml/lit at 15 days interval to control **Cercospora leaf spot**.

Cucurbits: **Late Blight** may happen due to rising day temperature. There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxchloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Mango: For control of mango **fruit borer** spray Neem oil 3ml + chloripyriphos 1 ml per litre of water along with stickers at marble stage of the fruit. The spraying should be preferably carried out during evening hours. Spray the foliage with Tebuconazole 50% + Trifloxystrobin 25% WG @ 1g/lit to control **Anthraco**se.

Animal husbandry:

As THI likely to be above 72, to escape dairy animals from thermal stress, make necessary arrangement for clean and sufficient drinking water with sufficient fodder crops for the animals. Keep farm animals indoor during the day time period.





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK



(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-496

Date- 21/05/2024

AGROMET ADVISORY BULLETIN FOR KANTAPADA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 2.6mm and district received average 01 mm rainfall during the last week. The mean daily maximum and minimum temperature was 37.6°C (+0.45 from normal) and 26.1°C (+1.14 from normal), respectively. The mean relative humidity was 43-97%.

Forecasted period: Today(8.30IST) to 26th May(8.30IST)2024

Weather advisory: Very Light to light rainfall/thundershower is forecasted for today to, Thursday and Tuesday and dry weather is forecasted for other 1 day in the block. The day temperature will be 36-37°C and night temperature will be 27-30°C for upcoming 5 days. Maximum and minimum relative humidity will be 72-83% and 29-36% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Kantapada (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	21/05/2024	22/05/2024	23/05/2024	24/05/2024	25/05/2024
Rainfall(mm)	4	2	6	0	8
T-Max(°C)	36	36	36	37	37
T-Min(°C)	28	27	27	29	30
Cloud Cover(octa)	1	1	2	6	8
RH Max (%)	78	82	83	74	72
RH Min (%)	36	35	36	29	33
Wind Speed (Kmph)	19	16	14	14	16
Wind Direction (deg)	198	193	162	225	203

Agromet Advisory

General advisory:

Farmers are advised to go for summer ploughing in the dry days as sufficient amount of rainfall was realized in the past and prepare the field for kharif sowing. As increase in day temperature condition likely to prevail in next 5 days, so farmers should cover the soil with natural/straw mulch and advised to strengthen the field bund to conserve present soil moisture.

SUMMER PLOUGHING:

In dry sunny days, go for summer ploughing at places receiving soaking rainfall to reduce weed infestation, to destroy eggs and larvae of hibernated soil insects and to increase soil profile recharge.

GREEN MANURING:

Optimum time for sowing of green manuring crops like Sesbania and sunhemp in the areas receiving convectional rain. Green manuring crops are sown in medium and low land condition at 6-8 weeks before transplanting of rice

Summer Maize:

Where Maize is a standing crop now, at their critical growth stage of soil moisture/irrigation i.e tasseling to silking stage to grain formation which stage demands high water/soil moisture content due to high LAI. Water deficit at that stage will negatively affect the fertilization, grain number per cob, and as a result, the final yield of maize is reduced. So Farmers are advised to maintain proper moisture in the maize crop by giving light irrigation as per requirement.

Vegetables:

To prevent from wilting or partial wilting in brinjal, Chilli and Okra apply Neem or Karanjaa cake and spray (basal + foliar) Tebuconazole fungicide @ 1ml/litre.

Okra: To control Yellow Vain Mosaic Virus (YVMV) in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease. Spray Metalaxyl+Mancozeb/Carboxin+Thiram @ 2g/litre to control rot or soil borne fungal disease. Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenconazole @ 1ml/lit at 15 days interval to control Cercospora leaf spot.

Cucurbits: Late Blight may happen due to rising day temperature. There are chances of infestation of Downy Mildew in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Present weather may aggravate fruit and vine rot (symptoms: leaf spot, leaf blight, stem rot, fruit rot) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxychloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Mango: For control of mango fruit borer spray Neem oil 3ml + chloripyriphos 1 ml per litre of water along with stickers at marble stage of the fruit. The sprayings should be preferably carried out during evening hours. Spray the foliage with Tebuconazole 50% + Trifloxystrobin 25% WG @ 1g/lit to control Anthracnose.

Animal husbandry:

As THI likely to be above 72, to escape dairy animals from thermal stress, make necessary arrangement for clean and sufficient drinking water with sufficient fodder crops for the animals. Keep farm animals indoor during the day time period.





**GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK**



(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-496
Date- 21/05/2024

AGROMET ADVISORY BULLETIN FOR MAHANGA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0.6mm and district received average 01 mm rainfall during the last week. The mean daily maximum and minimum temperature was 37.6°C (+0.45 from normal) and 26.1°C (+1.14 from normal), respectively. The mean relative humidity was 43-97%.

Forecasted period: Today(8.30IST) to 26thMay(8.30IST)2024

Weather advisory: Very Light to light rainfall/thundershower is forecasted for today to, Thursday and Tuesday and dry weather is forecasted for other 1 day in the block. The day temperature will be and it will be 36-38°C and night temperature will be 27-30°C for upcoming 5 days. Maximum and minimum relative humidity will be 61-80% and 28-33% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Mahanga (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	21/05/2024	22/05/2024	23/05/2024	24/05/2024	25/05/2024
Rainfall(mm)	5	2	6	0	8
T-Max(°C)	38	37	36	37	37
T-Min(°C)	28	27	27	29	30
Cloud Cover(octal)	1	2	2	7	8
RH Max (%)	78	79	80	73	61
RH Min (%)	30	32	33	32	28
Wind Speed (Kmph)	17	16	13	14	12
Wind Direction (deg)	193	180	124	240	206

Agromet Advisory

General advisory:

Farmers are advised to go for summer ploughing in the dry days as sufficient amount of rainfall was realized in the past and prepare the field for kharif sowing. As increase in day temperature condition likely to prevail in next 5 days, so farmers should cover the soil with natural/straw mulch and advised to strengthen the field bund to conserve present soil moisture.

SUMMER PLOUGHING:

In dry sunny days, go for summer ploughing at places receiving soaking rainfall to reduce weed infestation, to destroy eggs and larvae of hibernated soil insects and to increase soil profile recharge.

GREEN MANURING:

Optimum time for sowing of green manuring crops like Sesbania and sunhemp in the areas receiving convectional rain. Green manuring crops are sown in medium and low land condition at 6-8 weeks before transplanting of rice

Summer Maize:

Where Maize is a standing crop now, at their critical growth stage of soil moisture/irrigation i.e tasseling to silking stage to grain formation which stage demands high water/soil moisture content due to high LAI. Water deficit at that stage will negatively affect the fertilization, grain number per cob, and as a result, the final yield of maize is reduced. So Farmers are advised to maintain proper moisture in the maize crop by giving light irrigation as per requirement.

Vegetables:

To prevent from wilting or partial wilting in brinjal, Chilli and Okra apply Neem or Karanjaa cake and spray (basal + foliar) Tebuconazole fungicide @ 1ml/litre.

Okra: To control Yellow Vain Mosaic Virus (YVMV) in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease. Spray Metalaxyl+Mancozeb/Carboxin+Thiram @ 2g/litre to control rot or soil borne fungal disease. Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenconazole @ 1ml/lit at 15 days interval to control Cercospora leaf spot.

Cucurbits: Late Blight may happen due to rising day temperature. There are chances of infestation of Downy Mildew in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Present weather may aggravate fruit and vine rot (symptoms: leaf spot, leaf blight, stem rot, fruit rot) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxychloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Mango: For control of mango fruit borer spray Neem oil 3ml + chloripyriphos 1 ml per litre of water along with stickers at marble stage of the fruit. The sprayings should be preferably carried out during evening hours. Spray the foliage with Tebuconazole 50% + Trifloxystrobin 25% WG @ 1g/lit to control Anthracnose.

Animal husbandry:

As THI likely to be above 72, to escape dairy animals from thermal stress, make necessary arrangement for clean and sufficient drinking water with sufficient fodder crops for the animals. Keep farm animals indoor during the day time period.





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK



(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-496
Date- 21/05/2024

AGROMET ADVISORY BULLETIN FOR NARASINGHPUR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 01 mm rainfall during the last week. **The mean daily maximum and minimum temperature was 37.6°C (+0.45 from normal) and 26.1°C (+1.14 from normal), respectively.** The mean relative humidity was 43-97%.

Forecasted period: Today(8.30IST) to 26thMay(8.30IST)2024

Weather advisory: Very Light to light rainfall/thundershower is forecasted for today to, Thursday and Tuesday and dry weather is forecasted for other 1 day in the block. The day temperature will be 37-41°C and night temperature will be 24-28°C for upcoming 5 days. Maximum and minimum relative humidity will be 55-81% and 22-28% respectively. **There will be clear to partly cloudy sky for next 5 days.**

Block	Narasinghpur (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	21/05/2024	22/05/2024	23/05/2024	24/05/2024	25/05/2024
Rainfall(mm)	3	2	6	0	3
T-Max(°C)	41	41	40	39	37
T-Min(°C)	25	25	24	26	28
Cloud Cover(/octa)	3	3	3	5	8
RH Max (%)	81	72	81	73	55
RH Min (%)	22	22	26	28	28
Wind Speed (Kmph)	11	10	10	7	8
Wind Direction (deg)	161	156	161	180	203

Agromet Advisory

General advisory:

Farmers are advised to go for summer ploughing in the dry days as sufficient amount of rainfall was realized in the past and prepare the field for kharif sowing. As increase in day temperature condition likely to prevail in next 5 days, so farmers should cover the soil with natural/straw mulch and advised to strengthen the field bund to conserve present soil moisture.

SUMMER PLOUGHING:

In dry sunny days, go for summer ploughing at places receiving soaking rainfall to reduce weed infestation, to destroy eggs and larvae of hibernated soil insects and to increase soil profile recharge.

GREEN MANURING:

Optimum time for sowing of green manuring crops like Sesbania and sunhemp in the areas receiving convectional rain. Green manuring crops are sown in medium and low land condition at 6-8 weeks before transplanting of rice

Summer Maize:

Where Maize is a standing crop now, at their critical growth stage of soil moisture/irrigation i.e tasseling to silking stage to grain formation which stage demands high water/soil moisture content due to high LAI. Water deficit at that stage will negatively affect the fertilization, grain number per cob, and as a result, the final yield of maize is reduced, So Farmers are advised to maintain proper moisture in the maize crop by giving light irrigation as per requirement.

Vegetables:

To prevent from wilting or partial wilting in brinjal, Chilli and Okra apply Neem or Karanjaa cake and spray (basal + foliar) Tebuconazole fungicide @ 1ml/liter.

Okra: To control **Yellow Vain Mosaic Virus (YVMV)** in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease. Spray Metalaxyl+Mancozeb/Carboxin+Thiram @ 2g/litre to control **rot or soil borne fungal disease.** Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenoconazole @ 1ml/lit at 15 days interval to control **Cercospora leaf spot.**

Cucurbits: **Late Blight** may happen due to rising day temperature. There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxychloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Mango: For control of mango **fruit borer** spray Neem oil 3ml + chloripyriphos 1 ml per litre of water along with stickers at marble stage of the fruit. The sprayings should be preferably carried out during evening hours. Spray the foliage with Tebuconazole 50% + Trifloxystrobin 25% WG @ 1g/lit to control **Anthraxnose.**

Animal husbandry:

As THI likely to be above 72, to escape dairy animals from thermal stress, make necessary arrangement for clean and sufficient drinking water with sufficient fodder crops for the animals. Keep farm animals indoor during the day time period.





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK



(A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-496
 Date- 21/05/2024

AGROMET ADVISORY BULLETIN FOR NIALI

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 01 mm rainfall during the last week. **The mean daily maximum and minimum temperature was 37.6°C (+0.45 from normal) and 26.1°C (+1.14 from normal), respectively.** The mean relative humidity was 43-97%.

Forecasted period: Today(8.30IST) to 26thMay(8.30IST)2024

Weather advisory: Very Light to light rainfall/thundershower is forecasted for today to, Thursday and Tuesday and dry weather is forecasted for other 1 day in the block. The day temperature will be 36-38°C and night temperature will be 27-30°C for upcoming 5 days. Maximum and minimum relative humidity will be 74-83% and 32-41% respectively. **There will be clear to partly cloudy sky for next 5days.**

Block	Niali (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	21/05/2024	22/05/2024	23/05/2024	24/05/2024	25/05/2024
Rainfall(mm)	4	2	6	0	9
T-Max(°C)	36	36	36	37	38
T-Min(°C)	28	27	27	29	30
Cloud Cover(octa)	1	1	2	7	8
RH Max (%)	78	82	83	76	74
RH Min (%)	41	39	40	32	35
Wind Speed(Kmph)	19	16	14	16	18
Wind Direction(deg)	198	180	162	225	207

Agromet Advisory

General advisory:

Farmers are advised to go for summer ploughing in the dry days as sufficient amount of rainfall was realized in the past and prepare the field for kharif sowing. As increase in day temperature condition likely to prevail in next 5 days, so farmers should cover the soil with natural/straw mulch and advised to strengthen the field bund to conserve present soil moisture.

SUMMER PLOUGHING:

In dry sunny days, go for summer ploughing at places receiving soaking rainfall to reduce weed infestation, to destroy eggs and larvae of hibernated soil insects and to increase soil profile recharge.

GREEN MANURING:

Optimum time for sowing of green manuring crops like Sesbania and sunhemp in the areas receiving convectional rain. Green manuring crops are sown in medium and low land condition at 6-8 weeks before transplanting of rice

Summer Maize:

Where Maize is a standing crop now, at their critical growth stage of soil moisture/irrigation i.e tasseling to silking stage to grain formation which stage demands high water/soil moisture content due to high LAI. Water deficit at that stage will negatively affect the fertilization, grain number per cob, and as a result, the final yield of maize is reduced, So Farmers are advised to maintain proper moisture in the maize crop by giving light irrigation as per requirement.

Vegetables:

To prevent from wilting or partial wilting in brinjal, Chilli and Okra apply Neem or Karanjaa cake and spray (basal + foliar) Tebuconazole fungicide @ 1ml/litre.

Okra: To control **Yellow Vain Mosaic Virus (YVMV)** in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease. Spray Metalaxyl+Mancozeb/Carboxin+Thiram @ 2g/litre to control **rot or soil borne fungal disease.** Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenoconazole @ 1ml/lit at 15 days interval to control **Cercospora leaf spot.**

Cucurbits: **Late Blight** may happen due to rising day temperature. There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxychloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Mango: For control of mango **fruit borer** spray Neem oil 3ml + chloripyriphos 1 ml per litre of water along with stickers at marble stage of the fruit. The sprayings should be preferably carried out during evening hours. Spray the foliage with Tebuconazole 50% + Trifloxystrobin 25% WG @ 1g/lit to control **Anthracnose.**

Animal husbandry:

As THI likely to be above 72, to escape dairy animals from thermal stress, make necessary arrangement for clean and sufficient drinking water with sufficient fodder crops for the animals. Keep farm animals indoor during the day time period.





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK



(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-496
Date- 21/05/2024

AGROMET ADVISORY BULLETIN FOR NISCHINTAKOILI

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0.6mm and district received average 01 mm rainfall during the last week. **The mean daily maximum and minimum temperature was 37.6°C (+0.45 from normal) and 26.1°C (+1.14 from normal), respectively.** The mean relative humidity was 43-97%.

Forecasted period: Today(8.30IST) to 26thMay(8.30IST)2024

Weather advisory: *Very Light to light rainfall/thundershower is forecasted for today to, Thursday and Tuesday and dry weather is forecasted for other 1 day in the block.* The day temperature will be 36-37°C and night temperature will be 27-30°C for upcoming 5 days. Maximum and minimum relative humidity will be 66-81% and 30-34% respectively. **There will be clear to partly cloudy sky for next 5 days.**

Block	Nischintakoili (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	21/05/2024	22/05/2024	23/05/2024	24/05/2024	25/05/2024
Rainfall(mm)	5	2	6	0	9
T-Max(°C)	37	37	36	37	37
T-Min(°C)	28	27	27	29	30
Cloud Cover(octa)	1	1	2	7	8
RH Max (%)	78	80	81	74	66
RH Min (%)	34	33	34	32	30
Wind Speed (Kmph)	18	16	14	14	13
Wind Direction (deg)	193	180	124	240	207

Agromet Advisory

General advisory:

Farmers are advised to go for summer ploughing in the dry days as sufficient amount of rainfall was realized in the past and prepare the field for kharif sowing. As increase in day temperature condition likely to prevail in next 5 days, so farmers should cover the soil with natural/straw mulch and advised to strengthen the field bund to conserve present soil moisture.

SUMMER PLOUGHING:

In dry sunny days, go for summer ploughing at places receiving soaking rainfall to reduce weed infestation, to destroy eggs and larvae of hibernated soil insects and to increase soil profile recharge.

GREEN MANURING:

Optimum time for sowing of green manuring crops like Sesbania and sunhemp in the areas receiving convectional rain. Green manuring crops are sown in medium and low land condition at 6-8 weeks before transplanting of rice

Summer Maize:

Where Maize is a standing crop now, at their critical growth stage of soil moisture/irrigation i.e tasseling to silking stage to grain formation which stage demands high water/soil moisture content due to high LAI. Water deficit at that stage will negatively affect the fertilization, grain number per cob, and as a result, the final yield of maize is reduced, So Farmers are advised to maintain proper moisture in the maize crop by giving light irrigation as per requirement.

Vegetables:

To prevent from wilting or partial wilting in brinjal, Chilli and Okra apply Neem or Karanjaa cake and spray (basal + foliar) Tebuconazole fungicide @ 1ml/liter.

Okra: To control **Yellow Vain Mosaic Virus (YVMV)** in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease. Spray Metalaxyl+Mancozeb/Carboxin+Thiram @ 2g/litre to control **rot or soil borne fungal disease.** Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenconazole @ 1ml/lit at 15 days interval to control **Cercospora leaf spot.**

Cucurbits: **Late Blight** may happen due to rising day temperature. There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxchloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Mango: For control of mango **fruit borer** spray Neem oil 3ml + chloripyriphos 1 ml per litre of water along with stickers at marble stage of the fruit. The sprayings should be preferably carried out during evening hours. Spray the foliage with Tebuconazole 50% + Trifloxystrobin 25% WG @ 1g/lit to control **Anthracnose.**

Animal husbandry:

As THI likely to be above 72, to escape dairy animals from thermal stress, make necessary arrangement for clean and sufficient drinking water with sufficient fodder crops for the animals. Keep farm animals indoor during the day time period.





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-496
Date- 21/05/2024

AGROMET ADVISORY BULLETIN FOR SALEPUR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 2.6mm and district received average 01 mm rainfall during the last week. **The mean daily maximum and minimum temperature was 37.6°C (+0.45 from normal) and 26.1°C (+1.14 from normal), respectively.** The mean relative humidity was 43-97%.

Forecasted period: Today(8.30IST) to 26thMay(8.30IST)2024

Weather advisory: *Very Light to light rainfall/thundershower is forecasted for today to, Thursday and Tuesday and dry weather is forecasted for other 1 day in the block.* The day temperature will be 36-37°C and night temperature will be 27-30°C for upcoming 5 days. Maximum and minimum relative humidity will be 66-81% and 29-33% respectively. **There will be clear to partly cloudy sky for next 5 days.**

Block	Salepur (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	21/05/2024	22/05/2024	23/05/2024	24/05/2024	25/05/2024
Rainfall(mm)	4	2	6	0	8
T-Max(°C)	37	37	36	37	37
T-Min(°C)	28	27	27	29	30
Cloud Cover(octa)	1	1	2	7	8
RH Max (%)	78	80	81	74	66
RH Min (%)	31	32	33	30	29
Wind Speed (Kmph)	18	16	14	14	13
Wind Direction (deg)	193	180	124	240	207

Agromet Advisory

General advisory:

Farmers are advised to go for summer ploughing in the dry days as sufficient amount of rainfall was realized in the past and prepare the field for kharif sowing. As increase in day temperature condition likely to prevail in next 5 days, so farmers should cover the soil with natural/straw mulch and advised to strengthen the field bund to conserve present soil moisture.

SUMMER PLOUGHING:

In dry sunny days, go for summer ploughing at places receiving soaking rainfall to reduce weed infestation, to destroy eggs and larvae of hibernated soil insects and to increase soil profile recharge.

GREEN MANURING:

Optimum time for sowing of green manuring crops like Sesbania and sunhemp in the areas receiving convectional rain. Green manuring crops are sown in medium and low land condition at 6-8 weeks before transplanting of rice

Summer Maize:

Where Maize is a standing crop now, at their critical growth stage of soil moisture/irrigation i.e tasseling to silking stage to grain formation which stage demands high water/soil moisture content due to high LAI. Water deficit at that stage will negatively affect the fertilization, grain number per cob, and as a result, the final yield of maize is reduced, So Farmers are advised to maintain proper moisture in the maize crop by giving light irrigation as per requirement.

Vegetables:

To prevent from wilting or partial wilting in brinjal, Chilli and Okra apply Neem or Karanjaa cake and spray (basal + foliar) Tebuconazole fungicide @ 1ml/liter.

Okra: To control **Yellow Vain Mosaic Virus (YVMV)** in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease. Spray Metalaxyl+Mancozeb/Carboxin+Thiram @ 2g/litre to control **rot or soil borne fungal disease.** Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenconazole @ 1ml/lit at 15 days interval to control **Cercospora leaf spot.**

Cucurbits: **Late Blight** may happen due to rising day temperature. There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxchloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Mango: For control of mango **fruit borer** spray Neem oil 3ml + chloripyriphos 1 ml per litre of water along with stickers at marble stage of the fruit. The sprayings should be preferably carried out during evening hours. Spray the foliage with Tebuconazole 50% + Trifloxystrobin 25% WG @ 1g/lit to control **Anthracnose.**

Animal husbandry:

As THI likely to be above 72, to escape dairy animals from thermal stress, make necessary arrangement for clean and sufficient drinking water with sufficient fodder crops for the animals. Keep farm animals indoor during the day time period.

DrRanjan Kumar Mohanta
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK



(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-496

Date- 21/05/2024

AGROMET ADVISORY BULLETIN FOR TANGI-CHOUDWAR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 5mm and district received average 01 mm rainfall during the last week. The mean daily maximum and minimum temperature was 37.6°C (+0.45 from normal) and 26.1°C (+1.14 from normal), respectively. The mean relative humidity was 43-97%.

Forecasted period: Today(8.30IST) to 26thMay(8.30IST)2024

Weather advisory: Very Light to light rainfall/thundershower is forecasted for today to, Thursday and Tuesday and dry weather is forecasted for other 1 day in the block. The day temperature will be 37-41°C and night temperature will be 29-31°C for upcoming 5 days. Maximum and minimum relative humidity will be 74-80% and 40-50% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Tangi-Choudwar (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	21/05/2024	22/05/2024	23/05/2024	24/05/2024	25/05/2024
Rainfall(mm)	4	2	6	0	7
T-Max(°C)	38	37	37	39	41
T-Min(°C)	30	30	29	31	31
Cloud Cover(octa)	0	0	2	7	8
RH Max (%)	79	79	80	79	74
RH Min (%)	50	48	49	40	40
Wind Speed (Kmph)	20	17	13	18	21
Wind Direction (deg)	198	180	158	225	207

Agromet Advisory

General advisory:

Farmers are advised to go for summer ploughing in the dry days as sufficient amount of rainfall was realized in the past and prepare the field for kharif sowing. As increase in day temperature condition likely to prevail in next 5 days, so farmers should cover the soil with natural/straw mulch and advised to strengthen the field bund to conserve present soil moisture.

SUMMER PLOUGHING:

In dry sunny days, go for summer ploughing at places receiving soaking rainfall to reduce weed infestation, to destroy eggs and larvae of hibernated soil insects and to increase soil profile recharge.

GREEN MANURING:

Optimum time for sowing of green manuring crops like Sesbania and sunhemp in the areas receiving convectional rain. Green manuring crops are sown in medium and low land condition at 6-8 weeks before transplanting of rice

Summer Maize:

Where Maize is a standing crop now, at their critical growth stage of soil moisture/irrigation i.e tasseling to silking stage to grain formation which stage demands high water/soil moisture content due to high LAI. Water deficit at that stage will negatively affect the fertilization, grain number per cob, and as a result, the final yield of maize is reduced. So Farmers are advised to maintain proper moisture in the maize crop by giving light irrigation as per requirement.

Vegetables:

To prevent from wilting or partial wilting in brinjal, Chilli and Okra apply Neem or Karanjaa cake and spray (basal + foliar) Tebuconazole fungicide @ 1ml/liter.

Okra: To control Yellow Vain Mosaic Virus (YVMV) in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease. Spray Metalaxyl+Mancozeb/Carboxin+Thiram @ 2g/litre to control rot or soil borne fungal disease. Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenoconazole @ 1ml/lit at 15 days interval to control Cercospora leaf spot.

Cucurbits: Late Blight may happen due to rising day temperature. There are chances of infestation of Downy Mildew in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Present weather may aggravate fruit and vine rot (symptoms: leaf spot, leaf blight, stem rot, fruit rot) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxychloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Mango: For control of mango fruit borer spray Neem oil 3ml + chloripyriphos 1 ml per litre of water along with stickers at marble stage of the fruit. The spraying should be preferably carried out during evening hours. Spray the foliage with Tebuconazole 50% + Trifloxystrobin 25% WG @ 1g/lit to control Anthracnose.

Animal husbandry:

As THI likely to be above 72, to escape dairy animals from thermal stress, make necessary arrangement for clean and sufficient drinking water with sufficient fodder crops for the animals. Keep farm animals indoor during the day time period.





**GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK**



**(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com**

**Bulletin No.-496
Date- 21/05/2024**

AGROMET ADVISORY BULLETIN FOR TIGIRIA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 0mm and district received average 01 mm rainfall during the last week. The mean daily maximum and minimum temperature was 37.6°C (+0.45 from normal) and 26.1°C (+1.14 from normal), respectively. The mean relative humidity was 43-97%.

Forecasted period: Today(8.30IST) to 26thMay(8.30IST)2024

Weather advisory: Very Light to light rainfall/thundershower is forecasted for today to, Thursday and Tuesday and dry weather is forecasted for other 1 day in the block. The day temperature will be 37-39°C and night temperature will be 25-30°C for upcoming 5 days. Maximum and minimum relative humidity will be 54-79% and 24-29% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Tigiria (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	21/05/2024	22/05/2024	23/05/2024	24/05/2024	25/05/2024
Rainfall(mm)	3	2	6	0	4
T-Max(°C)	39	39	38	39	37
T-Min(°C)	26	25	25	27	30
Cloud Cover(octa)	3	3	3	5	8
RH Max (%)	79	77	78	71	54
RH Min (%)	26	28	29	24	28
Wind Speed (Kmph)	17	15	13	10	9
Wind Direction (deg)	193	159	159	165	199

Agro met Advisory

General advisory:

Farmers are advised to go for summer ploughing in the dry days as sufficient amount of rainfall was realized in the past and prepare the field for kharif sowing. As increase in day temperature condition likely to prevail in next 5 days, so farmers should cover the soil with natural/straw mulch and advised to strengthen the field bund to conserve present soil moisture.

SUMMER PLOUGHING:

In dry sunny days, go for summer ploughing at places receiving soaking rainfall to reduce weed infestation, to destroy eggs and larvae of hibernated soil insects and to increase soil profile recharge.

GREEN MANURING:

Optimum time for sowing of green manuring crops like Sesbania and sunhemp in the areas receiving convectional rain. Green manuring crops are sown in medium and low land condition at 6-8 weeks before transplanting of rice

Summer Maize:

Where Maize is a standing crop now, at their critical growth stage of soil moisture/irrigation i.e tasseling to silking stage to grain formation which stage demands high water/soil moisture content due to high LAI. Water deficit at that stage will negatively affect the fertilization, grain number per cob, and as a result, the final yield of maize is reduced, So Farmers are advised to maintain proper moisture in the maize crop by giving light irrigation as per requirement.

Vegetables:

To prevent from wilting or partial wilting in brinjal, Chilli and Okra apply Neem or Karanjaa cake and spray (basal + foliar) Tebuconazole fungicide @ 1ml/litre.

Okra: To control **Yellow Vain Mosaic Virus (YVMV)** in Okra apply Neem oil (10000 ppm) @ 20 ml/ 10 liter at dawn or use yellow sticky trap to control this disease. Spray Metalaxyl+Mancozeb/Carboxin+Thiram @ 2g/litre to control **rot or soil borne fungal disease**. Foliar spraying of Trifloxystrobin 25% + tebuconazole 50% WG @ 1g/lit or Difenconazole @ 1ml/lit at 15 days interval to control **Cercospora leaf spot**.

Cucurbits: **Late Blight** may happen due to rising day temperature. There are chances of infestation of **Downy Mildew** in cucurbit vegetables. To manage this disease in cucurbits spray Copper Hydroxide 77 % WP 12 @ 3-gram/litre or Carbendazim 12 % + Mancozeb 63 % WP @ 2-gram/litre.

Pointed gourd is at flowering and fruiting stage. Present weather may aggravate fruit and vine rot (symptoms: **leaf spot, leaf blight, stem rot, fruit rot**) of pointed gourd. If observe remove damaged fruit and vines from the field and should not be kept in and around the field. Spray alternately Copper Oxchloride 50% WP @ 3 g/litre of water and Metalaxyl 8% + Mancozeb 64% @ 2.5 g/litre of water in clear and dry weather.

Fruit:

Mango: For control of mango **fruit borer** spray Neem oil 3ml + chloripyriphos 1 ml per litre of water along with stickers at marble stage of the fruit. The spraying should be preferably carried out during evening hours. Spray the foliage with Tebuconazole 50% + Trifloxystrobin 25% WG @ 1g/lit to control **Anthraxnose**.

Animal husbandry:

As THI likely to be above 72, to escape dairy animals from thermal stress, make necessary arrangement for clean and sufficient drinking water with sufficient fodder crops for the animals. Keep farm animals indoor during the day time period.











GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-496
Date- 21/05/2024

				
MEGHDOOT APP	DAMINI APP	FACEBOOK (Damu Cuttack)	BLOCK SPECIFIC WHAT'S APP GROUPS	@DAMUCUTTACK

- Kindly install and use **“Meghdoot mobile App”** for Cuttack district weather forecast, warnings and weather based agromet advisory services.
<https://play.google.com/store/apps/details?id=com.aas.meghdoot>
- Kindly install and use **“Damini mobile App”** for location specific for lightening forecast and warnings.
<https://play.google.com/store/apps/details?id=com.lightening.live.damini>
- Follow the **block specific whats app groups** and join in your block wise groups for getting weather based agromet advisory services twice in a week (Tuesday and Friday) and other weather warning based farm management advisories.
- Follow the **Official Website links** to get the weather based agromet advisory service bulletins (**English & Odia**) twice in a week (Tuesday and Friday).

1	ICAR-NRRI, Cuttack, Odisha	https://icar-nrri.in/agromate/	
2	Agrimet division, IMD, Pune	https://imdagrmet.gov.in/imdproject/AGIndex.php	