




# *Cuttack District Block Wise Agro-met Advisory Service Bulletin*

*Friday, 14<sup>th</sup> June, 2024  
 (For the period 14<sup>th</sup> to 18<sup>th</sup> June, 2024)*



*Issued by  
 District Agro-met Unit,  
 KrishiVigyan Kendra Cuttack*





**STANDARD PRECIPITATION INDEX AND NDVI MAP**

Standard Precipitation Index (SPI)	Normalized difference vegetation Index (NDVI)
<p>MINISTRY OF EARTH SCIENCES INDIA METEOROLOGICAL DEPARTMENT HYDROMET SECTION, CRS PUNE</p> <p>STANDARDIZED PRECIPITATION INDEX FOR THE PERIOD 09 MAY - 05 JUNE 2024</p> <p>SPI VALUES</p> <ul style="list-style-type: none"> <li>0 to 0.49</li> <li>0.50 to 0.99 (Extremely Dry) 0% to 10% Area</li> <li>1.00 to 1.49 (Very Dry) 20% to 30% Area</li> <li>1.50 to 1.99 (Dry) 40% to 50% Area</li> <li>2.00 to 2.49 (Moderately Dry) 30% to 40% Area</li> <li>2.50 to 2.99 (Slightly Dry) 20% to 30% Area</li> <li>3.00 to 3.49 (Near Normal) 10% to 20% Area</li> <li>3.50 to 3.99 (Slightly Wet) 10% to 20% Area</li> <li>4.00 to 4.49 (Moderately Wet) 10% to 20% Area</li> <li>4.50 to 4.99 (Very Wet) 10% to 20% Area</li> <li>5.00 to 5.49 (Extremely Wet) 10% to 20% Area</li> </ul>	<p>NOAA/VIIRS/BLENDED NDVI Composite ending on Week no 23 (03.06.2024 to 09.06.2024) over Agricultural region</p> <p align="center"><b>Odisha</b></p> <ul style="list-style-type: none"> <li>-0.9999 - 0</li> <li>0 - 0.05</li> <li>0.05 - 0.1</li> <li>0.1 - 0.15</li> <li>0.15 - 0.2</li> <li>0.2 - 0.25</li> <li>0.25 - 0.3</li> <li>0.3 - 0.35</li> <li>0.35 - 0.4</li> <li>0.4 - 0.45</li> <li>0.45 - 0.5</li> <li>0.5 - 0.55</li> <li>0.55 - 0.6</li> <li>&gt; 0.6</li> </ul> <p>Bare soil, rocks, clouds, rain, snow, etc.</p> <p>Moderate</p> <p>Good</p> <p>Very good</p> <p>Agriculture vigour is moderate at few places of the state.</p>
<p><i>As per SPI value (cumulative 4weeks -09May to 05June 2024), mildly wet condition was there in Cuttack district.</i></p>	<p><i>According to NDVI map (03.06.2024 to 09.06.2024) the agriculture vigor was Moderate over Narasinghpur, Badamba and Banki-Dampada blocks and light over maximum blocks of Cuttack district.</i></p>



<b>LEGEND (Medium range block level rainfall forecast)</b>		
No Rainfall	00 mm	
Very light rainfall	01-05 mm	
Light rainfall	06-10 mm	
Light to moderate rainfall	11-30 mm	
Moderate to heavy rainfall	31-65 mm	
Heavy to very heavy rainfall	66-115 mm	
Extremely heavy rainfall	>116 mm	

<b>LEGEND (Past week temperature )</b>		
Color code	Temperature	Departure
Blue	Below Normal	< 2°C from normal
Green	Normal	-2°C to +2°C from normal
Yellow	Above Normal	>2°C from normal








**Past week (07-13 June, 2023) cumulative rainfall and its departure% from normal and (07-13 June) average maximum and minimum temperature for Cuttack district**



**Medium range (Today (8.30IST) to Wednesday (8.30IST)) Block level Rainfall (mm) forecast for Cuttack district**





**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**



(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-503

Date- 14/06/2024

**Weather (Past and Forecasted) based Disease and pest in Major Crops and vegetables**

Sl No	Crop	Stage and condition of the crop	Anticipated Disease and pest	
1	Jute	Sowing		Weed
2	Brinjal	Fruiting		Phomopsis Blight
				Fruit and shoot borer
3	Tomato	Fruiting		Bacterial wilt
				Seedling rot
4	Cowpea			Rust
5	Cucurbits	Flowering to Fruiting stage		Downy mildew
6	Okra	Fruiting		YMV
7	Banana			Banana leaf spot/sigatoka
				Panama wilt
8	Papaya	Fruting		phytopthera crown rot or foot rot





**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**



(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-503  
Date- 14/06/2024

**AGROMET ADVISORY BULLETIN FOR ATHAGAD**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 1.3mm and district received average 1 mm rainfall during the last week. The mean daily maximum and minimum temperature was 38.47°C (+2.5 from normal) and 26.5°C (+1.5 from normal), respectively. The mean relative humidity was 72-83%.

**Forecasted period: Today(8.30IST) to 19<sup>th</sup> June (8.30IST)2024**

**Weather advisory:** *Very light to light rain/Thundershower is forecasted for upcoming 5days in the block.* The day temperature will be 33-37°C and night temperature will be 25-31°C for upcoming 5 days. Maximum and minimum relative humidity will be 51-71% and 22-30% respectively. **There will be clear to partly cloudy sky for next 5days.**

Block	Athagad (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	14/06/2024	15/06/2024	16/06/2024	17/06/2024	18/06/2024
Rainfall (mm)	1	2	2	5	7
T-Max (°C)	37	34	36	33	35
T-Min (°C)	29	31	28	27	25
Cloud Cover (octa)	6	6	6	7	8
RH Max (%)	64	51	63	71	67
RH Min (%)	22	27	26	30	30
Wind Speed (kmph)	18	14	15	16	18
Wind Direction (deg)	200	214	206	205	212

**Agromet Advisory**

**General advisory:** **Farmer should wait for sufficient amount of rainfall (75-100 mm in a week) for sowing of Karif Rice.**

**Kharif rice:**

Sowing of rice in DSR method should be completed as soon as possible as rainfall is forecasted in between upcoming 5 days. Sowing should be started soon after receive of good amount of early monsoon showers (75-100 mm in a week) where direct seeding is practiced. Incorporate FYM or compost @ 2000 Kg/acre. Apply 27 kg DAP and 20 kg MoP per acre during last ploughing. Before sowing seed should be treated with Trichoderma viride @ 10 g or Carbendazim 50% WP @ 1.5 g/kg seed.

**Transplanting rice:**

After receive of sufficient rainfall, farmers are advised for sowing of dhanicha seeds @ 12 kg/acre prior to the transplanting of paddy so that the dhanicha crop will be 3-4 weeks old before 10-15 days of transplanting of rice. Apply all phosphatic fertilizers of Kharif rice to this green manuring crop in advance. Farmers who are sowing paddy nurseries are advised to treat the seeds with carbendizem @ 3 g/kg seed followed by shade drying for 24 hours before sowing in case of dry nursery and in case of wet nursery seeds should be treated with carbendizem @ 1 g/l of water by soaking for 24 hrs and another 24 hrs for sprouting and the seeds should be sown in wet nurseries.

**Maize:** It is time for sowing of maize for green cob purpose under irrigated condition Grow high yielding varieties like Navjot, Shakti, Decan-107, Decan-109, Use 6 kg seeds/acre. Apply one third of N and all P & K of the recommended dose of 50:24:24 kg. NPK/acre. Sow the seeds in lines with spacing of 45-60X20 cm at 5 cm soil depth.

**VEGETABLES AND FRUITS:**

- **Fruit and shoot borer** in **Brinjal**- Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@300g/acre at 15 days interval.
- Due to Past rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- After cessation of rain, **downy mildew disease** will appear in **cucurbits**. To control it, go for alternate spray of Metalaxyl+Mancozeb@2g/litre and Azoxystrobin+Tebuconazole @0.6g/litre of water at 10 days interval.
- **Rust** disease is likely to appear in **Cowpea**. Spray Chlorothalonil@2g/litre of water.
- **Wilting** may observe in **tomato and brinjal** field, to control it drenching with Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water.
- Current weather favours spread of **YMV** in **Ladies finger**. In case of appearance of diseases, rogue out affected plants and spray Thiometoxam 25WG @ 60g/acre.

**Fruit:**

**Banana:** For **leaf spot/sigatoka** as a precautionary measure spray carbendizim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalonil @ 2.0 g/ litre of water. Drenching soil with Metiram 75% WG @1.5g/4 lit or carbendazim 12%+ mancozeb 63% @ 2g/lit to prevent **Panama wilt**.

**Mango:** Harvest the mature mango fruits during morning or evening hours. Keep the harvested fruits in shade to prevent from spongy tissue disease and heat.

**Papaya:** After rain due to water logging there may be increase in the infestation of **phytophthora crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

**Animal Husbandry:** Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination. Keep the animal shed clean and dry.





**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**



(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-503  
Date- 14/06/2024

**AGROMET ADVISORY BULLETIN FOR BADAMBA**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 0.7 mm and district received average 1 mm rainfall during the last week. **The mean daily maximum and minimum temperature was 38.47°C (+2.5 from normal) and 26.5°C (+1.5 from normal), respectively.** The mean relative humidity was 72-83%.

**Forecasted period: Today(8.30IST) to 19<sup>th</sup> June (8.30IST)2024**

**Weatheradvisory:: Very light to light rain/Thundershower is forecasted for upcoming 5days in the block.** The day temperature will be 34-38°C and night temperature will be 24-29°C for upcoming 5 days. Maximum and minimum relative humidity will be 51-70% and 22-30% respectively. **There will be clear to partly cloudy sky for next 5days.**

Block	Badamba (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	14/06/2024	15/06/2024	16/06/2024	17/06/2024	18/06/2024
Rainfall(mm)	1	2	2	5	7
T-Max(°C)	38	34	36	34	36
T-Min(°C)	29	29	28	26	24
Cloud Cover(octa)	7	6	6	7	8
RH Max (%)	57	51	57	66	70
RH Min (%)	22	29	26	28	30
Wind Speed (Kmph)	15	12	12	13	15
Wind Direction (deg)	200	270	207	205	208

**Agromet Advisory**

**General advisory: Farmer should wait for sufficient amount of rainfall (75-100 mm in a week) for sowing of Karif Rice.**

**Kharif rice:**

Sowing of rice in DSR method should be completed as soon as possible as rainfall is forecasted in between upcoming 5 days. Sowing should be started soon after receive of good amount of early monsoon showers (75-100 mm in a week) where direct seeding is practiced. Incorporate FYM or compost @ 2000 Kg/acre. Apply 27 kg DAP and 20 kg MoP per acre during last ploughing. Before sowing seed should be treated with Trichoderma viride @ 10 g or Carbendazim 50% WP @ 1.5 g/kg seed.

**Transplanting rice:**

After receive of sufficient rainfall, farmers are advised for sowing of dhanicha seeds @ 12 kg/acre prior to the transplanting of paddy so that the dhanicha crop will be 3-4 weeks old before 10-15 days of transplanting of rice. Apply all phosphatic fertilizers of Kharif rice to this green manuring crop in advance. Farmers who are sowing paddy nurseries are advised to treat the seeds with carbendizem @3 g/kg seed followed by shade drying for 24 hours before sowing in case of dry nursery and in case of wet nursery seeds should be treated with carbendizem @ 1 g/l of water by soaking for 24 hrs and another 24 hrs for sprouting and the seeds should be sown in wet nurseries.

**Maize:** It is time for sowing of maize for green cob purpose under irrigated condition Grow high yielding varieties like Navjot, Shakti, Decan-107, Decan-109, Use 6 kg seeds/acre. Apply one third of N and all P & K of the recommended dose of 50:24:24 kg. NPK/acre. Sow the seeds in lines with spacing of 45-60X20 cm at 5 cm soil depth.

**VEGETABLES AND FRUITS:**

- **Fruit and shoot borer** in **Brinjal**- Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@300g/acre at 15 days interval.
- Due to Past rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- After cessation of rain, **downy mildew disease** will appear in **cucurbits**. To control it, go for alternate spray of Metalaxyl+Mancozeb@2g/litre and Azoxystrobin+Tebuconazole @0.6g/litre of water at 10 days interval.
- **Rust** disease is likely to appear in **Cowpea**. Spray Chlorothalonil@2g/litre of water.
- **Wilting** may observe in **tomato and brinjal** field, to control it drenching with Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water.
- Current weather favours spread of **YMV** in **Ladies finger**. In case of appearance of diseases, rogue out affected plants and spray Thiometoxam 25WG @ 60g/acre.

**Fruit:**

**Banana:** For **leaf spot/sigatoka** as a precautionary measure spray carbendizim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalonil@ 2.0 g/ litre of water. Drenching soil with Metiram 75% WG @1.5g/4 lit or carbendazim 12%+ mancozeb 63% @ 2g/lit to prevent **Panama wilt**.

**Mango:** Harvest the mature mango fruits during morning or evening hours. Keep the harvested fruits in shade to prevent from spongy tissue disease and heat.

**Papaya:** After rain due to water logging there may be increase in the infestation of **phytophthora crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

**Animal Husbandry:** Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination. Keep the animal shed clean and dry.

Dr Ranjan Kumar Mohanta  
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena  
SMS (Agrometeorology)  
Contact No-9861445080/7008453427



**GRAMIN KRISHI MAUSAM SEWA  
INDIA METEOROLOGICAL DEPARTMENT  
DISTRICT AGROMET UNIT  
KRISHI VIGYAN KENDRA CUTTACK**



(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-503  
Date- 14/06/2024

**AGROMET ADVISORY BULLETIN FOR BANKI**

**District Past week average weather condition and Block past week average Rainfall condition**

The district received average 1 mm rainfall during the last week. The mean daily maximum and minimum temperature was 38.47°C (+2.5 from normal) and 26.5°C (+1.5 from normal), respectively. The mean relative humidity was 72-83%.

**Forecasted period: Today(8.30IST) to 19<sup>th</sup> June (8.30IST)2024**

**Weather advisory:: Very light to light rain/Thundershower is forecasted for upcoming 5 days in the block.** The day temperature will be 34-38°C and night temperature will be 25-31°C for upcoming 5 days. Maximum and minimum relative humidity will be 50-69% and 22-32% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Banki (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	14/06/2024	15/06/2024	16/06/2024	17/06/2024	18/06/2024
Rainfall(mm)	1	3	3	7	7
T-Max(°C)	38	36	36	34	36
T-Min(°C)	29	31	28	27	25
Cloud Cover(octa)	7	6	6	7	8
RH Max (%)	64	50	61	69	67
RH Min (%)	22	27	26	28	32
Wind Speed (Kmph)	18	13	15	16	18
Wind Direction (deg)	203	225	203	203	212

**Agromet Advisory**

**General advisory: Farmer should wait for sufficient amount of rainfall (75-100 mm in a week) for sowing of Kharif Rice.**

**Kharif rice:**

Sowing of rice in DSR method should be completed as soon as possible as rainfall is forecasted in between upcoming 5 days. Sowing should be started soon after receive of good amount of early monsoon showers (75-100 mm in a week) where direct seeding is practiced. Incorporate FYM or compost @ 2000 Kg/acre. Apply 27 kg DAP and 20 kg MoP per acre during last ploughing. Before sowing seed should be treated with Trichoderma viride @ 10 g or Carbendazim 50% WP @ 1.5 g/kg seed.

**Transplanting rice:**

After receive of sufficient rainfall, farmers are advised for sowing of dhanicha seeds @ 12 kg/acre prior to the transplanting of paddy so that the dhanicha crop will be 3-4 weeks old before 10-15 days of transplanting of rice. Apply all phosphatic fertilizers of Kharif rice to this green manuring crop in advance. Farmers who are sowing paddy nurseries are advised to treat the seeds with carbendizem @3 g/kg seed followed by shade drying for 24 hours before sowing in case of dry nursery and in case of wet nursery seeds should be treated with carbendizem @ 1 g/l of water by soaking for 24 hrs and another 24 hrs for sprouting and the seeds should be sown in wet nurseries.

**Maize:** It is time for sowing of maize for green cob purpose under irrigated condition Grow high yielding varieties like Navjot, Shakti, Decan-107, Decan-109, Use 6 kg seeds/acre. Apply one third of N and all P & K of the recommended dose of 50:24:24 kg. NPK/acre. Sow the seeds in lines with spacing of 45-60X20 cm at 5 cm soil depth.

**VEGETABLES AND FRUITS:**

- Fruit and shoot borer** in **Brinjal**- Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@300g/acre at 15 days interval.
- Due to Past rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- After cessation of rain, **downy mildew disease** will appear in **cucurbits**. To control it, go for alternate spray of Metalaxyl+Mancozeb@2g/litre and Azoxystrobin+Tebuconazole @0.6g/litre of water at 10 days interval.
- Rust** disease is likely to appear in **Cowpea**. Spray Chlorothalonil@2g/litre of water.
- Wilting** may observe in **tomato and brinjal** field, to control it drenching with Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water.
- Current weather favours spread of **YMV** in **Ladies finger**. In case of appearance of diseases, rogue out affected plants and spray Thiometoxam 25WG @ 60g/acre.

**Fruit:**

**Banana:** For **leaf spot/sigatoka** as a precautionary measure spray carbendizim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalonil @ 2.0 g/ litre of water. Drenching soil with Metiram 75% WG @1.5g/4 lit or carbendazim 12%+ mancozeb 63% @ 2g/lit to prevent **Panama wilt**.

**Mango:** Harvest the mature mango fruits during morning or evening hours. Keep the harvested fruits in shade to prevent from spongy tissue disease and heat.

**Papaya:** After rain due to water logging there may be increase in the infestation of **phytophthora crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

**Animal Husbandry:** Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination. Keep the animal shed clean and dry.





**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**



(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-503  
Date- 14/06/2024

**AGROMET ADVISORY BULLETIN FOR BANKI-DAMPARA**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 0.6 mm and district received average 1 mm rainfall during the last week. The mean daily maximum and minimum temperature was 38.47°C (+2.5 from normal) and 26.5°C (+1.5 from normal), respectively. The mean relative humidity was 72-83%.

**Forecasted period: Today(8.30IST) to 19<sup>th</sup> June (8.30IST)2024**

**Weather Advisory:** Very light to light rain/Thundershower is forecasted for upcoming 5 days in the block. The day temperature will be 34-38°C and night temperature will be 26-31°C for upcoming 5 days. Maximum and minimum relative humidity will be 51-70% and 22-33% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Banki-Dampara (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	14/06/2024	15/06/2024	16/06/2024	17/06/2024	18/06/2024
Rainfall(mm)	2	3	3	7	8
T-Max(°C)	38	36	36	34	36
T-Min(°C)	29	31	28	27	26
Cloud Cover(octa)	7	6	6	7	8
RH Max (%)	65	51	62	70	66
RH Min (%)	22	26	26	28	33
Wind Speed (Kmph)	18	13	15	17	19
Wind Direction (deg)	205	225	199	203	212

**Agromet Advisory**

**General advisory:** Farmer should wait for sufficient amount of rainfall (75-100 mm in a week) for sowing of Karif Rice.

**Kharif rice:**

Sowing of rice in DSR method should be completed as soon as possible as rainfall is forecasted in between upcoming 5 days. Sowing should be started soon after receive of good amount of early monsoon showers (75-100 mm in a week) where direct seeding is practiced. Incorporate FYM or compost @ 2000 Kg/acre. Apply 27 kg DAP and 20 kg MoP per acre during last ploughing. Before sowing seed should be treated with Trichoderma viride @ 10 g or Carabendazim 50% WP @ 1.5 g/kg seed.

**Transplanting rice:**

After receive of sufficient rainfall, farmers are advised for sowing of dhanicha seeds @ 12 kg/acre prior to the transplanting of paddy so that the dhanicha crop will be 3-4 weeks old before 10-15 days of transplanting of rice. Apply all phosphatic fertilizers of Kharif rice to this green manuring crop in advance. Farmers who are sowing paddy nurseries are advised to treat the seeds with carbendizem @3 g/kg seed followed by shade drying for 24 hours before sowing in case of dry nursery and in case of wet nursery seeds should be treated with carbendizem @ 1 g/l of water by soaking for 24 hrs and another 24 hrs for sprouting and the seeds should be sown in wet nurseries.

**Maize:** It is time for sowing of maize for green cob purpose under irrigated condition Grow high yielding varieties like Navjot, Shakti, Decan-107, Decan-109, Use 6 kg seeds/acre. Apply one third of N and all P & K of the recommended dose of 50:24:24 kg. NPK/acre. Sow the seeds in lines with spacing of 45-60X20 cm at 5 cm soil depth.

**VEGETABLES AND FRUITS:**

- **Fruit and shoot borer** in **Brinjal**- Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@300g/acre at 15 days interval.
- Due to Past rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- After cessation of rain, **downy mildew disease** will appear in **cucurbits**. To control it, go for alternate spray of Metalaxyl+Mancozeb@2g/litre and Azoxystrobin+Tebuconazole @0.6g/litre of water at 10 days interval.
- **Rust** disease is likely to appear in **Cowpea**. Spray Chlorothalonil@2g/litre of water.
- **Wilting** may observe in **tomato and brinjal** field, to control it drenching with Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water.
- Current weather favours spread of **YMV** in **Ladies finger**. In case of appearance of diseases, rogue out affected plants and spray Thiometoxam 25WG @ 60g/acre.

**Fruit:**

**Banana:** For **leaf spot/sigatoka** as a precautionary measure spray carbendizim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalonil@ 2.0 g/ litre of water. Drenching soil with Metiram 75% WG @1.5g/4 lit or carbendazim 12%+ mancozeb 63% @ 2g/lit to prevent **Panama wilt**.

**Mango:** Harvest the mature mango fruits during morning or evening hours. Keep the harvested fruits in shade to prevent from spongy tissue disease and heat.

**Papaya:** After rain due to water logging there may be increase in the infestation of **phytophthora crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

**Animal Husbandry:** Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination. Keep the animal shed clean and dry.







**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**



(A Unit of ICAR-National Rice Research Institute)  
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-503  
 Date- 14/06/2024

**AGROMET ADVISORY BULLETIN FOR BARANGA**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 2.6mm and district received average 1 mm rainfall during the last week. **The mean daily maximum and minimum temperature was 38.47°C (+2.5 from normal) and 26.5°C (+1.5 from normal), respectively.** The mean relative humidity was 72-83%.

**Forecasted period: Today(8.30IST) to 19<sup>th</sup> June (8.30IST)2024**

**Weather advisory: Very light to light rain/Thundershower is forecasted for upcoming 5 days in the block.** The day temperature will be 34-38°C and night temperature will be 27-31°C for upcoming 5 days. Maximum and minimum relative humidity will be 57-73% and 22-31% respectively. **There will be clear to partly cloudy sky for next 5 days.**

Block	Baranga (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	14/06/2024	15/06/2024	16/06/2024	17/06/2024	18/06/2024
Rainfall(mm)	2	3	3	7	6
T-Max(°C)	38	37	37	34	35
T-Min(°C)	29	31	28	27	27
Cloud Cover	6	5	5	7	8
RH Max (%)	73	57	67	73	68
RH Min (%)	22	23	25	31	37
Wind Speed (Kmph)	22	17	20	22	25
Wind Direction (deg)	205	205	207	207	212

**Agromet Advisory**

**General advisory: Farmer should wait for sufficient amount of rainfall (75-100 mm in a week) for sowing of Karif Rice.**

**Kharif rice:**

Sowing of rice in DSR method should be completed as soon as possible as rainfall is forecasted in between upcoming 5 days. Sowing should be started soon after receive of good amount of early monsoon showers (75-100 mm in a week) where direct seeding is practiced. Incorporate FYM or compost @ 2000 Kg/acre. Apply 27 kg DAP and 20 kg MoP per acre during last ploughing. Before sowing seed should be treated with Trichoderma viride @ 10 g or Carbendazim 50% WP @ 1.5 g/kg seed.

**Transplanting rice:**

After receive of sufficient rainfall, farmers are advised for sowing of dhanicha seeds @ 12 kg/acre prior to the transplanting of paddy so that the dhanicha crop will be 3-4 weeks old before 10-15 days of transplanting of rice. Apply all phosphatic fertilizers of Kharif rice to this green manuring crop in advance. Farmers who are sowing paddy nurseries are advised to treat the seeds with carbendizem @3 g/kg seed followed by shade drying for 24 hours before sowing in case of dry nursery and in case of wet nursery seeds should be treated with carbendizem @ 1 g/l of water by soaking for 24 hrs and another 24 hrs for sprouting and the seeds should be sown in wet nurseries.

**Maize:** It is time for sowing of maize for green cob purpose under irrigated condition Grow high yielding varieties like Navjot, Shakti, Decan-107, Decan-109, Use 6 kg seeds/acre. Apply one third of N and all P & K of the recommended dose of 50:24:24 kg. NPK/acre. Sow the seeds in lines with spacing of 45-60X20 cm at 5 cm soil depth.

**VEGETABLES AND FRUITS:**

- **Fruit and shoot borer** in **Brinjal**- Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@300g/acre at 15 days interval.
- Due to Past rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- After cessation of rain, **downy mildew disease** will appear in **cucurbits**. To control it, go for alternate spray of Metalaxyl+Mancozeb@2g/litre and Azoxystrobin+Tebuconazole @0.6g/litre of water at 10 days interval.
- **Rust** disease is likely to appear in **Cowpea**. Spray Chlorothalonil@2g/litre of water.
- **Wilting** may observe in **tomato and brinjal** field, to control it drenching with Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water.
- Current weather favours spread of **YMV** in **Ladies finger**. In case of appearance of diseases, rogue out affected plants and spray Thiometoxam 25WG @ 60g/acre.

**Fruit:**

**Banana:** For **leaf spot/sigatoka** as a precautionary measure spray carbendizim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalonil@ 2.0 g/ litre of water. Drenching soil with Metiram 75% WG @1.5g/4 lit or carbendazim 12%+ mancozeb 63% @ 2g/lit to prevent **Panama wilt**.

**Mango:** Harvest the mature mango fruits during morning or evening hours. Keep the harvested fruits in shade to prevent from spongy tissue disease and heat.

**Papaya:** After rain due to water logging there may be increase in the infestation of **phytopthera crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

**Animal Husbandry:** Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination. Keep the animal shed clean and dry.





**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**



(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-503  
Date- 14/06/2024

**AGROMET ADVISORY BULLETIN FOR CUTTACK SADAR**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 4.1mm and district received average 1 mm rainfall during the last week. **The mean daily maximum and minimum temperature was 38.47°C (+2.5 from normal) and 26.5°C (+1.5 from normal), respectively.** The mean relative humidity was 72-83%.

**Forecasted period: Today(8.30IST) to 19<sup>th</sup> June (8.30IST)2024**

**Weather advisory: Very light to light rain/Thundershower is forecasted for upcoming 5days in the block.** The day temperature will be 33-37°C and night temperature will be 27-31°C for upcoming 5 days. Maximum and minimum relative humidity will be 56-72% and 22-32% respectively. **There will be clear to partly cloudy sky for next 5days.**

Block	Cuttack sadar (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	14/06/2024	15/06/2024	16/06/2024	17/06/2024	18/06/2024
Rainfall(mm)	1	2	2	8	7
T-Max(°C)	37	36	36	33	35
T-Min(°C)	29	31	28	27	27
Cloud Cover(octa)	6	5	5	7	8
RH Max (%)	71	56	65	72	67
RH Min (%)	22	25	25	31	32
Wind Speed (Kmph)	20	16	19	20	22
Wind Direction (deg)	205	225	207	208	212

**Agromet Advisory**

**General advisory: Farmer should wait for sufficient amount of rainfall (75-100 mm in a week) for sowing of Karif Rice.**

**Kharif rice:**

Sowing of rice in DSR method should be completed as soon as possible as rainfall is forecasted in between upcoming 5 days. Sowing should be started soon after receive of good amount of early monsoon showers (75-100 mm in a week) where direct seeding is practiced. Incorporate FYM or compost @ 2000 Kg/acre. Apply 27 kg DAP and 20 kg MoP per acre during last ploughing. Before sowing seed should be treated with Trichoderma viride @ 10 g or Carabendazim 50% WP @ 1.5 g/kg seed.

**Transplanting rice:**

After receive of sufficient rainfall, farmers are advised for sowing of dhanicha seeds @ 12 kg/acre prior to the transplanting of paddy so that the dhanicha crop will be 3-4 weeks old before 10-15 days of transplanting of rice. Apply all phosphatic fertilizers of Kharif rice to this green manuring crop in advance. Farmers who are sowing paddy nurseries are advised to treat the seeds with carbendizem @3 g/kg seed followed by shade drying for 24 hours before sowing in case of dry nursery and in case of wet nursery seeds should be treated with carbendizem @ 1 g/l of water by soaking for 24 hrs and another 24 hrs for sprouting and the seeds should be sown in wet nurseries.

**Maize:** It is time for sowing of maize for green cob purpose under irrigated condition Grow high yielding varieties like Navjot, Shakti, Decan-107, Decan-109, Use 6 kg seeds/acre. Apply one third of N and all P & K of the recommended dose of 50:24:24 kg. NPK/acre. Sow the seeds in lines with spacing of 45-60X20 cm at 5 cm soil depth.

**VEGETABLES AND FRUITS:**

- **Fruit and shoot borer** in **Brinjal**- Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@300g/acre at 15 days interval.
- Due to Past rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- After cessation of rain, **downy mildew disease** will appear in **cucurbits**. To control it, go for alternate spray of Metalaxyl+Mancozeb@2g/litre and Azoxystrobin+Tebuconazole @0.6g/litre of water at 10 days interval.
- **Rust** disease is likely to appear in **Cowpea**. Spray Chlorothalonil@2g/litre of water.
- **Wilting** may observe in **tomato and brinjal** field, to control it drenching with Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water.
- Current weather favours spread of **YMV** in **Ladies finger**. In case of appearance of diseases, rogue out affected plants and spray Thiometoxam 25WG @ 60g/acre.

**Fruit:**

**Banana:** For **leaf spot/sigatoka** as a precautionary measure spray carbendizim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalonil@ 2.0 g/ litre of water. Drenching soil with Metiram 75% WG @1.5g/4 lit or carbendazim 12%+ mancozeb 63% @ 2g/lit to prevent **Panama wilt**.

**Mango:** Harvest the mature mango fruits during morning or evening hours. Keep the harvested fruits in shade to prevent from spongy tissue disease and heat.

**Papaya:** After rain due to water logging there may be increase in the infestation of **phytophthora crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

**Animal Husbandry:** Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination. Keep the animal shed clean and dry.





**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**



(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-503  
Date- 14/06/2024

**AGROMET ADVISORY BULLETIN FOR KANTAPADA**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 0.3mm and district received average 1 mm rainfall during the last week. The mean daily maximum and minimum temperature was 38.47°C (+2.5 from normal) and 26.5°C (+1.5 from normal), respectively. The mean relative humidity was 72-83%.

**Forecasted period: Today(8.30IST) to 19<sup>th</sup> June (8.30IST)2024**

**Weather advisory: Very light to light rain/Thundershower is forecasted for upcoming 5 days in the block.** The day temperature will be 34-39°C and night temperature will be 27-31°C for upcoming 5 days. Maximum and minimum relative humidity will be 61-74% and 22-43% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Kantapada (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	14/06/2024	15/06/2024	16/06/2024	17/06/2024	18/06/2024
Rainfall(mm)	1	2	2	8	7
T-Max(°C)	38	39	37	35	34
T-Min(°C)	29	31	29	28	27
Cloud Cover(octa)	6	5	6	7	8
RH Max (%)	72	61	69	74	70
RH Min (%)	24	22	27	35	43
Wind Speed (Kmph)	24	17	23	26	27
Wind Direction (deg)	207	203	207	203	212

**Agromet Advisory**

**General advisory: Farmer should wait for sufficient amount of rainfall (75-100 mm in a week) for sowing of Kharif Rice.**

**Kharif rice:**

Sowing of rice in DSR method should be completed as soon as possible as rainfall is forecasted in between upcoming 5 days. Sowing should be started soon after receive of good amount of early monsoon showers (75-100 mm in a week) where direct seeding is practiced. Incorporate FYM or compost @ 2000 Kg/acre. Apply 27 kg DAP and 20 kg MoP per acre during last ploughing. Before sowing seed should be treated with Trichoderma viride @ 10 g or Carbendazim 50% WP @ 1.5 g/kg seed.

**Transplanting rice:**

After receive of sufficient rainfall, farmers are advised for sowing of dhanicha seeds @ 12 kg/acre prior to the transplanting of paddy so that the dhanicha crop will be 3-4 weeks old before 10-15 days of transplanting of rice. Apply all phosphatic fertilizers of Kharif rice to this green manuring crop in advance. Farmers who are sowing paddy nurseries are advised to treat the seeds with carbendizem @3 g/kg seed followed by shade drying for 24 hours before sowing in case of dry nursery and in case of wet nursery seeds should be treated with carbendizem @ 1 g/l of water by soaking for 24 hrs and another 24 hrs for sprouting and the seeds should be sown in wet nurseries.

**Maize:** It is time for sowing of maize for green cob purpose under irrigated condition Grow high yielding varieties like Navjot, Shakti, Decan-107, Decan-109, Use 6 kg seeds/acre. Apply one third of N and all P & K of the recommended dose of 50:24:24 kg. NPK/acre. Sow the seeds in lines with spacing of 45-60X20 cm at 5 cm soil depth.

**VEGETABLES AND FRUITS:**

- Fruit and shoot borer** in **Brinjal**- Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@300g/acre at 15 days interval.
- Due to Past rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- After cessation of rain, **downy mildew disease** will appear in **cucurbits**. To control it, go for alternate spray of Metalaxyl+Mancozeb@2g/litre and Azoxystrobin+Tebuconazole @0.6g/litre of water at 10 days interval.
- Rust** disease is likely to appear in **Cowpea**. Spray Chlorothalonil@2g/litre of water.
- Wilting** may observe in **tomato and brinjal** field, to control it drenching with Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water.
- Current weather favours spread of **YMV** in **Ladies finger**. In case of appearance of diseases, rogue out affected plants and spray Thiomethoxam 25WG @ 60g/acre.

**Fruit:**

**Banana:** For **leaf spot/sigatoka** as a precautionary measure spray carbendizim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalonil @ 2.0 g/ litre of water. Drenching soil with Metiram 75% WG @1.5g/4 lit or carbendazim 12%+ mancozeb 63% @ 2g/lit to prevent **Panama wilt**.

**Mango:** Harvest the mature mango fruits during morning or evening hours. Keep the harvested fruits in shade to prevent from spongy tissue disease and heat.

**Papaya:** After rain due to water logging there may be increase in the infestation of **phytophthora crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

**Animal Husbandry:** Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination. Keep the animal shed clean and dry.





**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**



(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-503  
Date- 14/06/2024

**AGROMET ADVISORY BULLETIN FOR MAHANGA**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 0.6mm and district received average 1 mm rainfall during the last week. The mean daily maximum and minimum temperature was 38.47°C (+2.5 from normal) and 26.5°C (+1.5 from normal), respectively. The mean relative humidity was 72-83%.

**Forecasted period: Today(8.30IST) to 19<sup>th</sup> June (8.30IST)2024**

**Weather advisory: Very light to light rain/Thundershower is forecasted for upcoming 5days in the block.** The day temperature will be and it will be 33-38°C and night temperature will be 27-31°C for upcoming 5 days. Maximum and minimum relative humidity will be 59-72% and 23-31% respectively. **There will be clear to partly cloudy sky for next 5days.**

Block	Mahanga (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	14/06/2024	15/06/2024	16/06/2024	17/06/2024	18/06/2024
Rainfall(mm)	1	2	2	8	8
T-Max(°C)	38	37	37	33	34
T-Min(°C)	29	31	29	27	27
Cloud Cover(octal)	5	5	5	7	8
RH Max (%)	70	59	64	72	67
RH Min (%)	23	24	25	31	35
Wind Speed (Kmph)	23	19	22	22	25
Wind Direction (deg)	205	225	212	208	212

**Agromet Advisory**

**General advisory: Farmer should wait for sufficient amount of rainfall (75-100 mm in a week) for sowing of Kharif Rice.**

**Kharif rice:**

Sowing of rice in DSR method should be completed as soon as possible as rainfall is forecasted in between upcoming 5 days. Sowing should be started soon after receive of good amount of early monsoon showers (75-100 mm in a week) where direct seeding is practiced. Incorporate FYM or compost @ 2000 Kg/acre. Apply 27 kg DAP and 20 kg MoP per acre during last ploughing. Before sowing seed should be treated with Trichoderma viride @ 10 g or Carbendazim 50% WP @ 1.5 g/kg seed.

**Transplanting rice:**

After receive of sufficient rainfall, farmers are advised for sowing of dhanicha seeds @ 12 kg/acre prior to the transplanting of paddy so that the dhanicha crop will be 3-4 weeks old before 10-15 days of transplanting of rice. Apply all phosphatic fertilizers of Kharif rice to this green manuring crop in advance. Farmers who are sowing paddy nurseries are advised to treat the seeds with carbendizem @3 g/kg seed followed by shade drying for 24 hours before sowing in case of dry nursery and in case of wet nursery seeds should be treated with carbendizem @ 1 g/l of water by soaking for 24 hrs and another 24 hrs for sprouting and the seeds should be sown in wet nurseries.

**Maize:** It is time for sowing of maize for green cob purpose under irrigated condition Grow high yielding varieties like Navjot, Shakti, Decan-107, Decan-109, Use 6 kg seeds/acre. Apply one third of N and all P & K of the recommended dose of 50:24:24 kg. NPK/acre. Sow the seeds in lines with spacing of 45-60X20 cm at 5 cm soil depth.

**VEGETABLES AND FRUITS:**

- Fruit and shoot borer** in **Brinjal**- Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@300g/acre at 15 days interval.
- Due to Past rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- After cessation of rain, **downy mildew disease** will appear in **cucurbits**. To control it, go for alternate spray of Metalaxyl+Mancozeb@2g/litre and Azoxystrobin+Tebuconazole @0.6g/litre of water at 10 days interval.
- Rust** disease is likely to appear in **Cowpea**. Spray Chlorothalonil@2g/litre of water.
- Wilting** may observe in **tomato and brinjal** field, to control it drenching with Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water.
- Current weather favours spread of **YMV** in **Ladies finger**. In case of appearance of diseases, rogue out affected plants and spray Thiometoxam 25WG @ 60g/acre.

**Fruit:**

**Banana:** For **leaf spot/sigatoka** as a precautionary measure spray carbendizim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalonil @ 2.0 g/ litre of water. Drenching soil with Metiram 75% WG @1.5g/4 lit or carbendazim 12%+ mancozeb 63% @ 2g/lit to prevent **Panama wilt**.

**Mango:** Harvest the mature mango fruits during morning or evening hours. Keep the harvested fruits in shade to prevent from spongy tissue disease and heat.

**Papaya:** After rain due to water logging there may be increase in the infestation of **phytophthora crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

**Animal Husbandry:** Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination. Keep the animal shed clean and dry.





**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**



(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-503  
Date- 14/06/2024

**AGROMET ADVISORY BULLETIN FOR NARASINGHPUR**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 0mm and district received average 1 mm rainfall during the last week. **The mean daily maximum and minimum temperature was 38.47°C (+2.5 from normal) and 26.5°C (+1.5 from normal), respectively.** The mean relative humidity was 72-83%.

**Forecasted period: Today(8.30IST) to 19<sup>th</sup> June (8.30IST)2024**

**Weather advisory: Very light to light rain/Thundershower is forecasted for upcoming 5days in the block.** The day temperature will be 32-37°C and night temperature will be 23-28°C for upcoming 5 days. Maximum and minimum relative humidity will be 53-78% and 21-34% respectively. **There will be clear to partly cloudy sky for next 5days.**

Block	Narasinghpur (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	14/06/2024	15/06/2024	16/06/2024	17/06/2024	18/06/2024
Rainfall(mm)	2	3	3	5	8
T-Max(°C)	37	33	35	32	37
T-Min(°C)	28	27	27	25	23
Cloud Cover(/octa)	7	6	6	8	8
RH Max (%)	53	55	55	62	78
RH Min (%)	21	31	25	34	27
Wind Speed (Kmph)	11	9	8	8	9
Wind Direction (deg)	203	261	240	225	216

**Agromet Advisory**

**General advisory: Farmer should wait for sufficient amount of rainfall (75-100 mm in a week) for sowing of Karif Rice.**

**Kharif rice:**

Sowing of rice in DSR method should be completed as soon as possible as rainfall is forecasted in between upcoming 5 days. Sowing should be started soon after receive of good amount of early monsoon showers (75-100 mm in a week) where direct seeding is practiced. Incorporate FYM or compost @ 2000 Kg/acre. Apply 27 kg DAP and 20 kg MoP per acre during last ploughing. Before sowing seed should be treated with Trichoderma viride @ 10 g or Carbendazim 50% WP @ 1.5 g/kg seed.

**Transplanting rice:**

After receive of sufficient rainfall, farmers are advised for sowing of dhanicha seeds @ 12 kg/acre prior to the transplanting of paddy so that the dhanicha crop will be 3-4 weeks old before 10-15 days of transplanting of rice. Apply all phosphatic fertilizers of Kharif rice to this green manuring crop in advance. Farmers who are sowing paddy nurseries are advised to treat the seeds with carbendizem @3 g/kg seed followed by shade drying for 24 hours before sowing in case of dry nursery and in case of wet nursery seeds should be treated with carbendizem @ 1 g/l of water by soaking for 24 hrs and another 24 hrs for sprouting and the seeds should be sown in wet nurseries.

**Maize:** It is time for sowing of maize for green cob purpose under irrigated condition Grow high yielding varieties like Navjot, Shakti, Decan-107, Decan-109, Use 6 kg seeds/acre. Apply one third of N and all P & K of the recommended dose of 50:24:24 kg. NPK/acre. Sow the seeds in lines with spacing of 45-60X20 cm at 5 cm soil depth.

**VEGETABLES AND FRUITS:**

- **Fruit and shoot borer** in **Brinjal**- Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@300g/acre at 15 days interval.
- Due to Past rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- After cessation of rain, **downy mildew disease** will appear in **cucurbits**. To control it, go for alternate spray of Metalaxyl+Mancozeb@2g/litre and Azoxystrobin+Tebuconazole @0.6g/litre of water at 10 days interval.
- **Rust** disease is likely to appear in **Cowpea**. Spay Chlorothalonil@2g/litre of water.
- **Wilting** may observe in **tomato and brinjal** field, to control it drenching with Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water.
- Current weather favours spread of **YMV** in **Ladies finger**. In case of appearance of diseases, rogue out affected plants and spray Thiometoxam 25WG @ 60g/acre.

**Fruit:**

**Banana:** For **leaf spot/sigatoka** as a precautionary measure spray carbendizim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalonil@ 2.0 g/ litre of water. Drenching soil with Metiram 75% WG @1.5g/4 lit or carbendazim 12%+ mancozeb 63% @ 2g/lit to prevent **Panama wilt**.

**Mango:** Harvest the mature mango fruits during morning or evening hours. Keep the harvested fruits in shade to prevent from spongy tissue disease and heat.

**Papaya:** After rain due to water logging there may be increase in the infestation of **phytopthera crown rot or foot rot** in papya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

**Animal Husbandry:** Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination. Keep the animal shed clean and dry.





**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**



(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-503  
Date- 14/06/2024

**AGROMET ADVISORY BULLETIN FOR NIALI**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 0.9mm and district received average 1 mm rainfall during the last week. **The mean daily maximum and minimum temperature was 38.47°C (+2.5 from normal) and 26.5°C (+1.5 from normal), respectively.** The mean relative humidity was 72-83%.

**Forecasted period: Today(8.30IST) to 19<sup>th</sup> June (8.30IST)2024**

**Weather advisory: Very light to light rain/Thundershower is forecasted for upcoming 5days in the block.** The day temperature will be 34-40°C and night temperature will be 27-30°C for upcoming 5 days. Maximum and minimum relative humidity will be 65-75% and 26-47% respectively. There will be clear to partly cloudy sky for next 5days.

Block	Niali (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	14/06/2024	15/06/2024	16/06/2024	17/06/2024	18/06/2024
Rainfall(mm)	2	2	3	7	7
T-Max(°C)	38	40	38	35	34
T-Min(°C)	29	30	28	28	27
Cloud Cover(octa)	6	4	5	7	8
RH Max (%)	73	65	72	75	72
RH Min (%)	26	26	30	42	47
Wind Speed(Kmph)	25	18	24	28	29
Wind Direction(deg)	207	207	203	203	212

**Agromet Advisory**

**General advisory: Farmer should wait for sufficient amount of rainfall (75-100 mm in a week) for sowing of Karif Rice.**

**Kharif rice:**

Sowing of rice in DSR method should be completed as soon as possible as rainfall is forecasted in between upcoming 5 days. Sowing should be started soon after receive of good amount of early monsoon showers (75-100 mm in a week) where direct seeding is practiced. Incorporate FYM or compost @ 2000 Kg/acre. Apply 27 kg DAP and 20 kg MoP per acre during last ploughing. Before sowing seed should be treated with Trichoderma viride @ 10 g or Carbendazim 50% WP @ 1.5 g/kg seed.

**Transplanting rice:**

After receive of sufficient rainfall, farmers are advised for sowing of dhanicha seeds @ 12 kg/acre prior to the transplanting of paddy so that the dhanicha crop will be 3-4 weeks old before 10-15 days of transplanting of rice. Apply all phosphatic fertilizers of Kharif rice to this green manuring crop in advance. Farmers who are sowing paddy nurseries are advised to treat the seeds with carbendizem @3 g/kg seed followed by shade drying for 24 hours before sowing in case of dry nursery and in case of wet nursery seeds should be treated with carbendizem @ 1 g/l of water by soaking for 24 hrs and another 24 hrs for sprouting and the seeds should be sown in wet nurseries.

**Maize:** It is time for sowing of maize for green cob purpose under irrigated condition Grow high yielding varieties like Navjot, Shakti, Decan-107, Decan-109, Use 6 kg seeds/acre. Apply one third of N and all P & K of the recommended dose of 50:24:24 kg. NPK/acre. Sow the seeds in lines with spacing of 45-60X20 cm at 5 cm soil depth.

**VEGETABLES AND FRUITS:**

- Fruit and shoot borer in Brinjal-** Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@300g/acre at 15 days interval.
- Due to Past rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- After cessation of rain, **downy mildew disease** will appear in **cucurbits**. To control it, go for alternate spray of Metalaxyl+Mancozeb@2g/litre and Azoxystrobin+Tebuconazole @0.6g/litre of water at 10 days interval.
- Rust** disease is likely to appear in **Cowpea**. Spay Chlorothalonil@2g/litre of water.
- Wilting** may observe in **tomato and brinjal** field, to control it drenching with Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water.
- Current weather favours spread of **YMV** in **Ladies finger**. In case of appearance of diseases, rogue out affected plants and spray Thiomethoxam 25WG @ 60g/acre.

**Fruit:**

**Banana:** For **leaf spot/sigatoka** as a precautionary measure spray carbendizim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalonil@ 2.0 g/ litre of water. Drenching soil with Metiram 75% WG @1.5g/4 lit or carbendazim 12%+ mancozeb 63% @ 2g/lit to prevent **Panama wilt**.

**Mango:** Harvest the mature mango fruits during morning or evening hours. Keep the harvested fruits in shade to prevent from spongy tissue disease and heat.

**Papaya:** After rain due to water logging there may be increase in the infestation of **phytopthera crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

**Animal Husbandry:** Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination. Keep the animal shed clean and dry.





**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**



(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-503  
Date- 14/06/2024

**AGROMET ADVISORY BULLETIN FOR NISCHINTAKOILI**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 0mm and district received average 1 mm rainfall during the last week. **The mean daily maximum and minimum temperature was 38.47°C (+2.5 from normal) and 26.5°C (+1.5 from normal), respectively.** The mean relative humidity was 72-83%.

**Forecasted period: Today(8.30IST) to 19<sup>th</sup> June (8.30IST)2024**

**Weather advisory: Very light to light rain/Thundershower is forecasted for upcoming 5 days in the block.** The day temperature will be 34-38°C and night temperature will be 27-31°C for upcoming 5 days. Maximum and minimum relative humidity will be 61-72% and 23-38% respectively. **There will be clear to partly cloudy sky for next 5 days.**

Block	Nischintakoili (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	14/06/2024	15/06/2024	16/06/2024	17/06/2024	18/06/2024
Rainfall(mm)	1	2	2	7	8
T-Max(°C)	38	38	37	34	34
T-Min(°C)	29	31	29	27	28
Cloud Cover(octa)	6	5	5	7	8
RH Max (%)	70	61	66	72	67
RH Min (%)	23	23	25	32	38
Wind Speed (Kmph)	23	19	22	23	26
Wind Direction (deg)	205	205	207	207	212

**Agromet Advisory**

**General advisory: Farmer should wait for sufficient amount of rainfall (75-100 mm in a week) for sowing of Karif Rice.**

**Kharif rice:**

Sowing of rice in DSR method should be completed as soon as possible as rainfall is forecasted in between upcoming 5 days. Sowing should be started soon after receive of good amount of early monsoon showers (75-100 mm in a week) where direct seeding is practiced. Incorporate FYM or compost @ 2000 Kg/acre. Apply 27 kg DAP and 20 kg MoP per acre during last ploughing. Before sowing seed should be treated with Trichoderma viride @ 10 g or Carbendazim 50% WP @ 1.5 g/kg seed.

**Transplanting rice:**

After receive of sufficient rainfall, farmers are advised for sowing of dhanicha seeds @ 12 kg/acre prior to the transplanting of paddy so that the dhanicha crop will be 3-4 weeks old before 10-15 days of transplanting of rice. Apply all phosphatic fertilizers of Kharif rice to this green manuring crop in advance. Farmers who are sowing paddy nurseries are advised to treat the seeds with carbendizem @3 g/kg seed followed by shade drying for 24 hours before sowing in case of dry nursery and in case of wet nursery seeds should be treated with carbendizem @ 1 g/l of water by soaking for 24 hrs and another 24 hrs for sprouting and the seeds should be sown in wet nurseries.

**Maize:** It is time for sowing of maize for green cob purpose under irrigated condition Grow high yielding varieties like Navjot, Shakti, Decan-107, Decan-109, Use 6 kg seeds/acre. Apply one third of N and all P & K of the recommended dose of 50:24:24 kg. NPK/acre. Sow the seeds in lines with spacing of 45-60X20 cm at 5 cm soil depth.

**VEGETABLES AND FRUITS:**

- **Fruit and shoot borer** in **Brinjal**- Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@300g/acre at 15 days interval.
- Due to Past rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- After cessation of rain, **downy mildew disease** will appear in **cucurbits**. To control it, go for alternate spray of Metalaxyl+Mancozeb@2g/litre and Azoxystrobin+Tebuconazole @0.6g/litre of water at 10 days interval.
- **Rust** disease is likely to appear in **Cowpea**. Spray Chlorothalonil@2g/litre of water.
- **Wilting** may observe in **tomato and brinjal** field, to control it drenching with Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water.
- Current weather favours spread of **YMV** in **Ladies finger**. In case of appearance of diseases, rogue out affected plants and spray Thiomethoxam 25WG @ 60g/acre.

**Fruit:**

**Banana:** For **leaf spot/sigatoka** as a precautionary measure spray carbendizim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalonil@ 2.0 g/ litre of water. Drenching soil with Metiram 75% WG @1.5g/4 lit or carbendazim 12%+ mancozeb 63% @ 2g/lit to prevent **Panama wilt**.

**Mango:** Harvest the mature mango fruits during morning or evening hours. Keep the harvested fruits in shade to prevent from spongy tissue disease and heat.

**Papaya:** After rain due to water logging there may be increase in the infestation of **phytophthora crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

**Animal Husbandry:** Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination. Keep the animal shed clean and dry.





**GRAMIN KRISHI MAUSAM SEWA  
INDIA METEOROLOGICAL DEPARTMENT  
DISTRICT AGROMET UNIT  
KRISHI VIGYAN KENDRA CUTTACK**



**(A Unit of ICAR-National Rice Research Institute)  
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com**

**Bulletin No.-503  
Date- 14/06/2024**

**AGROMET ADVISORY BULLETIN FOR SALEPUR**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 0.9 mm and district received average 1 mm rainfall during the last week. **The mean daily maximum and minimum temperature was 38.47°C (+2.5 from normal) and 26.5°C (+1.5 from normal), respectively.** The mean relative humidity was 72-83%.

**Forecasted period: Today(8.30IST) to 19<sup>th</sup> June (8.30IST)2024**

**Weather advisory: Very light to light rain/Thundershower is forecasted for upcoming 5days in the block.** The day temperature will be 33-38°C and night temperature will be 27-31°C for upcoming 5 days. Maximum and minimum relative humidity will be 60-72% and 23-37% respectively. **There will be clear to partly cloudy sky for next 5days.**

Block	Salepur (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	14/06/2024	15/06/2024	16/06/2024	17/06/2024	18/06/2024
Rainfall(mm)	1	2	2	8	7
T-Max(°C)	38	37	37	33	34
T-Min(°C)	29	31	29	27	27
Cloud Cover(octa)	6	5	5	7	8
RH Max (%)	71	60	66	72	67
RH Min (%)	23	23	25	31	37
Wind Speed (Kmph)	23	18	22	23	25
Wind Direction (deg)	205	210	212	212	212

**Agromet Advisory**

**General advisory: Farmer should wait for sufficient amount of rainfall (75-100 mm in a week) for sowing of Kharif Rice.**

**Kharif rice:**

Sowing of rice in DSR method should be completed as soon as possible as rainfall is forecasted in between upcoming 5 days. Sowing should be started soon after receive of good amount of early monsoon showers (75-100 mm in a week) where direct seeding is practiced. Incorporate FYM or compost @ 2000 Kg/acre. Apply 27 kg DAP and 20 kg MoP per acre during last ploughing. Before sowing seed should be treated with Trichoderma viride @ 10 g or Carbendazim 50% WP @ 1.5 g/kg seed.

**Transplanting rice:**

After receive of sufficient rainfall, farmers are advised for sowing of dhanicha seeds @ 12 kg/acre prior to the transplanting of paddy so that the dhanicha crop will be 3-4 weeks old before 10-15 days of transplanting of rice. Apply all phosphatic fertilizers of Kharif rice to this green manuring crop in advance. Farmers who are sowing paddy nurseries are advised to treat the seeds with carbendizem @3 g/kg seed followed by shade drying for 24 hours before sowing in case of dry nursery and in case of wet nursery seeds should be treated with carbendizem @ 1 g/l of water by soaking for 24 hrs and another 24 hrs for sprouting and the seeds should be sown in wet nurseries.

**Maize:** It is time for sowing of maize for green cob purpose under irrigated condition Grow high yielding varieties like Navjot, Shakti, Decan-107, Decan-109, Use 6 kg seeds/acre. Apply one third of N and all P & K of the recommended dose of 50:24:24 kg. NPK/acre. Sow the seeds in lines with spacing of 45-60X20 cm at 5 cm soil depth.

**VEGETABLES AND FRUITS:**

- **Fruit and shoot borer** in **Brinjal**- Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@300g/acre at 15 days interval.
- Due to Past rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- After cessation of rain, **downy mildew disease** will appear in **cucurbits**. To control it, go for alternate spray of Metalaxyl+Mancozeb@2g/litre and Azoxystrobin+Tebuconazole @0.6g/litre of water at 10 days interval.
- **Rust** disease is likely to appear in **Cowpea**. Spray Chlorothalonil@2g/litre of water.
- **Wilting** may observe in **tomato and brinjal** field, to control it drenching with Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water.
- Current weather favours spread of **YMV** in **Ladies finger**. In case of appearance of diseases, rogue out affected plants and spray Thiometoxam 25WG @ 60g/acre.

**Fruit:**

**Banana:** For **leaf spot/sigatoka** as a precautionary measure spray carbendizim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalonil@ 2.0 g/ litre of water. Drenching soil with Metiram 75% WG @1.5g/4 lit or carbendazim 12%+ mancozeb 63% @ 2g/lit to prevent **Panama wilt**.

**Mango:** Harvest the mature mango fruits during morning or evening hours. Keep the harvested fruits in shade to prevent from spongy tissue disease and heat.

**Papaya:** After rain due to water logging there may be increase in the infestation of **phytophthora crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

**Animal Husbandry:** Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination. Keep the animal shed clean and dry.







**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**



(A Unit of ICAR-National Rice Research Institute)  
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-503  
 Date- 14/06/2024

**AGROMET ADVISORY BULLETIN FOR TANGI-CHOUDWAR**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 0mm and district received average 1 mm rainfall during the last week. The mean daily maximum and minimum temperature was 38.47°C (+2.5 from normal) and 26.5°C (+1.5 from normal), respectively. The mean relative humidity was 72-83%.

**Forecasted period: Today(8.30IST) to 19<sup>th</sup> June (8.30IST)2024**

**Weather advisory:** *Very light to light rain/Thundershower is forecasted for upcoming 5days in the block.* The day temperature will be 37-42°C and night temperature will be 31-32°C for upcoming 5 days. Maximum and minimum relative humidity will be 71-78% and 34-54% respectively. There will be clear to partly cloudy sky for next 5days.

Block	Tangi-Choudwar (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	14/06/2024	15/06/2024	16/06/2024	17/06/2024	18/06/2024
Rainfall(mm)	1	2	2	7	7
T-Max(°C)	41	42	40	37	37
T-Min(°C)	32	32	32	31	31
Cloud Cover(octa)	6	5	5	7	8
RH Max (%)	75	71	76	78	72
RH Min (%)	34	36	42	54	53
Wind Speed (Kmph)	28	23	29	32	31
Wind Direction (deg)	207	204	203	203	212

**Agromet Advisory**

**General advisory:** Farmer should wait for sufficient amount of rainfall (75-100 mm in a week) for sowing of Kharif Rice.

**Kharif rice:**

Sowing of rice in DSR method should be completed as soon as possible as rainfall is forecasted in between upcoming 5 days. Sowing should be started soon after receive of good amount of early monsoon showers (75-100 mm in a week) where direct seeding is practiced. Incorporate FYM or compost @ 2000 Kg/acre. Apply 27 kg DAP and 20 kg MoP per acre during last ploughing. Before sowing seed should be treated with Trichoderma viride @ 10 g or Carbendazim 50% WP @ 1.5 g/kg seed.

**Transplanting rice:**

After receive of sufficient rainfall, farmers are advised for sowing of dhanicha seeds @ 12 kg/acre prior to the transplanting of paddy so that the dhanicha crop will be 3-4 weeks old before 10-15 days of transplanting of rice. Apply all phosphatic fertilizers of Kharif rice to this green manuring crop in advance. Farmers who are sowing paddy nurseries are advised to treat the seeds with carbendizem @3 g/kg seed followed by shade drying for 24 hours before sowing in case of dry nursery and in case of wet nursery seeds should be treated with carbendizem @ 1 g/l of water by soaking for 24 hrs and another 24 hrs for sprouting and the seeds should be sown in wet nurseries.

**Maize:** It is time for sowing of maize for green cob purpose under irrigated condition Grow high yielding varieties like Navjot, Shakti, Decan-107, Decan-109, Use 6 kg seeds/acre. Apply one third of N and all P & K of the recommended dose of 50:24:24 kg. NPK/acre. Sow the seeds in lines with spacing of 45-60X20 cm at 5 cm soil depth.

**VEGETABLES AND FRUITS:**

- Fruit and shoot borer in Brinjal-** Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@300g/acre at 15 days interval.
- Due to Past rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- After cessation of rain, **downy mildew disease** will appear in **cucurbits**. To control it, go for alternate spray of Metalaxyl+Mancozeb@2g/litre and Azoxytrobilin+Tebuconazole @0.6g/litre of water at 10 days interval.
- Rust** disease is likely to appear in **Cowpea**. Spray Chlorothalonil@2g/litre of water.
- Wilting** may observe in **tomato and brinjal** field, to control it drenching with Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water.
- Current weather favours spread of **YMV** in **Ladies finger**. In case of appearance of diseases, rogue out affected plants and spray Thiomethoxam 25WG @ 60g/acre.

**Fruit:**

**Banana:** For **leaf spot/sigatoka** as a precautionary measure spray carbendizim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalonil@ 2.0 g/ litre of water. Drenching soil with Metiram 75% WG @1.5g/4 lit or carbendazim 12%+ mancozeb 63% @ 2g/lit to prevent **Panama wilt**.

**Mango:** Harvest the mature mango fruits during morning or evening hours. Keep the harvested fruits in shade to prevent from spongy tissue disease and heat.

**Papaya:** After rain due to water logging there may be increase in the infestation of **phytophthora crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

**Animal Husbandry:** Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination. Keep the animal shed clean and dry.

Dr Ranjan Kumar Mohanta  
 (Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena  
 SMS (Agrometeorology)  
 Contact No-9861445080/7008453427



**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**



(A Unit of ICAR-National Rice Research Institute)  
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-503  
 Date- 14/06/2024

**AGROMET ADVISORY BULLETIN FOR TIGIRIA**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 0mm and district received average 1 mm rainfall during the last week. The mean daily maximum and minimum temperature was 38.47°C (+2.5 from normal) and 26.5°C (+1.5 from normal), respectively. The mean relative humidity was 72-83%.

**Forecasted period: Today(8.30IST) to 19<sup>th</sup> June (8.30IST)2024**

**Weather advisory: Very light to light rain/Thundershower is forecasted for upcoming 5days in the block.** The day temperature will be 33-37°C and night temperature will be 25-30°C for upcoming 5 days. Maximum and minimum relative humidity will be 52-70% and 21-30% respectively. **There will be clear to partly cloudy sky for next 5days.**

Block	Tigiria (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	14/06/2024	15/06/2024	16/06/2024	17/06/2024	18/06/2024
Rainfall(mm)	1	2	2	5	7
T-Max(°C)	37	34	36	33	36
T-Min(°C)	29	30	28	26	25
Cloud Cover(octa)	6	7	6	7	8
RH Max (%)	62	52	60	70	69
RH Min (%)	21	28	26	30	30
Wind Speed (Kmph)	16	13	13	14	16
Wind Direction (deg)	200	214	203	205	208

**Agro met Advisory**

**General advisory: Farmer should wait for sufficient amount of rainfall (75-100 mm in a week) for sowing of Karif Rice.**

**Kharif rice:**

Sowing of rice in DSR method should be completed as soon as possible as rainfall is forecasted in between upcoming 5 days. Sowing should be started soon after receive of good amount of early monsoon showers (75-100 mm in a week) where direct seeding is practiced. Incorporate FYM or compost @ 2000 Kg/acre. Apply 27 kg DAP and 20 kg MoP per acre during last ploughing. Before sowing seed should be treated with Trichoderma viride @ 10 g or Carbendazim 50% WP @ 1.5 g/kg seed.

**Transplanting rice:**

After receive of sufficient rainfall, farmers are advised for sowing of dhanicha seeds @ 12 kg/acre prior to the transplanting of paddy so that the dhanicha crop will be 3-4 weeks old before 10-15 days of transplanting of rice. Apply all phosphatic fertilizers of Kharif rice to this green manuring crop in advance. Farmers who are sowing paddy nurseries are advised to treat the seeds with carbendazim @ 3 g/kg seed followed by shade drying for 24 hours before sowing in case of dry nursery and in case of wet nursery seeds should be treated with carbendazim @ 1 g/l of water by soaking for 24 hrs and another 24 hrs for sprouting and the seeds should be sown in wet nurseries.

**Maize:** It is time for sowing of maize for green cob purpose under irrigated condition Grow high yielding varieties like Navjot, Shakti, Decan-107, Decan-109, Use 6 kg seeds/acre. Apply one third of N and all P & K of the recommended dose of 50:24:24 kg. NPK/acre. Sow the seeds in lines with spacing of 45-60X20 cm at 5 cm soil depth.

**VEGETABLES AND FRUITS:**

- **Fruit and shoot borer** in **Brinjal**- Spray Neem based pesticide 1500ppm@600ml/acre at initial stages. Later on go for alternate spray of Spinosad 45%SC@70ml/acre and Thiodicarb@300g/acre at 15 days interval.
- Due to Past rain, Chances of **fruit rot** infestation in **chilli and brinjal**, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- After cessation of rain, **downy mildew disease** will appear in **cucurbits**. To control it, go for alternate spray of Metalaxyl+Mancozeb@2g/litre and Azoxystrobin+Tebuconazole @0.6g/litre of water at 10 days interval.
- **Rust** disease is likely to appear in **Cowpea**. Spray Chlorothalonil@2g/litre of water.
- **Wilting** may observe in **tomato and brinjal** field, to control it drenching with Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water.
- Current weather favours spread of **YMV** in **Ladies finger**. In case of appearance of diseases, rogue out affected plants and spray Thiometoxam 25WG @ 60g/acre.

**Fruit:**

**Banana:** For **leaf spot/sigatoka** as a precautionary measure spray carbendazim + mancozeb @ 2.0 g / litre of water followed by (15 days after) chlorothalonil @ 2.0 g/ litre of water. Drenching soil with Metiram 75% WG @1.5g/4 lit or carbendazim 12%+ mancozeb 63% @ 2g/lit to prevent **Panama wilt**.

**Mango:** Harvest the mature mango fruits during morning or evening hours. Keep the harvested fruits in shade to prevent from spongy tissue disease and heat.

**Papaya:** After rain due to water logging there may be increase in the infestation of **phytophthora crown rot or foot rot** in papaya, so farmers should drain out excess water and drench the soil with fungicide like thiophanate methyl 70%wp 2-4g/ltr of water.

**Animal Husbandry:** Provide sufficient clean drinking water to animals. Contact departmental personnel for vaccination. Keep the animal shed clean and dry.











**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**  
 (A Unit of ICAR-National Rice Research Institute)  
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-503  
Date- 14/06/2024

				
MEGHDOOT APP	DAMINI APP	FACEBOOK (Damu Cuttack)	BLOCK SPECIFIC WHAT'S APP GROUPS	@DAMUCUTTACK

- Kindly install and use **“Meghdoot mobile App”** for Cuttack district weather forecast, warnings and weather based agromet advisory services.  
<https://play.google.com/store/apps/details?id=com.aas.meghdoot>
- Kindly install and use **“Damini mobile App”** for location specific for lightening forecast and warnings.  
<https://play.google.com/store/apps/details?id=com.lightening.live.damini>
- Follow the **block specific whats app groups** and join in your block wise groups for getting weather based agromet advisory services twice in a week (Tuesday and Friday) and other weather warning based farm management advisories.
- Follow the **Official Website links** to get the weather based agromet advisory service bulletins (**English & Odia**) twice in a week (Tuesday and Friday).

1	ICAR-NRRI, Cuttack, Odisha	<a href="https://icar-nrri.in/agromate/">https://icar-nrri.in/agromate/</a>	
2	Agrimet division, IMD, Pune	<a href="https://imdagrmet.gov.in/imdproject/AGIndex.php">https://imdagrmet.gov.in/imdproject/AGIndex.php</a>	