




Cuttack District Block Wise Agro-met Advisory Service Bulletin

*Tuesday, 30th July, 2024
 (For the period 30th July to 03rd August,
 2024)*




*Issued by
 District Agro-met Unit,
 KrishiVigyan Kendra Cuttack*





STANDARD PRECIPITATION INDEX AND NDVI MAP



LEGEND (Medium range block level rainfall forecast)		
	No Rainfall	00 mm
	Very light rainfall	01-05 mm
	Light rainfall	06-10 mm
	Light to moderate rainfall	11-30 mm
	Moderate to heavy rainfall	31-65 mm
	Heavy to very heavy rainfall	66-115 mm
	Extremely heavy rainfall	>116 mm

LEGEND (Past week temperature)		
Color code	Temperature	Departure
	Below Normal	< 2°C from normal
	Normal	-2°C to +2°C from normal
	Above Normal	>2°C from normal





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-516
Date- 30/07/2024

Met Sub division	Extended range outlook for subsequent 1 week (6 th to 12 th days) (04-10 August 2024) after 5 days				
	Parameter	Actual	Normal	Departure (%)	Category
Odisha	Rainfall (mm)	15	12.2	23.36	Above Normal
	Parameter	Actual	Normal	Anomaly	Category
	Maximum temperature (°C)	31.0	30.8	0.20	Normal
	Minimum temperature (°C)	24.6	24.2	0.45	Normal

***Agromet Advisory based upon subsequent week extended range forecast-As there will be above normal rainfall in the subsequent week (04-10 August,2024) after 5 days in Odisha condition,so farmers are advised to go for transplanting and beushaning of early sown kharif paddy in the subsequent week and are advised to make proper drainage facilities in nursery area of paddy and vegetables to drain out excess water from the field. Farmers should pause sowing activity in the subsequent week. They should monitor paddy field for seedling blight, Blast and Sheath blight diseases.**



DrRanjan Kumar Mohanta
(Nodal officer)













Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



Weather (Past and Forecasted) based Disease and pest in Major Crops and vegetables

Sl No	Crop	Stage and condition of the crop	Anticipated Disease and pest	
1	Rice	Nursery stage		Blast
				Seedling blight
				Sheath blight
2	Sugarcane	Vegetative to Tillering		Early shot borer
				Top borer
				Smut
3	Maize	5 week stage		Inter node borer
				Stem fly
				Leaf blight
3	Jute	30-50days		Stem rot
















**GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK**



(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-516
Date- 30/07/2024

				Indigo caterpillar
4	Brinjal, tomatoto, chilli	seedling		Seedling rot
				Stem rot
5	Tomato and brinjal	Fruting		Bacterial wilt
6	chilli	fruting		thrips
				Leaf curl
7	okra	Fruting		Fruit and shoot borer
				White fly
				Jassid
8	Banana			Sigatoka
9	Papaya			Yellow mosaic virus





**GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK**



(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

**Bulletin No.-516
Date- 30/07/2024**

AGROMET ADVISORY BULLETIN FOR ATHAGAD

District Past week average weather condition and Block past week average Rainfall condition

The block received average 14 mm and district received average 13 mm rainfall during the last week. The mean daily maximum and minimum temperature was 29.85°C (-2.37 from normal) and 24.85°C (+0.64 from normal), respectively. The mean relative humidity was 92-94%.

Forecasted period: Today(8.30IST) to 04th August(8.30IST)2024

Weather advisory: Moderate with isolated heavy rainfall is forecasted for tomorrow and light to moderate rainfall is forecasted for other 4 days in the block. The day temperature will be 31-33°C and night temperature will be 25-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 82-87% and 53-65% respectively. **There will be partly cloudy to overcast sky for next 5 days.**

Block	Athagad (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	30/07/2024	31/07/2024	01/08/2024	02/08/2024	03/08/2024
Rainfall (mm)	25	31	12	8	6
T-Max (°C)	33	32	31	31	31
T-Min (°C)	26	26	25	25	25
Cloud Cover (octa)	8	8	8	8	7
RH Max (%)	82	87	84	83	85
RH Min (%)	53	58	65	59	57
Wind Speed (kmph)	12	11	12	13	8
Wind Direction (deg)	225	248	238	212	210

Agromet Advisory

General advisory: Drain out excess water from rice nurseries, orchards, maize and jute fields. Postpone irrigation, intercultural operations, application of plant protection measures and fertilizers to the standing crops in upcoming 5 days. Keep the harvested produce at safer places. Cover the vegetable seedling with polythene to protect them from heavy shower.

Kharifrice:(DIRECT SEEDED RICE):

Direct seeded rice are at three to four week stage. As light to moderate amount of rainfall is forecasted in the upcoming days, farmers should go for beushanning. Apply 35 kg urea/ac at beushanning followed by weeding and gap filling. Drain out the excess water at the time of fertilizer application and pond it after one day. If basal fertilizer not applied, apply 18 kg urea, 100kg ssp and 26 kg MOP while beushanning followed by top dressing of 35 kg urea after 7-10 days.

TRANSPLANTED RICE:

Considering the anticipated rainfall, paddy transplanting may be completed before 30th July. Apply 400-gram Chlorantraniliprole 0.4 % G or 800-gram Cartap Hydrochloride 4% GR in the 10 decimal nursery area 7 days before transplanting to manage gall midge, stem borer, caseworm, leaf folder and root knot nematode up to 3 weeks after transplanting.

After rain in dry weather go for foliar spray of Hexaconazole with Plantomycin @ 1.0 g/l to prevent the spread of BLB and sheath blight. If leaf blast is observed in nursery, spray Tebuconazole 50% + Trifloxystrobin 25% WG @ 0.4 g or Isoprothiolane 40EC @ 1.5 ml per litre of water. Repeat the spray at 7-10 days interval.

Maize:

Stem fly in Maize infects the plants from sowing to 5 weeks after sowing, temperature between 25-30 deg. celsius and humidity above 60% favour pest incidence. In vegetative stage to control stem fly spray Methyl demeton @ 1 ml/lt or Dimethoate @ 2ml/lt or apply Carbofuran 3G granuals @ 14 kg/acre. High humid conditions, temperature between 18-27 Deg. Celsius favours the occurrence of **leaf blight** in Maize. To control spray Azoxystrobin + Mancozeb 300 g /ac.

VEGETABLES:

- June-july is the best time for sowing of tomato in raised bed for nursery seedlings. Do seed treatment with Trichoderma viride @ 4g/kg seeds. The seed treatment must be done 24 hours before sowing of seeds.
- Due to forecast of rain, Chances of **fruit rot** infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever **sucking pests** incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. **To prevent of soil borne diseases like seedling rot, stem rot and wilt diseases apply 3g of Copper oxychloride per litre of water around the main stem.**
- Incidence of **thrips** may observe in chilli, if observe spray Spinosad 45% SC @ 0.4 ml/litre of water or Imidacloprid 17.8% SL @ 0.3 ml/litre of water in clean and dry weather. The present weather conditions are favourable for Upward and downward leafcurl in chilli. To control spray phosalon 3 ml. or diphenhiuron @ 1g./L of water.
- To control **fruit and shoot borer** in okra, Spray Bacillus thuringiensis 5% W.P @ 1.5 gram/lit of water. Present weather condition favours incidence of white fly and jassid. To control this apply Thiomethoxam @ 0.4g/ litre of water.

Fruits:

Spray mancozeb 2g/litre, to control sigatoka leaf spot in **banana**. To control **papaya yellow mosaic virus**, spray Imidachloprid @ 0.5 ml / litre.





**GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK**



(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-516
Date- 30/07/2024

AGROMET ADVISORY BULLETIN FOR BADAMBA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 05 mm and district received average 13 mm rainfall during the last week. The mean daily maximum and minimum temperature was 29.85°C (-2.37 from normal) and 24.85°C (+0.64 from normal), respectively. The mean relative humidity was 92-94%.

Forecasted period: Today(8.30IST) to 04th August(8.30IST)2024

Weather advisory: Moderate with isolated heavy rainfall is forecasted for tomorrow and light to moderate rainfall is forecasted for other 4 in the block. The day temperature will be 31-32°C and night temperature will be 24-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 83-88% and 55-64% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Badamba (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	30/07/2024	31/07/2024	01/08/2024	02/08/2024	03/08/2024
Rainfall(mm)	28	32	12	6	6
T-Max(°C)	32	32	31	31	31
T-Min(°C)	26	25	24	24	24
Cloud Cover(octa)	8	8	8	8	7
RH Max (%)	83	88	83	83	86
RH Min (%)	55	58	64	59	58
Wind Speed (Kmph)	11	10	11	13	7
Wind Direction (deg)	243	244	238	212	238

Agromet Advisory

General advisory: Drain out excess water from rice nurseries, orchards, maize and jute fields. Postpone irrigation, intercultural operations, application of plant protection measures and fertilizers to the standing crops in upcoming 5 days. Keep the harvested produce at safer places. Cover the vegetable seedling with polythene to protect them from heavy shower.

Kharifrice:(DIRECT SEEDED RICE):

Direct seeded rice are at three to four week stage. As light to moderate amount of rainfall is forecasted in the upcoming days, farmers should go for beushanning. Apply 35 kg urea/ac at beushanning followed by weeding and gap filling. Drain out the excess water at the time of fertilizer application and pond it after one day. If basal fertilizer not applied, apply 18 kg urea, 100kg ssp and 26 kg MOP while beushanning followed by top dressing of 35 kg urea after 7-10 days.

TRANSPLANTED RICE:

Considering the anticipated rainfall, paddy transplanting may be completed before 30th July. Apply 400-gram Chlorantranilprole 0.4 % G or 800-gram Cartap Hydrochloride 4% GR in the 10 decimal nursery area 7 days before transplanting to manage gall midge, stem borer, caseworm, leaf folder and root knot nematode up to 3 weeks after transplanting.

After rain in dry weather go for foliar spray of Hexaconazole with Plantomycin @ 1.0 g/l to prevent the spread of BLB and sheath blight. If leaf blast is observed in nursery, spray Tebuconazole 50% + Trifloxystrobin 25% WG @ 0.4 g or Isoprothiolane 40EC @ 1.5 ml per litre of water. Repeat the spray at 7-10 days interval.

Maize:

Stem fly in Maize infects the plants from sowing to 5 weeks after sowing, temperature between 25-30 deg. celsius and humidity above 60% favour pest incidence. In vegetative stage to control stem fly spray Methyl demeton @ 1 ml/lit or Dimethoate @ 2ml/lit or apply Carbofuran 3G granuals @ 14 kg/acre. High humid conditions, temperature between 18-27 Deg. Celsius favours the occurrence of leaf blight in Maize. To control spray Azoxystrobin + Mancozeb 300 g /ac.

VEGETABLES:

- June-july is the best time for sowing of tomato in raised bed for nursery seedlings. Do seed treatment with Trichoderma viride @ 4g/kg seeds. The seed treatment must be done 24 hours before sowing of seeds.
- Due to forecast of rain, Chances of fruit rot infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever sucking pests incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. To prevent of soil borne diseases like seedling rot, stem rot and wilt diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Incidence of thrips may observe in chilli, if observe spray Spinosad 45% SC @ 0.4 ml/litre of water or Imidacloprid 17.8% SL @ 0.3 ml/litre of water in clean and dry weather. The present weather conditions are favourable for Upward and downward leafcurl in chilli. To control spray phosalon 3 ml. or diphenhiuron @ 1g./L of water.
- To control fruit and shoot borer in okra, Spray Bacillus thuringiensis 5% W.P @ 1.5 gram/lit of water. Present weather condition favours incidence of white fly and jassid. To control this apply Thiomethoxam @ 0.4g/ litre of water.

Fruits:

Spray mancozeb 2g/litre, to control sigatoka leaf spot in banana. To control papaya yellow mosaic virus, spray Imidachloprid @ 0.5 ml / litre.

DrRanjan Kumar Mohanta
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



**GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK**



**(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com**

**Bulletin No.-516
Date- 30/07/2024**

AGROMET ADVISORY BULLETIN FOR BANKI

District Past week average weather condition and Block past week average Rainfall condition

The district received average 13 mm rainfall during the last week. The mean daily maximum and minimum temperature was 29.85°C (-2.37 from normal) and 24.85°C (+0.64 from normal), respectively. The mean relative humidity was 92-94%.

Forecasted period: Today(8.30IST) to 04th August(8.30IST)2024

Weather advisory: Moderate with isolated heavy rainfall is forecasted for tomorrow and light to moderate rainfall is forecasted for other 4 in the block. The day temperature will be 31-33°C and night temperature will be 24-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 81-86% and 54-62% respectively. **There will be partly cloudy to overcast sky for next 5 days.**

Block	Banki (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	30/07/2024	31/07/2024	01/08/2024	02/08/2024	03/08/2024
Rainfall(mm)	28	31	11	6	6
T-Max(°C)	33	32	31	31	31
T-Min(°C)	26	26	24	25	24
Cloud Cover(octa)	8	8	8	8	7
RH Max (%)	82	86	82	81	84
RH Min (%)	54	56	62	57	57
Wind Speed (Kmph)	12	11	13	14	8
Wind Direction (deg)	243	244	243	212	216

Agromet Advisory

General advisory: Drain out excess water from rice nurseries, orchards, maize and jute fields. Postpone irrigation, intercultural operations, application of plant protection measures and fertilizers to the standing crops in upcoming 5 days. Keep the harvested produce at safer places. Cover the vegetable seedling with polythene to protect them from heavy shower.

Kharifrice:(DIRECT SEEDED RICE):

Direct seeded rice are at three to four week stage. As light to moderate amount of rainfall is forecasted in the upcoming days, farmers should go for beushanning. Apply 35 kg urea/ac at beushanning followed by weeding and gap filling. Drain out the excess water at the time of fertilizer application and pond it after one day. If basal fertilizer not applied, apply 18 kg urea, 100kg ssp and 26 kg MOP while beushanning followed by top dressing of 35 kg urea after 7-10 days.

TRANSPLANTED RICE:

Considering the anticipated rainfall, paddy transplanting may be completed before 30th July. Apply 400-gram Chlorantranilprole 0.4 % G or 800-gram Cartap Hydrochloride 4% GR in the 10 decimal nursery area 7 days before transplanting to manage gall midge, stem borer, caseworm, leaf folder and root knot nematode up to 3 weeks after transplanting.

After rain in dry weather go for foliar spray of Hexaconazole with Plantomycin@1.0 g/l to prevent the spread of BLB and sheath blight. If leaf blast is observed in nursery, spray Tebuconazole 50% + Trifloxystrobin 25% WG@ 0.4 g or Isoprothiolane 40EC @ 1.5 ml per litre of water. Repeat the spray at 7-10 days interval.

Maize:

Stem fly in Maize infects the plants from sowing to 5 weeks after sowing, temperature between 25-30 deg. celsius and humidity above 60% favour pest incidence. In vegetative stage to control stem fly spray Methyl demeton @1 ml/lit or Dimethoate @ 2ml/lit or apply Carbofuran 3G granuals @ 14 kg/acre. High humid conditions, temperature between 18-27 Deg. Celsius favours the occurrence of **leaf blight** in Maize. To control spray Azoxystrobin + Mancozeb 300 g /ac.

VEGETABLES:

- June-july is the best time for sowing of tomato in raised bed for nursery seedlings. Do seed treatment with Trichoderma viride @ 4g/kg seeds. The seed treatment must be done 24 hours before sowing of seeds.
- Due to forecast of rain, Chances of **fruit rot** infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever **sucking pests** incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. **To prevent of soil borne diseases like seedling rot, stem rot and wilt diseases apply 3g of Copper oxychloride per litre of water around the main stem.**
- Incidence of **thrips** may observe in chilli, if observe spray Spinosad 45% SC @ 0.4 ml/litre of water or Imidacloprid 17.8% SL @ 0.3 ml/litre of water in clean and dry weather. The present weather conditions are favourable for Upward and downward leaf curl in chilli. To control spray phosalon 3 ml. or diphenhiuron @ 1g./L of water.
- To control **fruit and shoot borer** in okra, Spray Bacillus thuringiensis 5% W.P @1.5 gram/lit of water. Present weather condition favours incidence of white fly and jassid. To control this apply Thiomethoxam @0.4g/ litre of water.

Fruits:

Spray mancozeb 2g/litre, to control sigatoka leaf spot in **banana**. To control **papaya yellow mosaic virus**, spray Imidachloprid @ 0.5 ml / litre.

**DrRanjan Kumar Mohanta
(Nodal officer)**



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



**Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427**



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK



(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-516

Date- 30/07/2024

AGROMET ADVISORY BULLETIN FOR BANKI-DAMPARA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 29 mm and district received average 13 mm rainfall during the last week. The mean daily maximum and minimum temperature was 29.85°C (-2.37 from normal) and 24.85°C (+0.64 from normal), respectively. The mean relative humidity was 92-94%.

Forecasted period: Today(8.30IST) to 04th August(8.30IST)2024

Weather Advisory: Moderate with isolated heavy rainfall is forecasted for tomorrow and light to moderate rainfall is forecasted for other 4 in the block. The day temperature will be 31-33°C and night temperature will be 25-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 81-86% and 54-63% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Banki-Dampara (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	30/07/2024	31/07/2024	01/08/2024	02/08/2024	03/08/2024
Rainfall(mm)	30	32	11	7	6
T-Max(°C)	33	32	31	31	31
T-Min(°C)	26	26	25	25	25
Cloud Cover(octa)	8	8	8	8	7
RH Max (%)	82	86	82	81	84
RH Min (%)	54	56	63	57	57
Wind Speed (Kmph)	12	11	13	13	9
Wind Direction (deg)	243	244	243	212	216

Agromet Advisory

General advisory: Drain out excess water from rice nurseries, orchards, maize and jute fields. Postpone irrigation, intercultural operations, application of plant protection measures and fertilizers to the standing crops in upcoming 5 days. Keep the harvested produce at safer places. Cover the vegetable seedling with polythene to protect them from heavy shower.

Kharifrice:(DIRECT SEEDED RICE):

Direct seeded rice are at three to four week stage. As light to moderate amount of rainfall is forecasted in the upcoming days, farmers should go for beushanning. Apply 35 kg urea/ac at beushanning followed by weeding and gap filling. Drain out the excess water at the time of fertilizer application and pond it after one day. If basal fertilizer not applied, apply 18 kg urea, 100kg ssp and 26 kg MOP while beushanning followed by top dressing of 35 kg urea after 7-10 days.

TRANSPLANTED RICE:

Considering the anticipated rainfall, paddy transplanting may be completed before 30th July. Apply 400-gram Chlorantraniliprole 0.4 % G or 800-gram Cartap Hydrochloride 4% GR in the 10 decimal nursery area 7 days before transplanting to manage gall midge, stem borer, caseworm, leaf folder and root knot nematode up to 3 weeks after transplanting.

After rain in dry weather go for foliar spray of Hexaconazole with Plantomycin @ 1.0 g/l to prevent the spread of BLB and sheath blight. If leaf blast is observed in nursery, spray Tebuconazole 50% + Trifloxystrobin 25% WG @ 0.4 g or Isoprothiolane 40EC @ 1.5 ml per litre of water. Repeat the spray at 7-10 days interval.

Maize:

Stem fly in Maize infects the plants from sowing to 5 weeks after sowing, temperature between 25-30 deg. celsius and humidity above 60% favour pest incidence. In vegetative stage to control stem fly spray Methyl demeton @ 1 ml/lt or Dimethoate @ 2ml/lt or apply Carbofuran 3G granuals @ 14 kg/acre. High humid conditions, temperature between 18-27 Deg. Celsius favours the occurrence of **leaf blight** in Maize. To control spray Azoxystrobin + Mancozeb 300 g /ac.

VEGETABLES:

- June-july is the best time for sowing of tomato in raised bed for nursery seedlings. Do seed treatment with Trichoderma viride @ 4g/kg seeds. The seed treatment must be done 24 hours before sowing of seeds.
- Due to forecast of rain, Chances of **fruit rot** infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever **sucking pests** incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. To prevent of soil borne diseases like seedling rot, stem rot and wilt diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Incidence of **thrips** may observe in chilli, if observe spray Spinosad 45% SC @ 0.4 ml/litre of water or Imidacloprid 17.8% SL @ 0.3 ml/litre of water in clean and dry weather. The present weather conditions are favourable for Upward and downward leafcurl in chilli. To control spray phosalon 3 ml. or diphenhiuron @ 1g./L of water.
- To control **fruit and shoot borer** in okra, Spray Bacillus thuringiensis 5% W.P @ 1.5 gram/lit of water. Present weather condition favours incidence of white fly and jassid. To control this apply Thiomethoxam @ 0.4g/ litre of water.

Fruits:

Spray mancozeb 2g/litre, to control sigatoka leaf spot in banana. To control papaya yellow mosaic virus, spray Imidachloprid @ 0.5 ml / litre.

DrRanjan Kumar Mohanta
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK



(A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-516
 Date- 30/07/2024

AGROMET ADVISORY BULLETIN FOR BARANGA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 17 mm and district received average 13 mm rainfall during the last week. The mean daily maximum and minimum temperature was 29.85°C (-2.37 from normal) and 24.85°C (+0.64 from normal), respectively. The mean relative humidity was 92-94%.

Forecasted period: Today(8.30IST) to 04th August(8.30IST)2024

Weather advisory: Moderate with isolated heavy rainfall is forecasted for tomorrow and light to moderate rainfall is forecasted for other 4 in the block. The day temperature will be 31-33°C and night temperature will be 26-27°C for upcoming 5 days. Maximum and minimum relative humidity will be 79-83% and 50-61% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Baranga (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	30/07/2024	31/07/2024	01/08/2024	02/08/2024	03/08/2024
Rainfall(mm)	18	33	11	7	6
T-Max(°C)	33	33	31	31	32
T-Min(°C)	27	27	26	26	26
Cloud Cover	8	8	8	8	7
RH Max (%)	79	83	80	80	81
RH Min (%)	50	55	61	56	54
Wind Speed (Kmph)	14	13	15	14	10
Wind Direction (deg)	216	244	243	212	214

Agromet Advisory

General advisory: Drain out excess water from rice nurseries, orchards, maize and jute fields. Postpone irrigation, intercultural operations, application of plant protection measures and fertilizers to the standing crops in upcoming 5 days. Keep the harvested produce at safer places. Cover the vegetable seedling with polythene to protect them from heavy shower.

Kharifrice:(DIRECT SEEDED RICE):

Direct seeded rice are at three to four week stage. As light to moderate amount of rainfall is forecasted in the upcoming days, farmers should go for beushanning. Apply 35 kg urea/ac at beushanning followed by weeding and gap filling. Drain out the excess water at the time of fertilizer application and pond it after one day. If basal fertilizer not applied, apply 18 kg urea, 100kg ssp and 26 kg MOP while beushanning followed by top dressing of 35 kg urea after 7-10 days.

TRANSPLANTED RICE:

Considering the anticipated rainfall, paddy transplanting may be completed before 30th July. Apply 400-gram Chlorantranilprole 0.4 % G or 800-gram Cartap Hydrochloride 4% GR in the 10 decimal nursery area 7 days before transplanting to manage gall midge, stem borer, caseworm, leaf folder and root knot nematode up to 3 weeks after transplanting.

After rain in dry weather go for foliar spray of Hexaconazole with Plantomycin @ 1.0 g/l to prevent the spread of BLB and sheath blight. If leaf blast is observed in nursery, spray Tebuconazole 50% + Trifloxystrobin 25% WG @ 0.4 g or Isoprothiolane 40EC @ 1.5 ml per litre of water. Repeat the spray at 7-10 days interval.

Maize:

Stem fly in Maize infects the plants from sowing to 5 weeks after sowing, temperature between 25-30 deg. celsius and humidity above 60% favour pest incidence. In vegetative stage to control stem fly spray Methyl demeton @ 1 ml/lit or Dimethoate @ 2ml/lit or apply Carbofuran 3G granuals @ 14 kg/acre. High humid conditions, temperature between 18-27 Deg. Celsius favours the occurrence of leaf blight in Maize. To control spray Azoxystrobin + Mancozeb 300 g /ac.

VEGETABLES:

- June-july is the best time for sowing of tomato in raised bed for nursery seedlings. Do seed treatment with Trichoderma viride @ 4g/kg seeds. The seed treatment must be done 24 hours before sowing of seeds.
- Due to forecast of rain, Chances of fruit rot infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever sucking pests incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. To prevent of soil borne diseases like seedling rot, stem rot and wilt diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Incidence of thrips may observe in chilli, if observe spray Spinosad 45% SC @ 0.4 ml/litre of water or Imidacloprid 17.8% SL @ 0.3 ml/litre of water in clean and dry weather. The present weather conditions are favourable for Upward and downward leafcurl in chilli. To control spray phosalon 3 ml. or diphenhiuron @ 1g./L of water.
- To control fruit and shoot borer in okra, Spray Bacillus thuringiensis 5% W.P @ 1.5 gram/lit of water. Present weather condition favours incidence of white fly and jassid. To control this apply Thiomethoxam @ 0.4g/ litre of water.

Fruits:

Spray mancozeb 2g/litre, to control sigatoka leaf spot in banana. To control papaya yellow mosaic virus, spray Imidachloprid @ 0.5 ml / litre.





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK



(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-516
Date- 30/07/2024

AGROMET ADVISORY BULLETIN FOR CUTTACK SADAR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 15 mm and district received average 13 mm rainfall during the last week. The mean daily maximum and minimum temperature was 29.85°C (-2.37 from normal) and 24.85°C (+0.64 from normal), respectively. The mean relative humidity was 92-94%.

Forecasted period: Today(8.30IST) to 04th August(8.30IST)2024

Weather advisory: Moderate with isolated heavy rainfall is forecasted for tomorrow and light to moderate rainfall is forecasted for other 4 in the block. The day temperature will be 31-33°C and night temperature will be 25-27°C for upcoming 5 days. Maximum and minimum relative humidity will be 79-85% and 51-62% respectively. **There will be partly cloudy to overcast sky for next 5 days.**

Block	Cuttack sadar (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	30/07/2024	31/07/2024	01/08/2024	02/08/2024	03/08/2024
Rainfall(mm)	19	31	11	8	6
T-Max(°C)	33	33	31	31	32
T-Min(°C)	27	27	25	26	26
Cloud Cover(octa)	8	8	8	8	7
RH Max (%)	79	85	83	82	83
RH Min (%)	51	56	62	57	54
Wind Speed (Kmph)	14	13	14	14	10
Wind Direction (deg)	210	248	243	212	259

Agromet Advisory

General advisory: Drain out excess water from rice nurseries, orchards, maize and jute fields. Postpone irrigation, intercultural operations, application of plant protection measures and fertilizers to the standing crops in upcoming 5 days. Keep the harvested produce at safer places. Cover the vegetable seedling with polythene to protect them from heavy shower.

Kharrifrice:(DIRECT SEEDED RICE):

Direct seeded rice are at three to four week stage. As light to moderate amount of rainfall is forecasted in the upcoming days, farmers should go for beushanning. Apply 35 kg urea/ac at beushanning followed by weeding and gap filling. Drain out the excess water at the time of fertilizer application and pond it after one day. If basal fertilizer not applied, apply 18 kg urea, 100kg ssp and 26 kg MOP while beushanning followed by top dressing of 35 kg urea after 7-10 days.

TRANSPLANTED RICE:

Considering the anticipated rainfall, paddy transplanting may be completed before 30th July. Apply 400-gram Chlorantraniliprole 0.4 % G or 800-gram Cartap Hydrochloride 4% GR in the 10 decimal nursery area 7 days before transplanting to manage gall midge, stem borer, caseworm, leaf folder and root knot nematode up to 3 weeks after transplanting.

After rain in dry weather go for foliar spray of Hexaconazole with Plantomycin @ 1.0 g/l to prevent the spread of BLB and sheath blight. If leaf blast is observed in nursery, spray Tebuconazole 50% + Trifloxystrobin 25% WG @ 0.4 g or Isoprothiolane 40EC @ 1.5 ml per litre of water. Repeat the spray at 7-10 days interval.

Suagrcane:

For Red rot and wilt disease, spray the crop with Mancozeb 0.3% or carbendazim 0.1% or Azoxystrobin 18.2% SC/ Difenconazole 11.45SC @ 1ml per liter water in dry days. weather is conducive for temperatures between 29-35 degree Celsius, humidity 60-70 %; this may favour the growth of Whiteflies in sugarcane; spray Thiamethoxam 25 WG @ 125g/ha or apply Chloropyriphos @ 2.5ml/litre water. Present rainy weather favors infestation Wilt disease in Sugarcane, to control this drain out excess water and apply Cupper oxychloride 50.0% 3 gram per liter of water.

VEGETABLES:

- June-july is the best time for sowing of tomato in raised bed for nursery seedlings. Do seed treatment with Trichoderma viride @ 4g/kg seeds. The seed treatment must be done 24 hours before sowing of seeds.
- Due to forecast of rain, Chances of fruit rot infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever sucking pests incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. To prevent of soil borne diseases like seedling rot, stem rot and wilt diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Incidence of thrips may observe in chilli, if observe spray Spinosad 45% SC @ 0.4 ml/litre of water or Imidacloprid 17.8% SL @ 0.3 ml/litre of water in clean and dry weather. The present weather conditions are favourable for Upward and downward leafcurl in chilli. To control spray phosalon 3 ml. or diphenhiuron @ 1g./L of water.
- To control fruit and shoot borer in okra, Spray Bacillus thuringiensis 5% W.P @ 1.5 gram/lit of water. Present weather condition favours incidence of white fly and jassid. To control this apply Thiomethoxam @ 0.4g/ litre of water.

Fruits:

Spray mancozeb 2g/litre, to control sigatoka leaf spot in banana. To control papaya yellow mosaic virus, spray Imidachloprid @ 0.5 ml / litre.





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK



(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-516
Date- 30/07/2024

AGROMET ADVISORY BULLETIN FOR KANTAPADA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 14 mm and district received average 13 mm rainfall during the last week. The mean daily maximum and minimum temperature was 29.85°C (-2.37 from normal) and 24.85°C (+0.64 from normal), respectively. The mean relative humidity was 92-94%.

Forecasted period: Today(8.30IST) to 04th August(8.30IST)2024

Weather advisory: Light to moderate rainfall is forecasted for upcoming 5 days in the block. The day temperature will be 31-33°C and night temperature will be 26-27°C for upcoming 5 days. Maximum and minimum relative humidity will be 75-79% and 49-59% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Kantapada (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	30/07/2024	31/07/2024	01/08/2024	02/08/2024	03/08/2024
Rainfall(mm)	15	25	8	6	6
T-Max(°C)	33	33	31	31	32
T-Min(°C)	27	27	26	26	26
Cloud Cover(octa)	8	8	8	8	8
RH Max (%)	78	79	75	78	77
RH Min (%)	49	54	59	54	52
Wind Speed (Kmph)	15	14	16	16	12
Wind Direction (deg)	208	244	243	212	214

Agromet Advisory

General advisory: Drain out excess water from rice nurseries, orchards, maize and jute fields. Postpone irrigation, intercultural operations, application of plant protection measures and fertilizers to the standing crops in upcoming 5 days. Keep the harvested produce at safer places. Cover the vegetable seedling with polythene to protect them from heavy shower.

Kharifrice:(DIRECT SEEDED RICE):

Direct seeded rice are at three to four week stage. As light to moderate amount of rainfall is forecasted in the upcoming days, farmers should go for beushanning. Apply 35 kg urea/ac at beushanning followed by weeding and gap filling. Drain out the excess water at the time of fertilizer application and pond it after one day. If basal fertilizer not applied, apply 18 kg urea, 100kg ssp and 26 kg MOP while beushanning followed by top dressing of 35 kg urea after 7-10 days.

TRANSPLANTED RICE:

Considering the anticipated rainfall, paddy transplanting may be completed before 30th July. Apply 400-gram Chlorantraniliprole 0.4 % G or 800-gram Cartap Hydrochloride 4% GR in the 10 decimal nursery area 7 days before transplanting to manage gall midge, stem borer, caseworm, leaf folder and root knot nematode up to 3 weeks after transplanting.

After rain in dry weather go for foliar spray of Hexaconazole with Plantomycin@1.0 g/l to prevent the spread of BLB and sheath blight. If leaf blast is observed in nursery, spray Tebuconazole 50% + Trifloxystrobin 25% WG@ 0.4 g or Isoprothiolane 40EC @ 1.5 ml per litre of water. Repeat the spray at 7-10 days interval.

Sugarcane:

For Red rot and wilt disease, spray the crop with Mancozeb 0.3% or carbendazim 0.1% or Azoxystrobin 18.2%SC/ Difenconazole 11.45SC @ 1ml per liter water in dry days. weather is conducive for temperatures between 29-35 degree Celsius, humidity 60-70 %; this may favour the growth of Whiteflies in sugarcane; spray Thiamethoxam 25 WG @ 125g/ha or apply Chloropyriphos @ 2.5ml/litre water. Present rainy weather favors infestation Wilt disease in Sugarcane, to control this drain out excess water and apply Copper oxychloride 50.0% 3 gram per liter of water.

VEGETABLES:

- June-july is the best time for sowing of tomato in raised bed for nursery seedlings. Do seed treatment with Trichoderma viride @ 4g/kg seeds. The seed treatment must be done 24 hours before sowing of seeds.
- Due to forecast of rain, Chances of fruit rot infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever sucking pests incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. To prevent of soil borne diseases like seedling rot, stem rot and wilt diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Incidence of thrips may observe in chilli, if observe spray Spinosad 45% SC @ 0.4 ml/litre of water or Imidacloprid 17.8% SL @ 0.3 ml/litre of water in clean and dry weather. The present weather conditions are favourable for Upward and downward leafcurl in chilli. To control spray phosalon 3 ml. or diphenhiuron @ 1g./L of water.
- To control fruit and shoot borer in okra, Spray Bacillus thuringiensis 5% W.P @1.5 gram/lit of water. Present weather condition favours incidence of white fly and jassid. To control this apply Thiomethoxam@0.4g/ litre of water.

Fruits:

Spray mancozeb 2g/litre, to control sigatoka leaf spot in banana. To control papaya yellow mosaic virus, spray Imidachloprid @ 0.5 ml / litre.





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK



(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-516

Date- 30/07/2024

AGROMET ADVISORY BULLETIN FOR MAHANGA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 11 mm and district received average 13 mm rainfall during the last week. The mean daily maximum and minimum temperature was 29.85°C (-2.37 from normal) and 24.85°C (+0.64 from normal), respectively. The mean relative humidity was 92-94%.

Forecasted period: Today(8.30IST) to 04th August(8.30IST)2024

Weather advisory: Moderate with isolated heavy rainfall is forecasted for tomorrow and light to moderate rainfall is forecasted for other 4 in the block. The day temperature will be and it will be 30-33°C and night temperature will be 26-27°C for upcoming 5 days. Maximum and minimum relative humidity will be 79-84% and 50-59% respectively. **There will be partly cloudy to overcast sky for next 5 days.**

Block	Mahanga (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	30/07/2024	31/07/2024	01/08/2024	02/08/2024	03/08/2024
Rainfall(mm)	25	31	11	7	6
T-Max(°C)	33	33	31	30	32
T-Min(°C)	27	27	26	26	26
Cloud Cover(octal)	8	8	8	8	8
RH Max (%)	79	84	81	81	82
RH Min (%)	50	54	59	57	52
Wind Speed (Kmph)	16	14	16	15	10
Wind Direction (deg)	216	244	243	207	270

Agromet Advisory

General advisory: Drain out excess water from rice nurseries, orchards, maize and jute fields. Postpone irrigation, intercultural operations, application of plant protection measures and fertilizers to the standing crops in upcoming 5 days. Keep the harvested produce at safer places. Cover the vegetable seedling with polythene to protect them from heavy shower.

Kharifrice:(DIRECT SEEDED RICE):

Direct seeded rice are at three to four week stage. As light to moderate amount of rainfall is forecasted in the upcoming days, farmers should go for beushanning. Apply 35 kg urea/ac at beushanning followed by weeding and gap filling. Drain out the excess water at the time of fertilizer application and pond it after one day. If basal fertilizer not applied, apply 18 kg urea, 100kg ssp and 26 kg MOP while beushanning followed by top dressing of 35 kg urea after 7-10 days.

TRANSPLANTED RICE:

Considering the anticipated rainfall, paddy transplanting may be completed before 30th July. Apply 400-gram Chlorantraniliprole 0.4 % G or 800-gram Cartap Hydrochloride 4% GR in the 10 decimal nursery area 7 days before transplanting to manage gall midge, stem borer, caseworm, leaf folder and root knot nematode up to 3 weeks after transplanting.

After rain in dry weather go for foliar spray of Hexaconazole with Plantomycin@1.0 g/l to prevent the spread of BLB and sheath blight. If leaf blast is observed in nursery, spray Tebuconazole 50% + Trifloxystrobin 25% WG@ 0.4 g or Isoprothiolane 40EC @ 1.5 ml per litre of water. Repeat the spray at 7-10 days interval.

Sugarcane:

For **Red rot and wilt disease**, spray the crop with Mancozeb 0.3% or carbendazim 0.1% or Azoxystrobin 18.2%SC/ Difenconazole 11.45SC @ 1ml per liter water in dry days. weather is conducive for temperatures between 29-35 degree Celsius, humidity 60-70 %; this may favour the growth of Whiteflies in sugarcane;; spray Thiamethoxam 25 WG @ 125g/ha or apply Chloropyriphos @ 2.5ml/litre water. Present rainy weather favors infestation Wilt disease in Sugarcane, to control this drain out excess water and apply Copper oxychloride 50.0% 3 gram per liter of water.

VEGETABLES:

- June-july is the best time for sowing of tomato in raised bed for nursery seedlings. Do seed treatment with Trichoderma viride @ 4g/kg seeds. The seed treatment must be done 24 hours before sowing of seeds.
- Due to forecast of rain, Chances of **fruit rot** infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever **sucking pests** incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. **To prevent of soil borne diseases like seedling rot, stem rot and wilt diseases apply 3g of Copper oxychloride per litre of water around the main stem.**
- Incidence of **thrips** may observe in chilli, if observe spray Spinosad 45% SC @ 0.4 ml/litre of water or Imidacloprid 17.8% SL @ 0.3 ml/litre of water in clean and dry weather. The present weather conditions are favourable for Upward and downward leafcurl in chilli. To control spray phosalon 3 ml. or diphenhiuron @ 1g./L of water.
- To control **fruit and shoot borer** in okra, Spray Bacillus thuringiensis 5% W.P @ 1.5 gram/lit of water. Present weather condition favours incidence of white fly and jassid. To control this apply Thiomethoxam @ 0.4g/ litre of water.

Fruits:

Spray mancozeb 2g/litre, to control sigatoka leaf spot in **banana**. To control **papaya yellow mosaic virus**, spray Imidachloprid @ 0.5 ml / litre.

DrRanjan Kumar Mohanta
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK



(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-516

Date- 30/07/2024

AGROMET ADVISORY BULLETIN FOR NARASINGHPUR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 10 mm and district received average 13 mm rainfall during the last week. The mean daily maximum and minimum temperature was 29.85°C (-2.37 from normal) and 24.85°C (+0.64 from normal), respectively. The mean relative humidity was 92-94%.

Forecasted period: Today(8.30IST) to 04th August(8.30IST)2024

Weather advisory: Moderate with isolated heavy rainfall is forecasted for tomorrow and light to moderate rainfall is forecasted for other 4 in the block. The day temperature will be 29-32°C and night temperature will be 23-25°C for upcoming 5 days. Maximum and minimum relative humidity will be 83-91% and 55-64% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Narasinghpur (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	30/07/2024	31/07/2024	01/08/2024	02/08/2024	03/08/2024
Rainfall(mm)	30	33	11	7	6
T-Max(°C)	32	32	30	29	30
T-Min(°C)	25	24	23	23	23
Cloud Cover(/octa)	8	8	8	8	7
RH Max (%)	84	91	83	83	88
RH Min (%)	55	58	64	61	61
Wind Speed (Kmph)	8	8	9	12	6
Wind Direction (deg)	243	248	225	225	225

Agromet Advisory

General advisory: Drain out excess water from rice nurseries, orchards, maize and jute fields. Postpone irrigation, intercultural operations, application of plant protection measures and fertilizers to the standing crops in upcoming 5 days. Keep the harvested produce at safer places. Cover the vegetable seedling with polythene to protect them from heavy shower.

Kharifrice:(DIRECT SEEDED RICE):

Direct seeded rice are at three to four week stage. As light to moderate amount of rainfall is forecasted in the upcoming days, farmers should go for beushanning. Apply 35 kg urea/ac at beushanning followed by weeding and gap filling. Drain out the excess water at the time of fertilizer application and pond it after one day. If basal fertilizer not applied, apply 18 kg urea, 100kg ssp and 26 kg MOP while beushanning followed by top dressing of 35 kg urea after 7-10 days.

TRANSPLANTED RICE:

Considering the anticipated rainfall, paddy transplanting may be completed before 30th July. Apply 400-gram Chlorantraniliprole 0.4 % G or 800-gram Cartap Hydrochloride 4% GR in the 10 decimal nursery area 7 days before transplanting to manage gall midge, stem borer, caseworm, leaf folder and root knot nematode up to 3 weeks after transplanting.

After rain in dry weather go for foliar spray of Hexaconazole with Plantomycin @ 1.0 g/l to prevent the spread of BLB and sheath blight. If leaf blast is observed in nursery, spray Tebuconazole 50% + Trifloxystrobin 25% WG @ 0.4 g or Isoprothiolane 40EC @ 1.5 ml per litre of water. Repeat the spray at 7-10 days interval.

Maize:

Stem fly in Maize infects the plants from sowing to 5 weeks after sowing, temperature between 25-30 deg. celsius and humidity above 60% favour pest incidence. In vegetative stage to control stem fly spray Methyl demeton @ 1 ml/lit or Dimethoate @ 2ml/lit or apply Carbofuran 3G granuals @ 14 kg/acre. High humid conditions, temperature between 18-27 Deg. Celsius favours the occurrence of leaf blight in Maize. To control spray Azoxystrobin + Mancozeb 300 g /ac.

VEGETABLES:

- June-july is the best time for sowing of tomato in raised bed for nursery seedlings. Do seed treatment with Trichoderma viride @ 4g/kg seeds. The seed treatment must be done 24 hours before sowing of seeds.
- Due to forecast of rain, Chances of fruit rot infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever sucking pests incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. To prevent of soil borne diseases like seedling rot, stem rot and wilt diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Incidence of thrips may observe in chilli, if observe spray Spinosad 45% SC @ 0.4 ml/litre of water or Imidacloprid 17.8% SL @ 0.3 ml/litre of water in clean and dry weather. The present weather conditions are favourable for Upward and downward leafcurl in chilli. To control spray phosalon 3 ml. or diphenhiuron @ 1g./L of water.
- To control fruit and shoot borer in okra, Spray Bacillus thuringiensis 5% W.P @ 1.5 gram/lit of water. Present weather condition favours incidence of white fly and jassid. To control this apply Thiomethoxam @ 0.4g/ litre of water.

Fruits:

Spray mancozeb 2g/litre, to control sigatoka leaf spot in banana. To control papaya yellow mosaic virus, spray Imidachloprid @ 0.5 ml / litre.

DrRanjan Kumar Mohanta
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK



(A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-516
 Date- 30/07/2024

AGROMET ADVISORY BULLETIN FOR NIALI

District Past week average weather condition and Block past week average Rainfall condition

The block received average 12 mm and district received average 13 mm rainfall during the last week. The mean daily maximum and minimum temperature was 29.85°C (-2.37 from normal) and 24.85°C (+0.64 from normal), respectively. The mean relative humidity was 92-94%.

Forecasted period: Today(8.30IST) to 04th August(8.30IST)2024

Weather advisory: Light to moderate rainfall is forecasted for upcoming 5 days in the block. The day temperature will be 31-33°C and night temperature will be 26-27°C for upcoming 5 days. Maximum and minimum relative humidity will be 73-78% and 49-58% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Niali (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	30/07/2024	31/07/2024	01/08/2024	02/08/2024	03/08/2024
Rainfall(mm)	14	18	8	7	6
T-Max(°C)	32	33	31	31	32
T-Min(°C)	27	27	27	26	26
Cloud Cover(octa)	8	8	8	8	8
RH Max (%)	78	78	73	77	75
RH Min (%)	49	54	58	54	51
Wind Speed(Kmph)	16	16	18	18	13
Wind Direction(deg)	208	244	243	212	225

Agromet Advisory

General advisory: Drain out excess water from rice nurseries, orchards, maize and jute fields. Postpone irrigation, intercultural operations, application of plant protection measures and fertilizers to the standing crops in upcoming 5 days. Keep the harvested produce at safer places. Cover the vegetable seedling with polythene to protect them from heavy shower.

Kharifrice:(DIRECT SEEDED RICE):

Direct seeded rice are at three to four week stage. As light to moderate amount of rainfall is forecasted in the upcoming days, farmers should go for beushanning. Apply 35 kg urea/ac at beushanning followed by weeding and gap filling. Drain out the excess water at the time of fertilizer application and pond it after one day. If basal fertilizer not applied, apply 18 kg urea, 100kg ssp and 26 kg MOP while beushanning followed by top dressing of 35 kg urea after 7-10 days.

TRANSPLANTED RICE:

Considering the anticipated rainfall, paddy transplanting may be completed before 30th July. Apply 400-gram Chlorantranilprole 0.4 % G or 800-gram Cartap Hydrochloride 4% GR in the 10 decimal nursery area 7 days before transplanting to manage gall midge, stem borer, caseworm, leaf folder and root knot nematode up to 3 weeks after transplanting.

After rain in dry weather go for foliar spray of Hexaconazole with Plantomycin@1.0 g/l to prevent the spread of BLB and sheath blight. If leaf blast is observed in nursery, spray Tebuconazole 50% + Trifloxystrobin 25% WG@ 0.4 g or Isoprothiolane 40EC @ 1.5 ml per litre of water. Repeat the spray at 7-10 days interval.

Sugarcane:

For Red rot and wilt disease, spray the crop with Mancozeb 0.3% or carbendazim 0.1% or Azoxystrobin 18.2%SC/ Difenconazole 11.45SC @ 1ml per liter water in dry days. weather is conducive for temperatures between 29-35 degree Celsius, humidity 60-70 %; this may favour the growth of Whiteflies in sugarcane; spray Thiamethoxam 25 WG @ 125g/ha or apply Chloropyrifos @ 2.5ml/litre water. Present rainy weather favors infestation Wilt disease in Sugarcane, to control this drain out excess water and apply Copper oxychloride 50.0% 3 gram per liter of water.

VEGETABLES:

- June-july is the best time for sowing of tomato in raised bed for nursery seedlings. Do seed treatment with Trichoderma viride @ 4g/kg seeds. The seed treatment must be done 24 hours before sowing of seeds.
- Due to forecast of rain, Chances of fruit rot infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever sucking pests incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. To prevent of soil borne diseases like seedling rot, stem rot and wilt diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Incidence of thrips may observe in chilli, if observe spray Spinosad 45% SC @ 0.4 ml/litre of water or Imidacloprid 17.8% SL @ 0.3 ml/litre of water in clean and dry weather. The present weather conditions are favourable for Upward and downward leafcurl in chilli. To control spray phosalon 3 ml. or diphenhiuron @ 1g./L of water.
- To control fruit and shoot borer in okra, Spray Bacillus thuringiensis 5% W.P @1.5 gram/lit of water. Present weather condition favours incidence of white fly and jassid. To control this apply Thiomethoxam@0.4g/ litre of water.

Fruits:

Spray mancozeb 2g/litre, to control sigatoka leaf spot in banana. To control papaya yellow mosaic virus, spray Imidachloprid @ 0.5 ml / litre.





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK



(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-516

Date- 30/07/2024

AGROMET ADVISORY BULLETIN FOR NISCHINTAKOILI

District Past week average weather condition and Block past week average Rainfall condition

The block received average 09 mm and district received average 13 mm rainfall during the last week. The mean daily maximum and minimum temperature was 29.85°C (-2.37 from normal) and 24.85°C (+0.64 from normal), respectively. The mean relative humidity was 92-94%.

Forecasted period: Today(8.30IST) to 04th August(8.30IST)2024

Weather advisory: *Light to moderate rainfall is forecasted for upcoming 5 days in the block.* The day temperature will be 30-33°C and night temperature will be 26-27°C for upcoming 5 days. Maximum and minimum relative humidity will be 79-83% and 49-58% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Nischintakoili (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	30/07/2024	31/07/2024	01/08/2024	02/08/2024	03/08/2024
Rainfall(mm)	15	19	9	6	6
T-Max(°C)	32	33	31	30	32
T-Min(°C)	27	27	26	26	26
Cloud Cover(octa)	8	8	8	8	8
RH Max (%)	80	83	79	81	81
RH Min (%)	49	54	58	56	51
Wind Speed (Kmph)	15	14	16	15	11
Wind Direction (deg)	208	244	243	207	270

Agromet Advisory

General advisory: *Drain out excess water from rice nurseries, orchards, maize and jute fields. Postpone irrigation, intercultural operations, application of plant protection measures and fertilizers to the standing crops in upcoming 5 days. Keep the harvested produce at safer places. Cover the vegetable seedling with polythene to protect them from heavy shower.*

Kharifrice:(DIRECT SEEDED RICE):

Direct seeded rice are at three to four week stage. As light to moderate amount of rainfall is forecasted in the upcoming days, farmers should go for beushanning. Apply 35 kg urea/ac at beushanning followed by weeding and gap filling. Drain out the excess water at the time of fertilizer application and pond it after one day. If basal fertilizer not applied, apply 18 kg urea, 100kg ssp and 26 kg MOP while beushanning followed by top dressing of 35 kg urea after 7-10 days.

TRANSPLANTED RICE:

Considering the anticipated rainfall, paddy transplanting may be completed before 30th July. Apply 400-gram Chlorantraniliprole 0.4 % G or 800-gram Cartap Hydrochloride 4% GR in the 10 decimal nursery area 7 days before transplanting to manage gall midge, stem borer, caseworm, leaf folder and root knot nematode up to 3 weeks after transplanting.

After rain in dry weather go for foliar spray of Hexaconazole with Plantomycin@1.0 g/l to prevent the spread of BLB and sheath blight. If leaf blast is observed in nursery, spray Tebuconazole 50% + Trifloxystrobin 25% WG@ 0.4 g or Isoprothiolane 40EC @ 1.5 ml per litre of water. Repeat the spray at 7-10 days interval.

Sugarcane:

For **Red rot and wilt disease**, spray the crop with Mancozeb 0.3% or carbendazim 0.1% or Azoxystrobin 18.2%SC/ Difenconazole 11.45SC @ 1ml per liter water in dry days. weather is conducive for temperatures between 29-35 degree Celsius, humidity 60-70 %; this may favour the growth of Whiteflies in sugarcane;; spray Thiamethoxam 25 WG @ 125g/ha or apply Chloropyriphos @ 2.5ml/litre water. Present rainy weather favors infestation Wilt disease in Sugarcane, to control this drain out excess water and apply Cupper oxychloride 50.0% 3 gram per liter of water.

VEGETABLES:

- June-july is the best time for sowing of tomato in raised bed for nursery seedlings. Do seed treatment with Trichoderma viride @ 4g/kg seeds. The seed treatment must be done 24 hours before sowing of seeds.
- Due to forecast of rain, Chances of **fruit rot** infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever **sucking pests** incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. **To prevent of soil borne diseases like seedling rot, stem rot and wilt diseases apply 3g of Copper oxychloride per litre of water around the main stem.**
- Incidence of **thrips** may observe in chilli, if observe spray Spinosad 45% SC @ 0.4 ml/litre of water or Imidacloprid 17.8% SL @ 0.3 ml/litre of water in clean and dry weather. The present weather conditions are favourable for Upward and downward leaf curl in chilli. To control spray phosalon 3 ml. or diphenthiuron @ 1g./L of water.
- To control **fruit and shoot borer** in okra, Spray Bacillus thuringiensis 5% W.P @ 1.5 gram/lit of water. Present weather condition favours incidence of white fly and jassid. To control this apply Thiomethoxam @ 0.4g/ litre of water.

Fruits:

Spray mancozeb 2g/litre, to control sigatoka leaf spot in **banana**. To control **papaya yellow mosaic virus**, spray Imidachloprid @ 0.5 ml / litre.





**GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK**



(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-516

Date- 30/07/2024

AGROMET ADVISORY BULLETIN FOR SALEPUR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 13 mm and district received average 13 mm rainfall during the last week. The mean daily maximum and minimum temperature was 29.85°C (-2.37 from normal) and 24.85°C (+0.64 from normal), respectively. The mean relative humidity was 92-94%.

Forecasted period: Today(8.30IST) to 04th August(8.30IST)2024

Weather advisory: Moderate with isolated heavy rainfall is forecasted for tomorrow and light to moderate rainfall is forecasted for other 4 in the block. The day temperature will be 30-33°C and night temperature will be 26-27°C for upcoming 5 days. Maximum and minimum relative humidity will be 79-83% and 50-59% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Salepur (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	30/07/2024	31/07/2024	01/08/2024	02/08/2024	03/08/2024
Rainfall(mm)	20	31	11	6	6
T-Max(°C)	32	33	31	30	32
T-Min(°C)	27	27	26	26	26
Cloud Cover(octa)	8	8	8	8	8
RH Max (%)	79	83	80	81	82
RH Min (%)	50	54	59	56	52
Wind Speed (Kmph)	15	14	16	15	10
Wind Direction (deg)	216	244	243	207	270

Agromet Advisory

General advisory: Drain out excess water from rice nurseries, orchards, maize and jute fields. Postpone irrigation, intercultural operations, application of plant protection measures and fertilizers to the standing crops in upcoming 5 days. Keep the harvested produce at safer places. Cover the vegetable seedling with polythene to protect them from heavy shower.

Kharifrice:(DIRECT SEEDED RICE):

Direct seeded rice are at three to four week stage. As light to moderate amount of rainfall is forecasted in the upcoming days, farmers should go for beushanning. Apply 35 kg urea/ac at beushanning followed by weeding and gap filling. Drain out the excess water at the time of fertilizer application and pond it after one day. If basal fertilizer not applied, apply 18 kg urea, 100kg ssp and 26 kg MOP while beushanning followed by top dressing of 35 kg urea after 7-10 days.

TRANSPLANTED RICE:

Considering the anticipated rainfall, paddy transplanting may be completed before 30th July. Apply 400-gram Chlorantranilprole 0.4 % G or 800-gram Cartap Hydrochloride 4% GR in the 10 decimal nursery area 7 days before transplanting to manage gall midge, stem borer, caseworm, leaf folder and root knot nematode up to 3 weeks after transplanting.

After rain in dry weather go for foliar spray of Hexaconazole with Plantomycin @ 1.0 g/l to prevent the spread of BLB and sheath blight. If leaf blast is observed in nursery, spray Tebuconazole 50% + Trifloxystrobin 25% WG @ 0.4 g or Isoprothiolane 40EC @ 1.5 ml per litre of water. Repeat the spray at 7-10 days interval.

Maize:

Stem fly in Maize infects the plants from sowing to 5 weeks after sowing, temperature between 25-30 deg. celsius and humidity above 60% favour pest incidence. In vegetative stage to control stem fly spray Methyl demeton @ 1 ml/lit or Dimethoate @ 2ml/lit or apply Carbofuran 3G granuals @ 14 kg/acre. High humid conditions, temperature between 18-27 Deg. Celsius favours the occurrence of leaf blight in Maize. To control spray Azoxystrobin + Mancozeb 300 g /ac.

VEGETABLES:

- June-july is the best time for sowing of tomato in raised bed for nursery seedlings. Do seed treatment with Trichoderma viride @ 4g/kg seeds. The seed treatment must be done 24 hours before sowing of seeds.
- Due to forecast of rain, Chances of fruit rot infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever sucking pests incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. To prevent of soil borne diseases like seedling rot, stem rot and wilt diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Incidence of thrips may observe in chilli, if observe spray Spinosad 45% SC @ 0.4 ml/litre of water or Imidacloprid 17.8% SL @ 0.3 ml/litre of water in clean and dry weather. The present weather conditions are favourable for Upward and downward leafcurl in chilli. To control spray phosalon 3 ml. or diphenhiuron @ 1g./L of water.
- To control fruit and shoot borer in okra, Spray Bacillus thuringiensis 5% W.P @ 1.5 gram/lit of water. Present weather condition favours incidence of white fly and jassid. To control this apply Thiomethoxam @ 0.4g/ litre of water.

Fruits:

Spray mancozeb 2g/litre, to control sigatoka leaf spot in banana. To control papaya yellow mosaic virus, spray Imidachloprid @ 0.5 ml / litre.

DrRanjan Kumar Mohanta
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK



(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-516
Date- 30/07/2024

AGROMET ADVISORY BULLETIN FOR TANGI-CHOUDWAR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 12 mm and district received average 13 mm rainfall during the last week. The mean daily maximum and minimum temperature was 29.85°C (-2.37 from normal) and 24.85°C (+0.64 from normal), respectively. The mean relative humidity was 92-94%.

Forecasted period: Today(8.30IST) to 04th August(8.30IST)2024

Weather advisory: Moderate with isolated heavy rainfall is forecasted for tomorrow and light to moderate rainfall is forecasted for other 4 in the block. The day temperature will be 33-36°C and night temperature will be 29-30°C for upcoming 5 days. Maximum and minimum relative humidity will be 74-80% and 53-61% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Tangi-Choudwar (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	30/07/2024	31/07/2024	01/08/2024	02/08/2024	03/08/2024
Rainfall(mm)	23	31	11	6	6
T-Max(°C)	34	36	34	33	35
T-Min(°C)	30	30	30	29	29
Cloud Cover(octa)	8	8	8	8	8
RH Max (%)	80	77	74	79	77
RH Min (%)	57	56	61	60	53
Wind Speed (Kmph)	19	19	21	21	15
Wind Direction (deg)	205	236	238	204	205

Agromet Advisory

General advisory: Drain out excess water from rice nurseries, orchards, maize and jute fields. Postpone irrigation, intercultural operations, application of plant protection measures and fertilizers to the standing crops in upcoming 5 days. Keep the harvested produce at safer places. Cover the vegetable seedling with polythene to protect them from heavy shower.

Kharifrice:(DIRECT SEEDED RICE):

Direct seeded rice are at three to four week stage. As light to moderate amount of rainfall is forecasted in the upcoming days, farmers should go for beushanning. Apply 35 kg urea/ac at beushanning followed by weeding and gap filling. Drain out the excess water at the time of fertilizer application and pond it after one day. If basal fertilizer not applied, apply 18 kg urea, 100kg ssp and 26 kg MOP while beushanning followed by top dressing of 35 kg urea after 7-10 days.

TRANSPLANTED RICE:

Considering the anticipated rainfall, paddy transplanting may be completed before 30th July. Apply 400-gram Chlorantraniliprole 0.4 % G or 800-gram Cartap Hydrochloride 4% GR in the 10 decimal nursery area 7 days before transplanting to manage gall midge, stem borer, caseworm, leaf folder and root knot nematode up to 3 weeks after transplanting.

After rain in dry weather go for foliar spray of Hexaconazole with Plantomycin @ 1.0 g/l to prevent the spread of BLB and sheath blight. If leaf blast is observed in nursery, spray Tebuconazole 50% + Trifloxystrobin 25% WG @ 0.4 g or Isoprothiolane 40EC @ 1.5 ml per litre of water. Repeat the spray at 7-10 days interval.

Maize:

Stem fly in Maize infects the plants from sowing to 5 weeks after sowing, temperature between 25-30 deg. celsius and humidity above 60% favour pest incidence. In vegetative stage to control stem fly spray Methyl demeton @ 1 ml/lit or Dimethoate @ 2ml/lit or apply Carbofuran 3G granuals @ 14 kg/acre. High humid conditions, temperature between 18-27 Deg. Celsius favours the occurrence of leaf blight in Maize. To control spray Azoxystrobin + Mancozeb 300 g /ac.

VEGETABLES:

- June-july is the best time for sowing of tomato in raised bed for nursery seedlings. Do seed treatment with Trichoderma viride @ 4g/kg seeds. The seed treatment must be done 24 hours before sowing of seeds.
- Due to forecast of rain, Chances of fruit rot infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever sucking pests incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. To prevent of soil borne diseases like seedling rot, stem rot and wilt diseases apply 3g of Copper oxychloride per litre of water around the main stem.
- Incidence of thrips may observe in chilli, if observe spray Spinosad 45% SC @ 0.4 ml/litre of water or Imidacloprid 17.8% SL @ 0.3 ml/litre of water in clean and dry weather. The present weather conditions are favourable for Upward and downward leafcurl in chilli. To control spray phosalon 3 ml. or diphenhiuron @ 1g./L of water.
- To control fruit and shoot borer in okra, Spray Bacillus thuringiensis 5% W.P @ 1.5 gram/lit of water. Present weather condition favours incidence of white fly and jassid. To control this apply Thiomethoxam @ 0.4g/ litre of water.

Fruits:

Spray mancozeb 2g/litre, to control sigatoka leaf spot in banana. To control papaya yellow mosaic virus, spray Imidachloprid @ 0.5 ml / litre.

DrRanjan Kumar Mohanta
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena
SMS (Agrometeorology)
Contact No-9861445080/7008453427



GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK



(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-516

Date- 30/07/2024

AGROMET ADVISORY BULLETIN FOR TIGIRIA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 16 mm and district received average 13 mm rainfall during the last week. The mean daily maximum and minimum temperature was 29.85°C (-2.37 from normal) and 24.85°C (+0.64 from normal), respectively. The mean relative humidity was 92-94%.

Forecasted period: Today(8.30IST) to 04th August(8.30IST)2024

Weather advisory: Moderate with isolated heavy rainfall is forecasted for tomorrow and light to moderate rainfall is forecasted for other 4 in the block. The day temperature will be 31-33°C and night temperature will be 24-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 83-88% and 54-64% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Tigiria (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	30/07/2024	31/07/2024	01/08/2024	02/08/2024	03/08/2024
Rainfall(mm)	26	32	11	6	6
T-Max(°C)	33	32	31	31	31
T-Min(°C)	26	26	24	25	24
Cloud Cover(octa)	8	8	8	8	7
RH Max (%)	83	88	85	83	86
RH Min (%)	54	59	64	60	58
Wind Speed (Kmph)	11	10	11	13	8
Wind Direction (deg)	248	248	238	212	216

Agro met Advisory

General advisory: Drain out excess water from rice nurseries, orchards, maize and jute fields. Postpone irrigation, intercultural operations, application of plant protection measures and fertilizers to the standing crops in upcoming 5 days. Keep the harvested produce at safer places. Cover the vegetable seedling with polythene to protect them from heavy shower.

Khharifrice:(DIRECT SEEDED RICE):

Direct seeded rice are at three to four week stage. As light to moderate amount of rainfall is forecasted in the upcoming days, farmers should go for beushanning. Apply 35 kg urea/ac at beushanning followed by weeding and gap filling. Drain out the excess water at the time of fertilizer application and pond it after one day. If basal fertilizer not applied, apply 18 kg urea, 100kg ssp and 26 kg MOP while beushanning followed by top dressing of 35 kg urea after 7-10 days.

TRANSPLANTED RICE:

Considering the anticipated rainfall, paddy transplanting may be completed before 30th July. Apply 400-gram Chlorantranilprole 0.4 % G or 800-gram Cartap Hydrochloride 4% GR in the 10 decimal nursery area 7 days before transplanting to manage gall midge, stem borer, caseworm, leaf folder and root knot nematode up to 3 weeks after transplanting.

After rain in dry weather go for foliar spray of Hexaconazole with Plantomycin@1.0 g/l to prevent the spread of BLB and sheath blight. If leaf blast is observed in nursery, spray Tebuconazole 50% + Trifloxystrobin 25% WG@ 0.4 g or Isoprothiolane 40EC @ 1.5 ml per litre of water. Repeat the spray at 7-10 days interval.

Maize:

Stem fly in Maize infects the plants from sowing to 5 weeks after sowing, temperature between 25-30 deg. celsius and humidity above 60% favour pest incidence. In vegetative stage to control stem fly spray Methyl demeton @1 ml/lit or Dimethoate @ 2ml/lit or apply Carbofuran 3G granuals @ 14 kg/acre. High humid conditions, temperature between 18-27 Deg. Celsius favours the occurrence of **leaf blight** in Maize. To control spray Azoxystrobin + Mancozeb 300 g /ac.

VEGETABLES:

- June-july is the best time for sowing of tomato in raised bed for nursery seedlings. Do seed treatment with Trichoderma viride @ 4g/kg seeds. The seed treatment must be done 24 hours before sowing of seeds.
- Due to forecast of rain, Chances of **fruit rot** infestation in chilli and brinjal, if observe spray Carbendazim 50% WP @ 2 g/litre of water or Copper oxychloride 50% WP @ 3 g/litre of water. If necessary spray second and third round at 15 days interval.
- Brinjal, tomato and chilli seedlings should be planted in the main field. Whenever **sucking pests** incidence was observed spray 5 ml Neem oil or Thiamethoxam 0.2 g. per litre of water. **To prevent of soil borne diseases like seedling rot, stem rot and wilt diseases apply 3g of Copper oxychloride per litre of water around the main stem.**
- Incidence of **thrips** may observe in chilli, if observe spray Spinosad 45% SC @ 0.4 ml/litre of water or Imidacloprid 17.8% SL @ 0.3 ml/litre of water in clean and dry weather. The present weather conditions are favourable for Upward and downward leaf curl in chilli. To control spray phosalon 3 ml. or diphenhiuron @ 1g./L of water.
- To control **fruit and shoot borer** in okra, Spray Bacillus thuringiensis 5% W.P @ 1.5 gram/lit of water. Present weather condition favours incidence of white fly and jassid. To control this apply Thiomethoxam @ 0.4g/ litre of water.

Fruits:

Spray mancozeb 2g/litre, to control sigatoka leaf spot in **banana**. To control **papaya yellow mosaic virus**, spray Imidachloprid @ 0.5 ml / litre.











GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-516
Date- 30/07/2024

				
MEGHDOOT APP	DAMINI APP	FACEBOOK (Damu Cuttack)	BLOCK SPECIFIC WHAT'S APP GROUPS	@DAMUCUTTACK

- Kindly install and use **“Meghdoot mobile App”** for Cuttack district weather forecast, warnings and weather based agromet advisory services.
<https://play.google.com/store/apps/details?id=com.aas.meghdoot>
- Kindly install and use **“Damini mobile App”** for location specific for lightening forecast and warnings.
<https://play.google.com/store/apps/details?id=com.lightening.live.damini>
- Follow the **block specific whats app groups** and join in your block wise groups for getting weather based agromet advisory services twice in a week (Tuesday and Friday) and other weather warning based farm management advisories.
- Follow the **Official Website links** to get the weather based agromet advisory service bulletins (**English & Odia**) twice in a week (Tuesday and Friday).

1	ICAR-NRRI, Cuttack, Odisha	https://icar-nrri.in/agromate/	
2	Agrimet division, IMD, Pune	https://imdagrmet.gov.in/imdproject/AGIndex.php	