




# *Cuttack District Block Wise Agro-met Advisory Service Bulletin*

*Friday, 18<sup>th</sup> October, 2024  
 (For the period 18<sup>th</sup> to 22<sup>th</sup> October,  
 2024)*





*Issued by  
 District Agro-met Unit,  
 KrishiVigyan Kendra Cuttack*





**STANDARD PRECIPITATION INDEX AND NDVI MAP**

Standard Precipitation Index (SPI)	Normalized difference vegetation Index (NDVI)
<div style="text-align: center;"> <p><b>MINISTRY OF EARTH SCIENCES</b>  <b>INDIA METEOROLOGICAL DEPARTMENT</b>  <b>HYDROMET SECTION, CRS PUNE</b></p> <p><b>STANDARDIZED PRECIPITATION INDEX</b>  <b>FOR THE PERIOD</b>  <b>05 SEPTEMBER TO 02 OCTOBER 2024</b></p>  </div>	<div style="text-align: center;"> <p><b>NOAA/VIIRS/BLENDED NDVI Composite ending on Week no 39</b>  <b>(23.09.2024 to 29.09.2024) over Agricultural region</b></p> <p><b>Odisha</b></p>  </div> <p style="text-align: center;">Agriculture vigour is moderate at isolated places of the state. NDVI could not be captured over remaining places due dense cloud cover or rainfall.</p>
<p><i>As per SPI value (cumulative 4weeks - 05 September to 02 October 2024), mildly wet condition was there in Cuttack district.</i></p>	<p><i>According to NDVI map (23.09.2024 to 29.09.2024) the agriculture vigor was light to moderate over most of the blocks of Cuttack district.</i></p>

<b>LEGEND (Medium range block level rainfall forecast)</b>		
	<b>No Rainfall</b>	<b>00 mm</b>
	<b>Very light rainfall</b>	<b>01-05 mm</b>
	<b>Light rainfall</b>	<b>06-10 mm</b>
	<b>Light to moderate rainfall</b>	<b>11-30 mm</b>
	<b>Moderate to heavy rainfall</b>	<b>31-65 mm</b>
	<b>Heavy to very heavy rainfall</b>	<b>66-115 mm</b>
	<b>Extremely heavy rainfall</b>	<b>&gt;116 mm</b>

<b>LEGEND (Past week temperature )</b>		
Color code	Temperature	Departure
	<b>Below Normal</b>	<b>&lt; 2°C from normal</b>
	<b>Normal</b>	<b>-2°C to +2°C from normal</b>
	<b>Above Normal</b>	<b>&gt;2°C from normal</b>



**DrRanjan Kumar Mohanta**  
 (Nodal officer)

**Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area**





**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**











(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-536  
Date- 18/10/2024

**Weather (Past and Forecasted) based Disease and pest in Major Crops and vegetables**

Sl No	Crop	Stage and condition of the crop	Anticipated Disease and pest	
1	Rice	Panicle initiation to flowering		Leaf streak
				Mealy bug
				Hispa
				Caseworm
				BPH
				Gundhi Bug
			2	Sugarcane
	Scales			
	White fly			



**DrRanjan Kumar Mohanta**  
(Nodal officer)

**Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area**















**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**  
 (A Unit of ICAR-National Rice Research Institute)  
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



**Bulletin No.-536**  
**Date- 18/10/2024**

3	Maize	Cob development stage		Aphid/thrips
				Stem borer
4	Vegetables			Aphid
				Leaf hopper
				White fly
5	Cabbage/Cauliflower	Seedling to vegetative Stage		Damping off
				White fly
				Tobacco caterpillar



**DrRanjan Kumar Mohanta**  
 (Nodal officer)

**Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area**





**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**



(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-536  
Date- 18/10/2024

**AGROMET ADVISORY BULLETIN FOR ATHAGAD**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 33.71°C (+1.03 from normal) and 24.98°C (+2.43 from normal), respectively. The mean relative humidity was 75-81%.

**Forecasted period: Today(8.30IST) to 23<sup>rd</sup> October (8.30IST)2024**

**Weather advisory:** Light to moderate rainfall/thundershower is forecasted for today to Sunday and dry weather is forecasted for other 2days in the block. The day temperature will be 32-34°C and night temperature will be 23-27°C for upcoming 5 days. Maximum and minimum relative humidity will be 78-94% and 48-63% respectively. There will be partly cloudy to overcastsky for next 5days.

Block	Athagad (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	18/10/2024	19/10/2024	20/10/2024	21/10/2024	22/10/2024
Rainfall (mm)	10	15	06	00	00
T-Max (°C)	34	34	32	34	34
T-Min (°C)	26	27	25	24	23
Cloud Cover (octa)	7	7	5	2	4
RH Max (%)	94	93	92	86	78
RH Min (%)	63	62	63	50	48
Wind Speed (kmph)	7	5	6	6	7
Wind Direction (deg)	153	150	23	342	45

**Agromet Advisory**

**General advisory:** As slight increase in temperature condition is forecasted in between upcoming 4-5 days and as the rice is at panicle initiation to flowering stage so farmers should maintain 5-10cm water level during the critical stage. Plug the holes in rice field and strengthen the field bunds. Farmers should continue irrigation, pesticide and fertilizer application activities and should Start land preparation and sowing of non paddy crops and vegetables like cabbage, cauliflower, radish by utilization of past rainfall associated soil moisture.

**Kharif Rice**

Maximum paddy crops are now mostly in panicle initiation to flowering stage. Moisture Stress during this stage will lead to decrease in yield. As weather to remain mostly dry, ensure proper water level of 4-5 centimeter in the field.

**Weather based Disease and pest of rice**

- Due to slight increase in temperature there may be infestation of **mealy bug** in upland rice. To control this spray Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200litr water for one acre.
- Due to fluctuation in day and night temperature there may be infestation of **hispa /case worm** in rice, to control this spray LambdaCyhalothrin 2.5%EC @ 200 ml/acre or Chlorpyrifos 20EC @ 400 ml/ acre in 200 ltr water.
- Current weather favors **Bacterial leaf streak infestation** in rice, apply Streptomycin sulphate (9%) + Tetracycline hydrochloride (1%) @ 200 g/acre along with Copper oxychloride @ 200g/acre. Use 200 litre solution for one acre area.
- If **Brown Plant Hopper (BPH) infestation** will observe in rice, go for alternate wetting and drying. Make a gape after 8 to 10 rows for entry of sunlight and air. Whenever the number of hoppers/hill reaches 5-10, spray Pymetrozine 50% WG @ 120 g or Buprofezin 25% SC @ 330 ml or Imidacloprid 17.8% SL @ 50 ml per acre in the base of the plant. Before spraying remove water from the field.
- **Chances of gundhi bug incidence for crop at flowering stage. If observe spray Ethofenoprox 10% EC @ 1 ml/litre of water.**

**Maize:**

- Maize is at cob development and harvesting stage. Harvest the matured maize in dry days based upon forecast of rain. There may be Aphids and Thrips infestation, then Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200litr water for one acre. To control stem borer apply fipronil 5SC 400ml/acre in 200 ltr water.

**VEGETABLES:**

- **Now the weather and soil moisture is optimum for land preparation and nursery sowing of cauliflower, cabbage, knolkhol and radish.**
- Fully decomposed FYM or compost should be added before sowing of vegetable. Transplant 35-40 days old seedlings in the main field.
- Undertake planting of **potato crop** on ridge after probable rain; 12 quintal seeds 14 (germinated tubers of 20- 30 g each), 40 kg Urea, 70 kg D.A.P., 80 kg MOP and 10 kg Sulphur are required for cultivation in 1 acre.
- Sow the seeds of **tomato** in nursery bed; farmers who have already raised the seedlings, may transplant them at a distance of 60 cm (row to row) and 40 cm (plant to plant) for HYV and for hybrid variety maintain the distance 60 x 60 cm.
- Monitor for sucking insect pests such as **white fly, leaf hopper and aphid** in vegetables nursery; to control, spray Imidacloprid @ 0.3ml/litre of water. Carry out transplanting of brinjal, tomato and chilli on raised bed.
- To prevent **damping off disease** in cabbage/Cauliflower nursery, treat 1 kg of seeds with 2-3 g Captan 75% WS or 2 g Carbendazim 50% WP before sowing; after sowing, drench the bed soil with 3 g Metalaxyl M 4% + Mancozeb 64% WP mixed in 1 litre of water.
- If **whitefly** will found in vegetative stage then to control spray Imidacloprid @ 0.3ml/litre of water.
- The dry weather favors infestation of **Tobacco caterpillar**, to control this apply 1ml/10 ltr water.

**Animal husbandry:**

Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide dry and clean shelter to the animals. Provide sufficient clean drinking water to the animals.



**DrRanjan Kumar Mohanta**  
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area





**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**



(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-536

Date- 18/10/2024

**AGROMET ADVISORY BULLETIN FOR BADAMBA**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 00 mm and district received average 00 mm rainfall during the last week. **The mean daily maximum and minimum temperature was 33.71°C (+1.03 from normal) and 24.98°C (+2.43 from normal), respectively.** The mean relative humidity was 75-81%.

**Forecasted period: Today(8.30IST) to 23<sup>th</sup> October (8.30IST)2024**

**Weather advisory: Light to moderate rainfall/thundershower is forecasted for today to Sunday and dry weather is forecasted for other 2days in the block.** The day temperature will be 32-34°C and night temperature will be 22-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 77-96% and 48-65% respectively. **There will be partly cloudy to overcast sky for next 5 days.**

Block	Badamba (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	18/10/2024	19/10/2024	20/10/2024	21/10/2024	22/10/2024
Rainfall(mm)	10	18	07	00	00
T-Max(°C)	34	34	32	33	33
T-Min(°C)	25	26	24	24	22
Cloud Cover(octa)	7	8	6	3	4
RH Max (%)	96	94	92	88	77
RH Min (%)	65	64	63	50	48
Wind Speed (Kmph)	6	4	5	6	6
Wind Direction (deg)	159	146	108	68	360

**Agromet Advisory**

**General advisory:** As slight increase in temperature condition is forecasted in between upcoming 4-5 days and as the rice is at panicle initiation to flowering stage so farmers should maintain 5-10cm water level during the critical stage. Plug the holes in rice field and strengthen the field bunds. Farmers should continue irrigation, pesticide and fertilizer application activities and should start land preparation and sowing of non paddy crops and vegetables like cabbage, cauliflower, radish by utilization of past rainfall associated soil moisture.

**Kharif Rice**

Maximum paddy crops are now mostly in panicle initiation to flowering stage. Moisture Stress during this stage will lead to decrease in yield. As weather to remain mostly dry, ensure proper water level of 4-5 centimeter in the field.

**Weather based Disease and pest of rice**

- Due to slight increase in temperature there may be infestation of **mealy bug** in upland rice. To control this spray Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200 liter water for one acre.
- Due to fluctuation in day and night temperature there may be infestation of **hispa /case worm** in rice, to control this spray LambdaCyhalothrin 2.5% EC @ 200 ml/acre or Chlorpyrifos 20EC @ 400 ml/ acre in 200 liter water.
- Current weather favors **Bacterial leaf streak infestation** in rice, apply Streptomycin sulphate (9%) + Tetracycline hydrochloride (1%) @ 200 g/acre along with Copper oxychloride @ 200g/acre. Use 200 litre solution for one acre area.
- If **Brown Plant Hopper (BPH) infestation** will observe in rice, go for alternate wetting and drying. Make a gape after 8 to 10 rows for entry of sunlight and air. Whenever the number of hoppers/hill reaches 5-10, spray Pymetrozine 50% WG @ 120 g or Buprofezin 25% SC @ 330 ml or Imidacloprid 17.8% SL @ 50 ml per acre in the base of the plant. Before spraying remove water from the field.
- **Chances of gundhi bug incidence for crop at flowering stage. If observe spray Ethofenoprox 10% EC @ 1 ml/litre of water.**

**Maize:**

- Maize is at cob development and harvesting stage. Harvest the matured maize in dry days based upon forecast of rain. There may be Aphids and Thrips infestation, then Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200 liter water for one acre. To control stem borer apply fipronil 5SC 400ml/acre in 200 liter water.

**VEGETABLES:**

**Now the weather and soil moisture is optimum for land preparation and nursery sowing of cauliflower, cabbage, knolkhol and radish.**

- Fully decomposed FYM or compost should be added before sowing of vegetable. Transplant 35-40 days old seedlings in the main field.
- Undertake planting of **potato crop** on ridge after probable rain; 12 quintal seeds 14 (germinated tubers of 20- 30 g each), 40 kg Urea, 70 kg D.A.P., 80 kg MOP and 10 kg Sulphur are required for cultivation in 1 acre.
- Sow the seeds of **tomato** in nursery bed; farmers who have already raised the seedlings, may transplant them at a distance of 60 cm (row to row) and 40 cm (plant to plant) for HYV and for hybrid variety maintain the distance 60 x 60 cm.
- Monitor for sucking insect pests such as **white fly, leaf hopper and aphid** in vegetables nursery; to control, spray Imidacloprid @ 0.3ml/litre of water. Carry out transplanting of brinjal, tomato and chilli on raised bed.
- To prevent **damping off disease** in cabbage/Cauliflower nursery, treat 1 kg of seeds with 2-3 g Captan 75% WS or 2 g Carbendazim 50% WP before sowing; after sowing, drench the bed soil with 3 g Metalaxyl M 4% + Mancozeb 64% WP mixed in 1 litre of water.
- If **whitefly** will found in vegetative stage then to control spray Imidacloprid @ 0.3ml/litre of water.
- The dry weather favors infestation of **Tobacco caterpillar**, to control this apply 1ml/10 liter water.

**Animal husbandry:**

Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide dry and clean shelter to the animals. Provide sufficient clean drinking water to the animals.



**DrRanjan Kumar Mohanta**  
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area







**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**



(A Unit of ICAR-National Rice Research Institute)  
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-536  
 Date- 18/10/2024

**AGROMET ADVISORY BULLETIN FOR BANKI**

**District Past week average weather condition and Block past week average Rainfall condition**

The district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 33.71°C (+1.03 from normal) and 24.98°C (+2.43 from normal), respectively. The mean relative humidity was 75-81%.

**Forecasted period: Today(8.30IST) to 23<sup>th</sup> October (8.30IST)2024**

**Weatheradvisory:: Light to moderate rainfall/thundershower is forecasted for today to Sunday and dry weather is forecasted for other 2days in the block.** The day temperature will be 33-34°C and night temperature will be 23-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 78-94% and 48-64% respectively. **There will be partly cloudy to overcast sky for next 5days.**

Block	Banki(VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	18/10/2024	19/10/2024	20/10/2024	21/10/2024	22/10/2024
Rainfall(mm)	10	20	10	00	00
T-Max(°C)	34	34	33	34	34
T-Min(°C)	26	26	24	24	23
Cloud Cover(octa)	7	8	5	2	4
RH Max (%)	94	94	92	87	78
RH Min (%)	64	63	62	51	48
Wind Speed (Kmph)	7	5	6	7	7
Wind Direction (deg)	156	146	23	342	56

**Agromet Advisory**

**General advisory:** As slight increase in temperature condition is forecasted in between upcoming 4-5 days and as the rice is at panicle initiation to flowering stage so farmers should maintain 5-10cm water level during the critical stage. Plug the holes in rice field and strengthen the field bunds. Farmers should continue irrigation, pesticide and fertilizer application activities and should Start land preparation and sowing of non paddy crops and vegetables like cabbage, cauliflower, radish by utilization of past rainfall associated soil moisture.

**Kharif Rice**

Maximum paddy crops are now mostly in panicle initiation to flowering stage. Moisture Stress during this stage will lead to decrease in yield. As weather to remain mostly dry, ensure proper water level of 4-5 centimeter in the field.

**Weather based Disease and pest of rice**

- Due to slight increase in temperature there may be infestation of **mealy bug** in upland rice. To control this spray Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200litr water for one acre.
- Due to fluctuation in day and night temperature there may be infestation of **hispa /case worm** in rice, to control this spray LambdaCyhalothrin 2.5%EC @ 200 ml/acre or Chlorpyrifos 20EC @ 400 ml/ acre in 200 ltr water.
- Current weather favors **Bacterial leaf streak infestation** in rice, apply Streptomycin sulphate (9%) + Tetracycline hydrochloride (1%) @ 200 g/acre along with Copper oxychloride @ 200g/acre. Use 200 litre solution for one acre area.
- If **Brown Plant Hopper (BPH) infestation** will observe in rice, go for alternate wetting and drying. Make a gape after 8 to 10 rows for entry of sunlight and air. Whenever the number of hoppers/hill reaches 5-10, spray Pymetrozine 50% WG @ 120 g or Buprofezin 25% SC @ 330 ml or Imidacloprid 17.8% SL @ 50 ml per acre in the base of the plant. Before spraying remove water from the field.
- **Chances of gundhi bug incidence for crop at flowering stage. If observe spray Ethofenoprox 10% EC @ 1 ml/litre of water.**

**Maize:**

- Maize is at cob development and harvesting stage. Harvest the matured maize in dry days based upon forecast of rain. There may be Aphids and Thrips infestation, then Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200litr water for one acre. To control stem borer apply fipronil 5SC 400ml/acre in 200 ltr water.

**VEGETABLES:**

**Now the weather and soil moisture is optimum for land preparation and nursery sowing of cauliflower, cabbage, knolkhol and radish.**

- Fully decomposed FYM or compost should be added before sowing of vegetable. Transplant 35-40 days old seedlings in the main field.
- Undertake planting of **potato crop** on ridge after probable rain; 12 quintal seeds 14 (germinated tubers of 20- 30 g each), 40 kg Urea, 70 kg D.A.P., 80 kg MOP and 10 kg Sulphur are required for cultivation in 1 acre.
- Sow the seeds of **tomato** in nursery bed; farmers who have already raised the seedlings, may transplant them at a distance of 60 cm (row to row) and 40 cm (plant to plant) for HYV and for hybrid variety maintain the distance 60 x 60 cm.
- Monitor for sucking insect pests such as **white fly, leaf hopper and aphid** in vegetables nursery; to control, spray Imidacloprid @ 0.3ml/litre of water. Carry out transplanting of brinjal, tomato and chilli on raised bed.
- To prevent **damping off disease** in cabbage/Cauliflower nursery, treat 1 kg of seeds with 2-3 g Captan 75% WS or 2 g Carbendazim 50% WP before sowing; after sowing, drench the bed soil with 3 g Metalaxyl M 4% + Mancozeb 64% WP mixed in 1 litre of water.
- If **whitefly** will found in vegetative stage then to control spray Imidacloprid @ 0.3ml/litre of water.
- The dry weather favors infestation of **Tobacco caterpillar**, to control this apply 1ml/10 ltr water.

**Animal husbandry:**

Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide dry and clean shelter to the animals. Provide sufficient clean drinking water to the animals.



**DrRanjan Kumar Mohanta**  
 (Nodal officer)

Follow Meghdoot Mobile APP for Agromet  
 advisories in local language and DAMINI  
 APP for Lightening alert in your area





**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**



(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-536

Date- 18/10/2024

**AGROMET ADVISORY BULLETIN FOR BANKI-DAMPARA**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 33.71°C (+1.03 from normal) and 24.98°C (+2.43 from normal), respectively. The mean relative humidity was 75-81%.

**Forecasted period: Today(8.30IST) to 23<sup>rd</sup> October (8.30IST)2024**

**Weather Advisory:** Light to moderate rainfall/thundershower is forecasted for today to Sunday and dry weather is forecasted for other 2 days in the block. The day temperature will be 33-34°C and night temperature will be 23-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 78-94% and 48-64% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Banki-Dampara (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	18/10/2024	19/10/2024	20/10/2024	21/10/2024	22/10/2024
Rainfall(mm)	10	20	10	00	00
T-Max(°C)	34	34	33	34	34
T-Min(°C)	26	26	24	24	23
Cloud Cover(octa)	7	8	5	2	4
RH Max (%)	94	94	92	86	78
RH Min (%)	64	63	63	51	48
Wind Speed (Kmph)	7	5	6	7	7
Wind Direction (deg)	156	146	23	342	56

**Agromet Advisory**

**General advisory:** As slight increase in temperature condition is forecasted in between upcoming 4-5 days and as the rice is at panicle initiation to flowering stage so farmers should maintain 5-10cm water level during the critical stage. Plug the holes in rice field and strengthen the field bunds. Farmers should continue irrigation, pesticide and fertilizer application activities and should start land preparation and sowing of non paddy crops and vegetables like cabbage, cauliflower, radish by utilization of past rainfall associated soil moisture.

**Kharif Rice**

Maximum paddy crops are now mostly in panicle initiation to flowering stage. Moisture Stress during this stage will lead to decrease in yield. As weather to remain mostly dry, ensure proper water level of 4-5 centimeter in the field.

**Weather based Disease and pest of rice**

- Due to slight increase in temperature there may be infestation of **mealy bug** in upland rice. To control this spray Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200litre water for one acre.
- Due to fluctuation in day and night temperature there may be infestation of **hispa /case worm** in rice, to control this spray LambdaCyhalothrin 2.5%EC @ 200 ml/acre or Chlorpyrifos 20EC @ 400 ml/ acre in 200 ltr water.
- Current weather favors **Bacterial leaf streak infestation** in rice, apply Streptomycin sulphate (9%) + Tetracycline hydrochloride (1%) @ 200 g/acre along with Copper oxychloride @ 200g/acre. Use 200 litre solution for one acre area.
- If **Brown Plant Hopper (BPH) infestation** will observe in rice, go for alternate wetting and drying. Make a gape after 8 to 10 rows for entry of sunlight and air. Whenever the number of hoppers/hill reaches 5-10, spray Pymetrozine 50% WG @ 120 g or Buprofezin 25% SC @ 330 ml or Imidacloprid 17.8% SL @ 50 ml per acre in the base of the plant. Before spraying remove water from the field.
- **Chances of gundhi bug incidence for crop at flowering stage. If observe spray Ethofenoprox 10% EC @ 1 ml/litre of water.**

**Maize:**

- Maize is at cob development and harvesting stage. Harvest the matured maize in dry days based upon forecast of rain. There may be Aphids and Thrips infestation, then Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200litre water for one acre. To control stem borer apply fipronil 5SC 400ml/acre in 200 litre water.

**VEGETABLES:**

- **Now the weather and soil moisture is optimum for land preparation and nursery sowing of cauliflower, cabbage, knolkhol and radish.**
- Fully decomposed FYM or compost should be added before sowing of vegetable. Transplant 35-40 days old seedlings in the main field.
- Undertake planting of **potato crop** on ridge after probable rain; 12 quintal seeds 14 (germinated tubers of 20- 30 g each), 40 kg Urea, 70 kg D.A.P., 80 kg MOP and 10 kg Sulphur are required for cultivation in 1 acre.
- Sow the seeds of **tomato** in nursery bed; farmers who have already raised the seedlings, may transplant them at a distance of 60 cm (row to row) and 40 cm (plant to plant) for HYV and for hybrid variety maintain the distance 60 x 60 cm.
- Monitor for sucking insect pests such as **white fly, leaf hopper and aphid** in vegetables nursery; to control, spray Imidacloprid @ 0.3ml/litre of water. Carry out transplanting of brinjal, tomato and chilli on raised bed.
- To prevent **damping off disease** in cabbage/Cauliflower nursery, treat 1 kg of seeds with 2-3 g Captan 75% WS or 2 g Carbendazim 50% WP before sowing; after sowing, drench the bed soil with 3 g Metalaxyl M 4% + Mancozeb 64% WP mixed in 1 litre of water.
- If **whitefly** will found in vegetative stage then to control spray Imidacloprid @ 0.3ml/litre of water.
- The dry weather favors infestation of **Tobacco caterpillar**, to control this apply 1ml/10 litre water.

**Animal husbandry:**

Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide dry and clean shelter to the animals. Provide sufficient clean drinking water to the animals.



**DrRanjan Kumar Mohanta**  
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area







**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**



(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-536  
Date- 18/10/2024

**AGROMET ADVISORY BULLETIN FOR BARANGA**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 00 mm and district received average 00 mm rainfall during the last week. **The mean daily maximum and minimum temperature was 33.71°C (+1.03 from normal) and 24.98°C (+2.43 from normal), respectively.** The mean relative humidity was 75-81%.

**Forecasted period: Today(8.30IST) to 23<sup>th</sup> October (8.30IST)2024**

**Weather advisory:** *Light to moderate rainfall/thundershower is forecasted for today to Sunday and dry weather is forecasted for other 2 days in the block.* The day temperature will be 32-35°C and night temperature will be 24-27°C for upcoming 5 days. Maximum and minimum relative humidity will be 78-93% and 50-67% respectively. **There will be partly cloudy to overcast sky for next 5 days.**

Block	Baranga (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	18/10/2024	19/10/2024	20/10/2024	21/10/2024	22/10/2024
Rainfall(mm)	09	17	09	00	00
T-Max(°C)	34	35	32	34	33
T-Min(°C)	27	27	25	25	24
Cloud Cover	7	8	6	2	5
RH Max (%)	91	93	91	83	78
RH Min (%)	64	58	67	52	50
Wind Speed (Kmph)	7	5	6	8	9
Wind Direction (deg)	158	146	18	11	23

**Agromet Advisory**

**General advisory:** *As slight increase in temperature condition is forecasted in between upcoming 4-5 days and as the rice is at panicle initiation to flowering stage so farmers should maintain 5-10cm water level during the critical stage. Plug the holes in rice field and strengthen the field bunds. Farmers should continue irrigation, pesticide and fertilizer application activities and should start land preparation and sowing of non paddy crops and vegetables like cabbage, cauliflower, radish by utilization of past rainfall associated soil moisture.*

**Kharif Rice**

Maximum paddy crops are now mostly in panicle initiation to flowering stage. Moisture Stress during this stage will lead to decrease in yield. As weather to remain mostly dry, ensure proper water level of 4-5 centimeter in the field.

**Weather based Disease and pest of rice**

- Due to slight increase in temperature there may be infestation of **mealy bug** in upland rice. To control this spray Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200litre water for one acre.
- Due to fluctuation in day and night temperature there may be infestation of **hispa /case worm** in rice, to control this spray LambdaCyhalothrin 2.5%EC @ 200 ml/acre or Chlorpyrifos 20EC @ 400 ml/ acre in 200 ltr water.
- Current weather favors **Bacterial leaf streak infestation** in rice, apply Streptomycin sulphate (9%) + Tetracycline hydrochloride (1%) @ 200 g/acre along with Copper oxychloride @ 200g/acre. Use 200 litre solution for one acre area.
- If **Brown Plant Hopper (BPH) infestation** will observe in rice, go for alternate wetting and drying. Make a gape after 8 to 10 rows for entry of sunlight and air. Whenever the number of hoppers/hill reaches 5-10, spray Pymetrozine 50% WG @ 120 g or Buprofezin 25% SC @ 330 ml or Imidacloprid 17.8% SL @ 50 ml per acre in the base of the plant. Before spraying remove water from the field.
- **Chances of gundhi bug incidence for crop at flowering stage. If observe spray Ethofenoprox 10% EC @ 1 ml/litre of water.**

**Maize:**

- Maize is at cob development and harvesting stage. Harvest the matured maize in dry days based upon forecast of rain. There may be Aphids and Thrips infestation, then Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200litre water for one acre. To control stem borer apply fipronil 5SC 400ml/acre in 200 litre water.

**VEGETABLES:**

- **Now the weather and soil moisture is optimum for land preparation and nursery sowing of cauliflower, cabbage, knolkhol and radish.**
- Fully decomposed FYM or compost should be added before sowing of vegetable. Transplant 35-40 days old seedlings in the main field.
- Undertake planting of **potato crop** on ridge after probable rain; 12 quintal seeds 14 (germinated tubers of 20- 30 g each), 40 kg Urea, 70 kg D.A.P., 80 kg MOP and 10 kg Sulphur are required for cultivation in 1 acre.
- Sow the seeds of **tomato** in nursery bed; farmers who have already raised the seedlings, may transplant them at a distance of 60 cm (row to row) and 40 cm (plant to plant) for HYV and for hybrid variety maintain the distance 60 x 60 cm.
- Monitor for sucking insect pests such as **white fly, leaf hopper and aphid** in vegetables nursery; to control, spray Imidacloprid @ 0.3ml/litre of water. Carry out transplanting of brinjal, tomato and chilli on raised bed.
- To prevent **damping off disease** in cabbage/Cauliflower nursery, treat 1 kg of seeds with 2-3 g Captan 75% WS or 2 g Carbendazim 50% WP before sowing; after sowing, drench the bed soil with 3 g Metalaxyl M 4% + Mancozeb 64% WP mixed in 1 litre of water.
- If **whitefly** will found in vegetative stage then to control spray Imidacloprid @ 0.3ml/litre of water.
- The dry weather favors infestation of **Tobacco caterpillar**, to control this apply 1ml/10 litre water.

**Animal husbandry:**

Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide dry and clean shelter to the animals. Provide sufficient clean drinking water to the animals.



**DrRanjan Kumar Mohanta**  
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area





**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**



(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-536  
Date- 18/10/2024

**AGROMET ADVISORY BULLETIN FOR CUTTACK SADAR**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 00 mm and district received average 00 mm rainfall during the last week. **The mean daily maximum and minimum temperature was 33.71°C (+1.03 from normal) and 24.98°C (+2.43 from normal), respectively.** The mean relative humidity was 75-81%.

**Forecasted period: Today(8.30IST) to 23<sup>th</sup> October (8.30IST)2024**

**Weather advisory: Light to moderate rainfall/thundershower is forecasted for today to Sunday and dry weather is forecasted for other 2 days in the block.** The day temperature will be 32-35°C and night temperature will be 24-27°C for upcoming 5 days. Maximum and minimum relative humidity will be 78-93% and 48-66% respectively. **There will be partly cloudy to overcast sky for next 5 days.**

Block	Cuttack sadar (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	18/10/2024	19/10/2024	20/10/2024	21/10/2024	22/10/2024
Rainfall(mm)	10	19	08	00	00
T-Max(°C)	34	35	32	34	33
T-Min(°C)	27	27	26	25	24
Cloud Cover(octa)	7	8	6	2	5
RH Max (%)	92	93	90	83	78
RH Min (%)	63	59	66	51	48
Wind Speed (Kmph)	7	5	6	7	8
Wind Direction (deg)	158	153	15	90	45

**Agromet Advisory**

**General advisory:** As slight increase in temperature condition is forecasted in between upcoming 4-5 days and as the rice is at panicle initiation to flowering stage so farmers should maintain 5-10cm water level during the critical stage. Plug the holes in rice field and strengthen the field bunds. Farmers should continue irrigation, pesticide and fertilizer application activities and should start land preparation and sowing of non paddy crops and vegetables like cabbage, cauliflower, radish by utilization of past rainfall associated soil moisture.

**Kharif Rice**

Maximum paddy crops are now mostly in panicle initiation to flowering stage. Moisture Stress during this stage will lead to decrease in yield. As weather to remain mostly dry, ensure proper water level of 4-5 centimeter in the field.

**Weather based Disease and pest of rice**

- Due to slight increase in temperature there may be infestation of **mealy bug** in upland rice. To control this spray Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200litre water for one acre.
- Due to fluctuation in day and night temperature there may be infestation of **hispa /case worm** in rice, to control this spray LambdaCyhalothrin 2.5%EC @ 200 ml/acre or Chlorpyrifos 20EC @ 400 ml/ acre in 200 ltr water.
- Current weather favors **Bacterial leaf streak infestation** in rice, apply Streptomycin sulphate (9%) + Tetracycline hydrochloride (1%) @ 200 g/acre along with Copper oxychloride @ 200g/acre. Use 200 litre solution for one acre area.
- If **Brown Plant Hopper (BPH) infestation** will observe in rice, go for alternate wetting and drying. Make a gape after 8 to 10 rows for entry of sunlight and air. Whenever the number of hoppers/hill reaches 5-10, spray Pymetrozine 50% WG @ 120 g or Buprofezin 25% SC @ 330 ml or Imidacloprid 17.8% SL @ 50 ml per acre in the base of the plant. Before spraying remove water from the field.
- **Chances of gundhi bug incidence for crop at flowering stage. If observe spray Ethofenoprox 10% EC @ 1 ml/litre of water.**

**Sugarcane:**

- Prevailing hot weather favours the infestation of **Mites, Mealy bug, Whitefly and scale insect** in sugarcane. If mites will be found the spray spiromesifen 22.9%SC 1ml/litre of water. To control mealy bug spray Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200litre water for one acre. To control scale insects apply Dimethoate 2ml/litre water or methyl demeton 2ml/litre water. To control white fly go for spraying of Thiamethoxam 40 gram in 200litre water for one acre.

**VEGETABLES:**

**Now the weather and soil moisture is optimum for land preparation and nursery sowing of cauliflower, cabbage, knolkhol and radish.**

- Fully decomposed FYM or compost should be added before sowing of vegetable. Transplant 35-40 days old seedlings in the main field.
- Undertake planting of **potato crop** on ridge after probable rain; 12 quintal seeds 14 (germinated tubers of 20- 30 g each), 40 kg Urea, 70 kg D.A.P., 80 kg MOP and 10 kg Sulphur are required for cultivation in 1 acre.
- Sow the seeds of **tomato** in nursery bed; farmers who have already raised the seedlings, may transplant them at a distance of 60 cm (row to row) and 40 cm (plant to plant) for HYV and for hybrid variety maintain the distance 60 x 60 cm.
- Monitor for sucking insect pests such as **white fly, leaf hopper and aphid** in vegetables nursery; to control, spray Imidacloprid @ 0.3ml/litre of water. Carry out transplanting of brinjal, tomato and chilli on raised bed.
- To prevent **damping off disease** in cabbage/Cauliflower nursery, treat 1 kg of seeds with 2-3 g Captan 75% WS or 2 g Carbendazim 50% WP before sowing; after sowing, drench the bed soil with 3 g Metalaxyl M 4% + Mancozeb 64% WP mixed in 1 litre of water.
- If **whitefly** will found in vegetative stage then to control spray Imidacloprid @ 0.3ml/litre of water.
- The dry weather favors infestation of **Tobacco caterpillar**, to control this apply 1ml/10 litre water.

**Animal husbandry:**

Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide dry and clean shelter to the animals. Provide sufficient clean drinking water to the animals.



**DrRanjan Kumar Mohanta**  
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area





**GRAMIN KRISHI MAUSAM SEWA  
INDIA METEOROLOGICAL DEPARTMENT  
DISTRICT AGROMET UNIT  
KRISHI VIGYAN KENDRA CUTTACK**



(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-536

Date- 18/10/2024

**AGROMET ADVISORY BULLETIN FOR KANTAPADA**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 33.71°C (+1.03 from normal) and 24.98°C (+2.43 from normal), respectively. The mean relative humidity was 75-81%.

**Forecasted period: Today(8.30IST) to 23<sup>th</sup> October (8.30IST)2024**

**Weather advisory: Light to moderate rainfall/thundershower is forecasted for today to Sunday and dry weather is forecasted for other 2days in the block.** The day temperature will be 33-35°C and night temperature will be 23-27°C for upcoming 5 days. Maximum and minimum relative humidity will be 78-93% and 51-66% respectively. **There will be partly cloudy to overcast sky for next 5days.**

Block	Kantapada (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	18/10/2024	19/10/2024	20/10/2024	21/10/2024	22/10/2024
Rainfall(mm)	06	16	09	00	00
T-Max(°C)	34	35	33	34	34
T-Min(°C)	27	27	26	25	23
Cloud Cover(octa)	7	8	7	2	5
RH Max (%)	88	93	90	82	78
RH Min (%)	64	56	66	52	51
Wind Speed (Kmph)	7	6	7	9	10
Wind Direction (deg)	158	124	23	15	23

**Agromet Advisory**

**General advisory: As slight increase in temperature condition is forecasted in between upcoming 4-5 days and as the rice is at panicle initiation to flowering stage so farmers should maintain 5-10cm water level during the critical stage. Plug the holes in rice field and strengthen the field bunds. Farmers should continue irrigation, pesticide and fertilizer application activities and should Start land preparation and sowing of non paddy crops and vegetables like cabbage, cauliflower, radish by utilization of past rainfall associated soil moisture.**

**Kharif Rice**

Maximum paddy crops are now mostly in panicle initiation to flowering stage. Moisture Stress during this stage will lead to decrease in yield. As weather to remain mostly dry, ensure proper water level of 4-5 centimeter in the field.

**Weather based Disease and pest of rice**

- Due to slight increase in temperature there may be infestation of **mealy bug** in upland rice. To control this spray Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200litr water for one acre.
- Due to fluctuation in day and night temperature there may be infestation of **hispa /case worm** in rice, to control this spray LambdaCyhalothrin 2.5%EC @ 200 ml/acre or Chlorpyrifos 20EC @ 400 ml/ acre in 200 ltr water.
- Current weather favors **Bacterial leaf streak infestation** in rice, apply Streptomycin sulphate (9%) + Tetracycline hydrochloride (1%) @ 200 g/acre along with Copper oxychloride @ 200g/acre. Use 200 litre solution for one acre area.
- If **Brown Plant Hopper (BPH) infestation** will observe in rice, go for alternate wetting and drying. Make a gape after 8 to 10 rows for entry of sunlight and air. Whenever the number of hoppers/hill reaches 5-10, spray Pymetrozine 50% WG @ 120 g or Buprofezin 25% SC @ 330 ml or Imidacloprid 17.8% SL @ 50 ml per acre in the base of the plant. Before spraying remove water from the field.
- **Chances of gundhi bug incidence for crop at flowering stage. If observe spray Ethofenoprox 10% EC @ 1 ml/litre of water.**

**Sugarcane:**

- Prevailing hot weather favours the infestation of **Mites, Mealy bug, Whitefly and scale insect** in sugarcane. If mites will be found the spray spiromesifen 22.9%SC 1ml/litr of water. To control mealy bug spray Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200litr water for one acre. To control scale insects apply Dimethoate 2ml/litr water or methyl demeton 2ml/litrwater. To control white fly go for spraying of Thiamethoxam 40 gram in 200litr water for one acre.

**VEGETABLES:**

- **Now the weather and soil moisture is optimum for land preparation and nursery sowing of cauliflower, cabbage, knolkhol and radish.**
- Fully decomposed FYM or compost should be added before sowing of vegetable. Transplant 35-40 days old seedlings in the main field.
- Undertake planting of **potato crop** on ridge after probable rain; 12 quintal seeds 14 (germinated tubers of 20- 30 g each), 40 kg Urea, 70 kg D.A.P., 80 kg MOP and 10 kg Sulphur are required for cultivation in 1 acre.
- Sow the seeds of **tomato** in nursery bed; farmers who have already raised the seedlings, may transplant them at a distance of 60 cm (row to row) and 40 cm (plant to plant) for HYV and for hybrid variety maintain the distance 60 x 60 cm.
- Monitor for sucking insect pests such as **white fly, leaf hopper and aphid** in vegetables nursery; to control, spray Imidacloprid @ 0.3ml/litre of water. Carry out transplanting of brinjal, tomato and chilli on raised bed.
- To prevent **damping off disease** in cabbage/Cauliflower nursery, treat 1 kg of seeds with 2-3 g Captan 75% WS or 2 g Carbendazim 50% WP before sowing; after sowing, drench the bed soil with 3 g Metalaxyl M 4% + Mancozeb 64% WP mixed in 1 litre of water.
- If **whitefly** will found in vegetative stage then to control spray Imidacloprid @ 0.3ml/litre of water.
- The dry weather favors infestation of **Tobacco caterpillar**, to control this apply 1ml/10 liter water.

**Animal husbandry:**

Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide dry and clean shelter to the animals. Provide sufficient clean drinking water to the animals.



**DrRanjan Kumar Mohanta  
(Nodal officer)**

**Follow Meghdoot Mobile APP for Agromet  
advisories in local language and DAMINI  
APP for Lightening alert in your area**







**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**



(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-536  
Date- 18/10/2024

**AGROMET ADVISORY BULLETIN FOR MAHANGA**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 33.71°C (+1.03 from normal) and 24.98°C (+2.43 from normal), respectively. The mean relative humidity was 75-81%.

**Forecasted period: Today(8.30IST) to 23<sup>th</sup> October (8.30IST)2024**

**Weather advisory: Light to moderate rainfall/thundershower is forecasted for today to Sunday and dry weather is forecasted for other 2days in the block.** The day temperature will be and it will be 32-35°C and night temperature will be 24-27°C for upcoming 5 days. Maximum and minimum relative humidity will be 76-93% and 51-69% respectively. **There will be partly cloudy to overcast sky for next 5days.**

Block	Mahanga (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	18/10/2024	19/10/2024	20/10/2024	21/10/2024	22/10/2024
Rainfall(mm)	06	20	08	00	00
T-Max(°C)	34	35	32	34	33
T-Min(°C)	27	27	25	24	24
Cloud Cover(octal)	6	8	7	2	5
RH Max (%)	90	93	90	80	76
RH Min (%)	65	57	69	52	51
Wind Speed (Kmph)	7	5	7	8	10
Wind Direction (deg)	180	158	349	360	23

**Agromet Advisory**

**General advisory: As slight increase in temperature condition is forecasted in between upcoming 4-5 days and as the rice is at panicle initiation to flowering stage so farmers should maintain 5-10cm water level during the critical stage. Plug the holes in rice field and strengthen the field bunds. Farmers should continue irrigation, pesticide and fertilizer application activities and should Start land preparation and sowing of non paddy crops and vegetables like cabbage, cauliflower, radish by utilization of past rainfall associated soil moisture.**

**Kharif Rice**

Maximum paddy crops are now mostly in panicle initiation to flowering stage. Moisture Stress during this stage will lead to decrease in yield. As weather to remain mostly dry, ensure proper water level of 4-5 centimeter in the field.

**Weather based Disease and pest of rice**

- Due to slight increase in temperature there may be infestation of **mealy bug** in upland rice. To control this spray Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200litr water for one acre.
- Due to fluctuation in day and night temperature there may be infestation of **hispa /case worm** in rice, to control this spray LambdaCyhalothrin 2.5%EC @ 200 ml/acre or Chlorpyrifos 20EC @ 400 ml/ acre in 200 ltr water.
- Current weather favors **Bacterial leaf streak infestation** in rice, apply Streptomycin sulphate (9%) + Tetracycline hydrochloride (1%) @ 200 g/acre along with Copper oxychloride @ 200g/acre. Use 200 litre solution for one acre area.
- If **Brown Plant Hopper (BPH) infestation** will observe in rice, go for alternate wetting and drying. Make a gape after 8 to 10 rows for entry of sunlight and air. Whenever the number of hoppers/hill reaches 5-10, spray Pymetrozine 50% WG @ 120 g or Buprofezin 25% SC @ 330 ml or Imidacloprid 17.8% SL @ 50 ml per acre in the base of the plant. Before spraying remove water from the field.
- **Chances of gundhi bug incidence for crop at flowering stage. If observe spray Ethofenoprox 10% EC @ 1 ml/litre of water.**

**Sugarcane:**

- Prevailing hot weather favours the infestation of **Mites, Mealy bug, Whitefly and scale insect** in sugarcane. If mites will be found the spray spiromesifen 22.9%SC 1ml/litr of water. To control mealy bug spray Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200litr water for one acre. To control scale insects apply Dimethoate 2ml/litr water or methyl demeton 2ml/litrwater. To control white fly go for spraying of Thiamethoxam 40 gram in 200litr water for one acre.

**VEGETABLES:**

- **Now the weather and soil moisture is optimum for land preparation and nursery sowing of cauliflower, cabbage, knolkhol and radish.**
- Fully decomposed FYM or compost should be added before sowing of vegetable. Transplant 35-40 days old seedlings in the main field.
- Undertake planting of **potato crop** on ridge after probable rain; 12 quintal seeds 14 (germinated tubers of 20- 30 g each), 40 kg Urea, 70 kg D.A.P., 80 kg MOP and 10 kg Sulphur are required for cultivation in 1 acre.
- Sow the seeds of **tomato** in nursery bed; farmers who have already raised the seedlings, may transplant them at a distance of 60 cm (row to row) and 40 cm (plant to plant) for HYV and for hybrid variety maintain the distance 60 x 60 cm.
- Monitor for sucking insect pests such as **white fly, leaf hopper and aphid** in vegetables nursery; to control, spray Imidacloprid @ 0.3ml/litre of water. Carry out transplanting of brinjal, tomato and chilli on raised bed.
- To prevent **damping off disease** in cabbage/Cauliflower nursery, treat 1 kg of seeds with 2-3 g Captan 75% WS or 2 g Carbendazim 50% WP before sowing; after sowing, drench the bed soil with 3 g Metalaxyl M 4% + Mancozeb 64% WP mixed in 1 litre of water.
- If **whitefly** will found in vegetative stage then to control spray Imidacloprid @ 0.3ml/litre of water.
- The dry weather favors infestation of **Tobacco caterpillar**, to control this apply 1ml/10 liter water.

**Animal husbandry:**

Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide dry and clean shelter to the animals. Provide sufficient clean drinking water to the animals.



**DrRanjan Kumar Mohanta**  
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area





**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**



(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-536  
Date- 18/10/2024

**AGROMET ADVISORY BULLETIN FOR NARASINGHPUR**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 00 mm and district received average 00 mm rainfall during the last week. **The mean daily maximum and minimum temperature was 33.71°C (+1.03 from normal) and 24.98°C (+2.43 from normal), respectively.** The mean relative humidity was 75-81%.

**Forecasted period: Today(8.30IST) to 23<sup>th</sup> October (8.30IST)2024**

**Weather advisory: Light to moderate rainfall/thundershower is forecasted for today to Sunday and dry weather is forecasted for other 2days in the block.** The day temperature will be 31-34°C and night temperature will be 21-25°C for upcoming 5 days. Maximum and minimum relative humidity will be 77-95% and 45-65% respectively. **There will be partly cloudy to overcast sky for next 5days.**

Block	Narasinghpur (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	18/10/2024	19/10/2024	20/10/2024	21/10/2024	22/10/2024
Rainfall(mm)	09	18	08	00	00
T-Max(°C)	33	34	31	33	33
T-Min(°C)	24	25	23	23	21
Cloud Cover(/octa)	6	7	5	2	4
RH Max (%)	97	95	94	90	77
RH Min (%)	65	59	61	46	45
Wind Speed (Kmph)	5	4	4	5	6
Wind Direction (deg)	167	150	90	68	11

**Agromet Advisory**

**General advisory:** As slight increase in temperature condition is forecasted in between upcoming 4-5 days and as the rice is at panicle initiation to flowering stage so farmers should maintain 5-10cm water level during the critical stage. Plug the holes in rice field and strengthen the field bunds. Farmers should continue irrigation, pesticide and fertilizer application activities and should start land preparation and sowing of non paddy crops and vegetables like cabbage, cauliflower, radish by utilization of past rainfall associated soil moisture.

**Kharif Rice**

Maximum paddy crops are now mostly in panicle initiation to flowering stage. Moisture Stress during this stage will lead to decrease in yield. As weather to remain mostly dry, ensure proper water level of 4-5 centimeter in the field.

**Weather based Disease and pest of rice**

- Due to slight increase in temperature there may be infestation of **mealy bug** in upland rice. To control this spray Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200ltr water for one acre.
- Due to fluctuation in day and night temperature there may be infestation of **hispa /case worm** in rice, to control this spray LambdaCyhalothrin 2.5%EC @ 200 ml/acre or Chlorpyrifos 20EC @ 400 ml/ acre in 200 ltr water.
- Current weather favors **Bacterial leaf streak infestation** in rice, apply Streptomycin sulphate (9%) + Tetracycline hydrochloride (1%) @ 200 g/acre along with Copper oxychloride @ 200g/acre. Use 200 litre solution for one acre area.
- If **Brown Plant Hopper (BPH) infestation** will observe in rice, go for alternate wetting and drying. Make a gape after 8 to 10 rows for entry of sunlight and air. Whenever the number of hoppers/hill reaches 5-10, spray Pymetrozine 50% WG @ 120 g or Buprofezin 25% SC @ 330 ml or Imidacloprid 17.8% SL @ 50 ml per acre in the base of the plant. Before spraying remove water from the field.
- **Chances of gundhi bug incidence for crop at flowering stage. If observe spray Ethofenoprox 10% EC @ 1 ml/litre of water.**

**Maize:**

- Maize is at cob development and harvesting stage. Harvest the matured maize in dry days based upon forecast of rain. There may be Aphids and Thrips infestation, then Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200ltr water for one acre. To control stem borer apply fipronil 5SC 400ml/acre in 200 litre water.

**VEGETABLES:**

- **Now the weather and soil moisture is optimum for land preparation and nursery sowing of cauliflower, cabbage, knolkhol and radish.**
- Fully decomposed FYM or compost should be added before sowing of vegetable. Transplant 35-40 days old seedlings in the main field.
- Undertake planting of **potato crop** on ridge after probable rain; 12 quintal seeds 14 (germinated tubers of 20- 30 g each), 40 kg Urea, 70 kg D.A.P., 80 kg MOP and 10 kg Sulphur are required for cultivation in 1 acre.
- Sow the seeds of **tomato** in nursery bed; farmers who have already raised the seedlings, may transplant them at a distance of 60 cm (row to row) and 40 cm (plant to plant) for HYV and for hybrid variety maintain the distance 60 x 60 cm.
- Monitor for sucking insect pests such as **white fly, leaf hopper and aphid** in vegetables nursery; to control, spray Imidacloprid @ 0.3ml/litre of water. Carry out transplanting of brinjal, tomato and chilli on raised bed.
- To prevent **damping off disease** in cabbage/Cauliflower nursery, treat 1 kg of seeds with 2-3 g Captan 75% WS or 2 g Carbendazim 50% WP before sowing; after sowing, drench the bed soil with 3 g Metalaxyl M 4% + Mancozeb 64% WP mixed in 1 litre of water.
- If **whitefly** will found in vegetative stage then to control spray Imidacloprid @ 0.3ml/litre of water.
- The dry weather favors infestation of **Tobacco caterpillar**, to control this apply 1ml/10 litre water.

**Animal husbandry:**

Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide dry and clean shelter to the animals. Provide sufficient clean drinking water to the animals.



**DrRanjan Kumar Mohanta**  
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area





**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**



(A Unit of ICAR-National Rice Research Institute)  
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-536  
 Date- 18/10/2024

**AGROMET ADVISORY BULLETIN FOR NIALI**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 00 mm and district received average 00 mm rainfall during the last week. **The mean daily maximum and minimum temperature was 33.71°C (+1.03 from normal) and 24.98°C (+2.43 from normal), respectively.** The mean relative humidity was 75-81%.

**Forecasted period: Today(8.30IST) to 23<sup>th</sup> October (8.30IST)2024**

**Weather advisory: Light to moderate rainfall/thundershower is forecasted for today to Sunday and dry weather is forecasted for other 2days in the block.** The day temperature will be 33-35°C and night temperature will be 23-27°C for upcoming 5 days. Maximum and minimum relative humidity will be 77-93% and 51-65% respectively. **There will be partly cloudy to overcast sky for next 5days.**

Block	Niali (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	18/10/2024	19/10/2024	20/10/2024	21/10/2024	22/10/2024
Rainfall(mm)	06	17	08	00	00
T-Max(°C)	34	35	33	35	34
T-Min(°C)	27	27	26	24	23
Cloud Cover(octa)	7	8	7	3	5
RH Max (%)	85	93	91	81	77
RH Min (%)	65	57	65	52	51
Wind Speed(Kmph)	7	7	7	10	11
Wind Direction(deg)	162	154	45	23	23

**Agromet Advisory**

**General advisory:** As slight increase in temperature condition is forecasted in between upcoming 4-5 days and as the rice is at panicle initiation to flowering stage so farmers should maintain 5-10cm water level during the critical stage. Plug the holes in rice field and strengthen the field bunds. Farmers should continue irrigation, pesticide and fertilizer application activities and should start land preparation and sowing of non paddy crops and vegetables like cabbage, cauliflower, radish by utilization of past rainfall associated soil moisture.

**Kharif Rice**

Maximum paddy crops are now mostly in panicle initiation to flowering stage. Moisture Stress during this stage will lead to decrease in yield. As weather to remain mostly dry, ensure proper water level of 4-5 centimeter in the field.

**Weather based Disease and pest of rice**

- Due to slight increase in temperature there may be infestation of **mealy bug** in upland rice. To control this spray Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200litre water for one acre.
- Due to fluctuation in day and night temperature there may be infestation of **hispa /case worm** in rice, to control this spray LambdaCyhalothrin 2.5%EC @ 200 ml/acre or Chlorpyrifos 20EC @ 400 ml/ acre in 200 ltr water.
- Current weather favors **Bacterial leaf streak infestation** in rice, apply Streptomycin sulphate (9%) + Tetracycline hydrochloride (1%) @ 200 g/acre along with Copper oxychloride @ 200g/acre. Use 200 litre solution for one acre area.
- If **Brown Plant Hopper (BPH) infestation** will observe in rice, go for alternate wetting and drying. Make a gape after 8 to 10 rows for entry of sunlight and air. Whenever the number of hoppers/hill reaches 5-10, spray Pymetrozine 50% WG @ 120 g or Buprofezin 25% SC @ 330 ml or Imidacloprid 17.8% SL @ 50 ml per acre in the base of the plant. Before spraying remove water from the field.
- **Chances of gundhi bug incidence for crop at flowering stage. If observe spray Ethofenoprox 10% EC @ 1 ml/litre of water.**

**Sugarcane:**

- Prevailing hot weather favours the infestation of **Mites, Mealy bug, Whitefly and scale insect** in sugarcane. If mites will be found the spray spiromesifen 22.9%SC 1ml/litre of water. To control mealy bug spray Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200litre water for one acre. To control scale insects apply Dimethoate 2ml/litre water or methyl demeton 2ml/litre water. To control white fly go for spraying of Thiamethoxam 40 gram in 200litre water for one acre.

**VEGETABLES:**

**Now the weather and soil moisture is optimum for land preparation and nursery sowing of cauliflower, cabbage, knolkhol and radish.**

- Fully decomposed FYM or compost should be added before sowing of vegetable. Transplant 35-40 days old seedlings in the main field.
- Undertake planting of **potato crop** on ridge after probable rain; 12 quintal seeds 14 (germinated tubers of 20- 30 g each), 40 kg Urea, 70 kg D.A.P., 80 kg MOP and 10 kg Sulphur are required for cultivation in 1 acre.
- Sow the seeds of **tomato** in nursery bed; farmers who have already raised the seedlings, may transplant them at a distance of 60 cm (row to row) and 40 cm (plant to plant) for HYV and for hybrid variety maintain the distance 60 x 60 cm.
- Monitor for sucking insect pests such as **white fly, leaf hopper and aphid** in vegetables nursery; to control, spray Imidacloprid @ 0.3ml/litre of water. Carry out transplanting of brinjal, tomato and chilli on raised bed.
- To prevent **damping off disease** in cabbage/Cauliflower nursery, treat 1 kg of seeds with 2-3 g Captan 75% WS or 2 g Carbendazim 50% WP before sowing; after sowing, drench the bed soil with 3 g Metalaxyl M 4% + Mancozeb 64% WP mixed in 1 litre of water.
- If **whitefly** will found in vegetative stage then to control spray Imidacloprid @ 0.3ml/litre of water.
- The dry weather favors infestation of **Tobacco caterpillar**, to control this apply 1ml/10 litre water.

**Animal husbandry:**

Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide dry and clean shelter to the animals. Provide sufficient clean drinking water to the animals.



**DrRanjan Kumar Mohanta**  
 (Nodal officer)

Follow Meghdoot Mobile APP for Agromet  
 advisories in local language and DAMINI  
 APP for Lightening alert in your area







**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**



(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-536  
Date- 18/10/2024

**AGROMET ADVISORY BULLETIN FOR NISCHINTAKOILI**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 00 mm and district received average 00 mm rainfall during the last week. **The mean daily maximum and minimum temperature was 33.71°C (+1.03 from normal) and 24.98°C (+2.43 from normal), respectively.** The mean relative humidity was 75-81%.

**Forecasted period: Today(8.30IST) to 23<sup>th</sup> October (8.30IST)2024**

**Weather advisory: Light to moderate rainfall/thundershower is forecasted for today to Sunday and dry weather is forecasted for other 2 days in the block.** The day temperature will be 32-35°C and night temperature will be 24-27°C for upcoming 5 days. Maximum and minimum relative humidity will be 76-93% and 51-68% respectively. **There will be partly cloudy to overcast sky for next 5 days.**

Block	Nischintakoili (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	18/10/2024	19/10/2024	20/10/2024	21/10/2024	22/10/2024
Rainfall(mm)	06	20	07	00	00
T-Max(°C)	34	35	32	34	34
T-Min(°C)	27	27	26	25	24
Cloud Cover(octa)	6	8	7	2	5
RH Max (%)	89	93	90	80	76
RH Min (%)	65	57	68	52	51
Wind Speed (Kmph)	7	5	7	9	10
Wind Direction (deg)	180	158	349	360	23

**Agromet Advisory**

**General advisory: As slight increase in temperature condition is forecasted in between upcoming 4-5 days and as the rice is at panicle initiation to flowering stage so farmers should maintain 5-10cm water level during the critical stage. Plug the holes in rice field and strengthen the field bunds. Farmers should continue irrigation, pesticide and fertilizer application activities and should start land preparation and sowing of non paddy crops and vegetables like cabbage, cauliflower, radish by utilization of past rainfall associated soil moisture.**

**Kharif Rice**

Maximum paddy crops are now mostly in panicle initiation to flowering stage. Moisture Stress during this stage will lead to decrease in yield. As weather to remain mostly dry, ensure proper water level of 4-5 centimeter in the field.

**Weather based Disease and pest of rice**

- Due to slight increase in temperature there may be infestation of **mealy bug** in upland rice. To control this spray Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200litre water for one acre.
- Due to fluctuation in day and night temperature there may be infestation of **hispa /case worm** in rice, to control this spray LambdaCyhalothrin 2.5%EC @ 200 ml/acre or Chlorpyrifos 20EC @ 400 ml/ acre in 200 ltr water.
- Current weather favors **Bacterial leaf streak infestation** in rice, apply Streptomycin sulphate (9%) + Tetracycline hydrochloride (1%) @ 200 g/acre along with Copper oxychloride @ 200g/acre. Use 200 litre solution for one acre area.
- If **Brown Plant Hopper (BPH) infestation** will observe in rice, go for alternate wetting and drying. Make a gape after 8 to 10 rows for entry of sunlight and air. Whenever the number of hoppers/hill reaches 5-10, spray Pymetrozine 50% WG @ 120 g or Buprofezin 25% SC @ 330 ml or Imidacloprid 17.8% SL @ 50 ml per acre in the base of the plant. Before spraying remove water from the field.
- **Chances of gundhi bug incidence for crop at flowering stage. If observe spray Ethofenoprox 10% EC @ 1 ml/litre of water.**

**Sugarcane:**

- Prevailing hot weather favours the infestation of **Mites, Mealy bug, Whitefly and scale insect** in sugarcane. If mites will be found the spray spiromesifen 22.9%SC 1ml/litre of water. To control mealy bug spray Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200litre water for one acre. To control scale insects apply Dimethoate 2ml/litre water or methyl demeton 2ml/litre water. To control white fly go for spraying of Thiamethoxam 40 gram in 200litre water for one acre.

**VEGETABLES:**

**Now the weather and soil moisture is optimum for land preparation and nursery sowing of cauliflower, cabbage, knolkhol and radish.**

- Fully decomposed FYM or compost should be added before sowing of vegetable. Transplant 35-40 days old seedlings in the main field.
- Undertake planting of **potato crop** on ridge after probable rain; 12 quintal seeds 14 (germinated tubers of 20- 30 g each), 40 kg Urea, 70 kg D.A.P., 80 kg MOP and 10 kg Sulphur are required for cultivation in 1 acre.
- Sow the seeds of **tomato** in nursery bed; farmers who have already raised the seedlings, may transplant them at a distance of 60 cm (row to row) and 40 cm (plant to plant) for HYV and for hybrid variety maintain the distance 60 x 60 cm.
- Monitor for sucking insect pests such as **white fly, leaf hopper and aphid** in vegetables nursery; to control, spray Imidacloprid @ 0.3ml/litre of water. Carry out transplanting of brinjal, tomato and chilli on raised bed.
- To prevent **damping off disease** in cabbage/Cauliflower nursery, treat 1 kg of seeds with 2-3 g Captan 75% WS or 2 g Carbendazim 50% WP before sowing; after sowing, drench the bed soil with 3 g Metalaxyl M 4% + Mancozeb 64% WP mixed in 1 litre of water.
- If **whitefly** will found in vegetative stage then to control spray Imidacloprid @ 0.3ml/litre of water.
- The dry weather favors infestation of **Tobacco caterpillar**, to control this apply 1ml/10 litre water.

**Animal husbandry:**

Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide dry and clean shelter to the animals. Provide sufficient clean drinking water to the animals.



**DrRanjan Kumar Mohanta**  
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area





**GRAMIN KRISHI MAUSAM SEWA  
INDIA METEOROLOGICAL DEPARTMENT  
DISTRICT AGROMET UNIT  
KRISHI VIGYAN KENDRA CUTTACK**



(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-536

Date- 18/10/2024

**AGROMET ADVISORY BULLETIN FOR SALEPUR**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 00 mm and district received average 00 mm rainfall during the last week. **The mean daily maximum and minimum temperature was 33.71°C (+1.03 from normal) and 24.98°C (+2.43 from normal), respectively.** The mean relative humidity was 75-81%.

**Forecasted period: Today(8.30IST) to 23<sup>th</sup> October (8.30IST)2024**

**Weather advisory: Light to moderate rainfall/thundershower is forecasted for today to Sunday and dry weather is forecasted for other 2days in the block.** The day temperature will be 32-35°C and night temperature will be 24-27°C for upcoming 5 days. Maximum and minimum relative humidity will be 76-93% and 51-69% respectively. **There will be partly cloudy to overcast sky for next 5days.**

Block	Salepur (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	18/10/2024	19/10/2024	20/10/2024	21/10/2024	22/10/2024
Rainfall(mm)	06	20	07	00	00
T-Max(°C)	34	35	32	34	34
T-Min(°C)	27	27	26	25	24
Cloud Cover(octa)	6	8	6	2	5
RH Max (%)	90	93	90	80	76
RH Min (%)	65	57	69	52	51
Wind Speed (Kmph)	7	5	7	8	10
Wind Direction (deg)	193	158	349	360	23

**Agromet Advisory**

**General advisory: As slight increase in temperature condition is forecasted in between upcoming 4-5 days and as the rice is at panicle initiation to flowering stage so farmers should maintain 5-10cm water level during the critical stage. Plug the holes in rice field and strengthen the field bunds. Farmers should continue irrigation, pesticide and fertilizer application activities and should Start land preparation and sowing of non paddy crops and vegetables like cabbage, cauliflower, radish by utilization of past rainfall associated soil moisture.**

**Kharif Rice**

Maximum paddy crops are now mostly in panicle initiation to flowering stage. Moisture Stress during this stage will lead to decrease in yield. As weather to remain mostly dry, ensure proper water level of 4-5 centimeter in the field.

**Weather based Disease and pest of rice**

- Due to slight increase in temperature there may be infestation of **mealy bug** in upland rice. To control this spray Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200litr water for one acre.
- Due to fluctuation in day and night temperature there may be infestation of **hispa /case worm** in rice, to control this spray LambdaCyhalothrin 2.5%EC @ 200 ml/acre or Chlorpyrifos 20EC @ 400 ml/ acre in 200 ltr water.
- Current weather favors **Bacterial leaf streak infestation** in rice, apply Streptomycin sulphate (9%) + Tetracycline hydrochloride (1%) @ 200 g/acre along with Copper oxychloride @ 200g/acre. Use 200 litre solution for one acre area.
- If **Brown Plant Hopper (BPH) infestation** will observe in rice, go for alternate wetting and drying. Make a gape after 8 to 10 rows for entry of sunlight and air. Whenever the number of hoppers/hill reaches 5-10, spray Pymetrozine 50% WG @ 120 g or Buprofezin 25% SC @ 330 ml or Imidacloprid 17.8% SL @ 50 ml per acre in the base of the plant. Before spraying remove water from the field.
- **Chances of gundhi bug incidence for crop at flowering stage. If observe spray Ethofenoprox 10% EC @ 1 ml/litre of water.**

**Sugarcane:**

- Prevailing hot weather favours the infestation of Mites, Mealy bug, Whitefly and scale insect in sugarcane. If mites will be found the spray spiromesifen 22.9%SC 1ml/litr of water. To control mealy bug spray Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200litr water for one acre. To control scale insects apply Dimethoate 2ml/litr water or methyl demeton 2ml/litrwater. To control white fly go for spraying of Thiamethoxam 40 gram in 200litr water for one acre.

**VEGETABLES:**

**Now the weather and soil moisture is optimum for land preparation and nursery sowing of cauliflower, cabbage, knolkhol and radish.**

- Fully decomposed FYM or compost should be added before sowing of vegetable. Transplant 35-40 days old seedlings in the main field.
- Undertake planting of **potato crop** on ridge after probable rain; 12 quintal seeds 14 (germinated tubers of 20- 30 g each), 40 kg Urea, 70 kg D.A.P., 80 kg MOP and 10 kg Sulphur are required for cultivation in 1 acre.
- Sow the seeds of **tomato** in nursery bed; farmers who have already raised the seedlings, may transplant them at a distance of 60 cm (row to row) and 40 cm (plant to plant) for HYV and for hybrid variety maintain the distance 60 x 60 cm.
- Monitor for sucking insect pests such as **white fly, leaf hopper and aphid** in vegetables nursery; to control, spray Imidacloprid @ 0.3ml/litre of water. Carry out transplanting of brinjal, tomato and chilli on raised bed.
- To prevent **damping off disease** in cabbage/Cauliflower nursery, treat 1 kg of seeds with 2-3 g Captan 75% WS or 2 g Carbendazim 50% WP before sowing; after sowing, drench the bed soil with 3 g Metalaxyl M 4% + Mancozeb 64% WP mixed in 1 litre of water.
- If **whitefly** will found in vegetative stage then to control spray Imidacloprid @ 0.3ml/litre of water.
- The dry weather favors infestation of **Tobacco caterpillar**, to control this apply 1ml/10 liter water.

**Animal husbandry:**

Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide dry and clean shelter to the animals. Provide sufficient clean drinking water to the animals.



**DrRanjan Kumar Mohanta  
(Nodal officer)**

**Follow Meghdoot Mobile APP for Agromet  
advisories in local language and DAMINI  
APP for Lightening alert in your area**





**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**



(A Unit of ICAR-National Rice Research Institute)

Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-536  
Date- 18/10/2024

**AGROMET ADVISORY BULLETIN FOR TANGI-CHOUDWAR**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 33.71°C (+1.03 from normal) and 24.98°C (+2.43 from normal), respectively. The mean relative humidity was 75-81%.

**Forecasted period: Today(8.30IST) to 23<sup>th</sup> October (8.30IST)2024**

**Weather advisory: Light to moderate rainfall/thundershower is forecasted for today to Sunday and dry weather is forecasted for other 2 days in the block.** The day temperature will be 32-33°C and night temperature will be 25-27°C for upcoming 5 days. Maximum and minimum relative humidity will be 71-92% and 53-67% respectively. **There will be partly cloudy to overcast sky for next 5 days.**

Block	Tangi-Choudwar (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	18/10/2024	19/10/2024	20/10/2024	21/10/2024	22/10/2024
Rainfall(mm)	06	15	07	00	00
T-Max(°C)	32	33	32	33	32
T-Min(°C)	27	27	26	25	25
Cloud Cover(octa)	6	7	7	3	5
RH Max (%)	83	92	88	82	71
RH Min (%)	67	60	63	57	53
Wind Speed (Kmph)	8	8	9	11	15
Wind Direction (deg)	180	169	45	60	26

**Agromet Advisory**

**General advisory: As slight increase in temperature condition is forecasted in between upcoming 4-5 days and as the rice is at panicle initiation to flowering stage so farmers should maintain 5-10cm water level during the critical stage. Plug the holes in rice field and strengthen the field bunds. Farmers should continue irrigation, pesticide and fertilizer application activities and should start land preparation and sowing of non paddy crops and vegetables like cabbage, cauliflower, radish by utilization of past rainfall associated soil moisture.**

**Kharif Rice**

Maximum paddy crops are now mostly in panicle initiation to flowering stage. Moisture Stress during this stage will lead to decrease in yield. As weather to remain mostly dry, ensure proper water level of 4-5 centimeter in the field.

**Weather based Disease and pest of rice**

- Due to slight increase in temperature there may be infestation of **mealy bug** in upland rice. To control this spray Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200litre water for one acre.
- Due to fluctuation in day and night temperature there may be infestation of **hispa /case worm** in rice, to control this spray LambdaCyhalothrin 2.5%EC @ 200 ml/acre or Chlorpyrifos 20EC @ 400 ml/ acre in 200 ltr water.
- Current weather favors **Bacterial leaf streak infestation** in rice, apply Streptomycin sulphate (9%) + Tetracycline hydrochloride (1%) @ 200 g/acre along with Copper oxychloride @ 200g/acre. Use 200 litre solution for one acre area.
- If **Brown Plant Hopper (BPH) infestation** will observe in rice, go for alternate wetting and drying. Make a gape after 8 to 10 rows for entry of sunlight and air. Whenever the number of hoppers/hill reaches 5-10, spray Pymetrozine 50% WG @ 120 g or Buprofezin 25% SC @ 330 ml or Imidacloprid 17.8% SL @ 50 ml per acre in the base of the plant. Before spraying remove water from the field.
- **Chances of gundhi bug incidence for crop at flowering stage. If observe spray Ethofenoprox 10% EC @ 1 ml/litre of water.**

**Maize:**

- Maize is at cob development and harvesting stage. Harvest the matured maize in dry days based upon forecast of rain. There may be Aphids and Thrips infestation, then Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200litre water for one acre. To control stem borer apply fipronil 5SC 400ml/acre in 200 litre water.

**VEGETABLES:**

- **Now the weather and soil moisture is optimum for land preparation and nursery sowing of cauliflower, cabbage, knolkhol and radish.**
- Fully decomposed FYM or compost should be added before sowing of vegetable. Transplant 35-40 days old seedlings in the main field.
- Undertake planting of **potato crop** on ridge after probable rain; 12 quintal seeds 14 (germinated tubers of 20- 30 g each), 40 kg Urea, 70 kg D.A.P., 80 kg MOP and 10 kg Sulphur are required for cultivation in 1 acre.
- Sow the seeds of **tomato** in nursery bed; farmers who have already raised the seedlings, may transplant them at a distance of 60 cm (row to row) and 40 cm (plant to plant) for HYV and for hybrid variety maintain the distance 60 x 60 cm.
- Monitor for sucking insect pests such as **white fly, leaf hopper and aphid** in vegetables nursery; to control, spray Imidacloprid @ 0.3ml/litre of water. Carry out transplanting of brinjal, tomato and chilli on raised bed.
- To prevent **damping off disease** in cabbage/Cauliflower nursery, treat 1 kg of seeds with 2-3 g Captan 75% WS or 2 g Carbendazim 50% WP before sowing; after sowing, drench the bed soil with 3 g Metalaxyl M 4% + Mancozeb 64% WP mixed in 1 litre of water.
- If **whitefly** will found in vegetative stage then to control spray Imidacloprid @ 0.3ml/litre of water.
- The dry weather favors infestation of **Tobacco caterpillar**, to control this apply 1ml/10 litre water.

**Animal husbandry:**

Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide dry and clean shelter to the animals. Provide sufficient clean drinking water to the animals.



**DrRanjan Kumar Mohanta**  
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area







**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**



(A Unit of ICAR-National Rice Research Institute)  
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-536  
 Date- 18/10/2024

**AGROMET ADVISORY BULLETIN FOR TIGIRIA**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 00 mm and district received average 00 mm rainfall during the last week. The mean daily maximum and minimum temperature was 33.71°C (+1.03 from normal) and 24.98°C (+2.43 from normal), respectively. The mean relative humidity was 75-81%.

**Forecasted period: Today(8.30IST) to 23<sup>th</sup> October (8.30IST)2024**

**Weatheradvisory: Light to moderate rainfall/thundershower is forecasted for today to Sunday and dry weather is forecasted for other 2 days in the block.** The day temperature will be 32-34°C and night temperature will be 23-26°C for upcoming 5 days. Maximum and minimum relative humidity will be 78-96% and 48-64% respectively. **There will be partly cloudy to overcast sky for next 5 days.**

Block	Tigiria (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	18/10/2024	19/10/2024	20/10/2024	21/10/2024	22/10/2024
Rainfall(mm)	09	20	08	00	00
T-Max(°C)	34	34	32	34	34
T-Min(°C)	26	26	24	24	23
Cloud Cover(octa)	7	8	5	2	4
RH Max (%)	96	94	93	88	78
RH Min (%)	64	64	64	51	48
Wind Speed (Kmph)	7	5	5	6	7
Wind Direction (deg)	153	135	108	105	56

**Agro met Advisory**

**General advisory:** As slight increase in temperature condition is forecasted in between upcoming 4-5 days and as the rice is at panicle initiation to flowering stage so farmers should maintain 5-10cm water level during the critical stage. Plug the holes in rice field and strengthen the field bunds. Farmers should continue irrigation, pesticide and fertilizer application activities and should start land preparation and sowing of non paddy crops and vegetables like cabbage, cauliflower, radish by utilization of past rainfall associated soil moisture.

**Kharif Rice**

Maximum paddy crops are now mostly in panicle initiation to flowering stage. Moisture Stress during this stage will lead to decrease in yield. As weather to remain mostly dry, ensure proper water level of 4-5 centimeter in the field.

**Weather based Disease and pest of rice**

- Due to slight increase in temperature there may be infestation of **mealy bug** in upland rice. To control this spray Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200litr water for one acre.
- Due to fluctuation in day and night temperature there may be infestation of **hispa /case worm** in rice, to control this spray LambdaCyhalothrin 2.5%EC @ 200 ml/acre or Chlorpyrifos 20EC @ 400 ml/ acre in 200 ltr water.
- Current weather favors **Bacterial leaf streak infestation** in rice, apply Streptomycin sulphate (9%) + Tetracycline hydrochloride (1%) @ 200 g/acre along with Copper oxychloride @ 200g/acre. Use 200 litre solution for one acre area.
- If **Brown Plant Hopper (BPH) infestation** will observe in rice, go for alternate wetting and drying. Make a gape after 8 to 10 rows for entry of sunlight and air. Whenever the number of hoppers/hill reaches 5-10, spray Pymetrozine 50% WG @ 120 g or Buprofezin 25% SC @ 330 ml or Imidacloprid 17.8% SL @ 50 ml per acre in the base of the plant. Before spraying remove water from the field.
- **Chances of gundhi bug incidence for crop at flowering stage. If observe spray Ethofenoprox 10% EC @ 1 ml/litre of water.**

**Maize:**

- Maize is at cob development and harvesting stage. Harvest the matured maize in dry days based upon forecast of rain. There may be Aphids and Thrips infestation, then Imidacloprid 17.8 SL 50 ml or Thiamethoxam 40 gram in 200litr water for one acre. To control stem borer apply fipronil 5SC 400ml/acre in 200 ltr water.

**VEGETABLES:**

**Now the weather and soil moisture is optimum for land preparation and nursery sowing of cauliflower, cabbage, knolkhol and radish.**

- Fully decomposed FYM or compost should be added before sowing of vegetable. Transplant 35-40 days old seedlings in the main field.
- Undertake planting of **potato crop** on ridge after probable rain; 12 quintal seeds 14 (germinated tubers of 20- 30 g each), 40 kg Urea, 70 kg D.A.P., 80 kg MOP and 10 kg Sulphur are required for cultivation in 1 acre.
- Sow the seeds of **tomato** in nursery bed; farmers who have already raised the seedlings, may transplant them at a distance of 60 cm (row to row) and 40 cm (plant to plant) for HYV and for hybrid variety maintain the distance 60 x 60 cm.
- Monitor for sucking insect pests such as **white fly, leaf hopper and aphid** in vegetables nursery; to control, spray Imidacloprid @ 0.3ml/litre of water. Carry out transplanting of brinjal, tomato and chilli on raised bed.
- To prevent **damping off disease** in cabbage/Cauliflower nursery, treat 1 kg of seeds with 2-3 g Captan 75% WS or 2 g Carbendazim 50% WP before sowing; after sowing, drench the bed soil with 3 g Metalaxyl M 4% + Mancozeb 64% WP mixed in 1 litre of water.
- If **whitefly** will found in vegetative stage then to control spray Imidacloprid @ 0.3ml/litre of water.
- The dry weather favors infestation of **Tobacco caterpillar**, to control this apply 1ml/10 ltr water.

**Animal husbandry:**

Vaccinate calves above 3 months and sheep and goat above two months against Foot and Mouth disease. Provide dry and clean shelter to the animals. Provide sufficient clean drinking water to the animals.



**DrRanjan Kumar Mohanta**  
 (Nodal officer)

Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area







**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**  
 (A Unit of ICAR-National Rice Research Institute)  
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



**Bulletin No.-536**  
**Date- 18/10/2024**

				
<b>MEGHDOOT APP</b>	<b>DAMINI APP</b>	<b>FACEBOOK (Damu Cuttack)</b>	<b>BLOCK SPECIFIC WHAT'S APP GROUPS</b>	<b>@DAMUCUTTACK</b>

- Kindly install and use **“Meghdoot mobile App”** for Cuttack district weather forecast, warnings and weather based agromet advisory services.  
<https://play.google.com/store/apps/details?id=com.aas.meghdoot>
- Kindly install and use **“Damini mobile App”** for location specific for lightening forecast and warnings.  
<https://play.google.com/store/apps/details?id=com.lightening.live.damini>
- Follow the **block specific whats app groups** and join in your block wise groups for getting weather based agromet advisory services twice in a week (Tuesday and Friday) and other weather warning based farm management advisories.
- Follow the **Official Website links** to get the weather based agromet advisory service bulletins (**English & Odia**) twice in a week (Tuesday and Friday).

1	<b>ICAR-NRRI, Cuttack, Odisha</b>	<a href="https://icar-nrri.in/agromate/">https://icar-nrri.in/agromate/</a>	
2	<b>Agrimet division, IMD, Pune</b>	<a href="https://imdagrmet.gov.in/imdproject/AGIndex.php">https://imdagrmet.gov.in/imdproject/AGIndex.php</a>	



**DrRanjan Kumar Mohanta**  
 (Nodal officer)

**Follow Meghdoot Mobile APP for Agromet  
advisories in local language and DAMINI  
APP for Lightening alert in your area**

