




# *Cuttack District Block Wise Agro-met Advisory Service Bulletin*

*Tuesday, 24<sup>th</sup> December, 2024  
 (For the period 24<sup>th</sup> to 28<sup>th</sup> December,  
 2024)*





*Issued by  
 District Agromet Unit,  
 KrishiVigyan Kendra Cuttack*





**STANDARD PRECIPITATION INDEX AND NDVI MAP**

Standard Precipitation Index (SPI)	Normalized difference vegetation Index (NDVI)
<p style="text-align: center;">MINISTRY OF EARTH SCIENCES INDIA METEOROLOGICAL DEPARTMENT HYDROMET SECTION, CRS PUNE</p>  <p style="font-size: small;">STANDARDISED PRECIPITATION INDEX(SPI) FOR THE PERIOD 14-11-2024 to 11-12-2024</p> <p style="font-size: x-small;">SPI Values        ○ No Data        ● Extremely Dry (-1.4 to -2) 0.0% Dist 0.0% Area        ● Severely Dry (-1.39 to -1.5) 0.0% Dist 0.0% Area        ● Moderately Dry (-1.49 to -1) 2.0% Dist 2.2% Area        ● Mildly Dry (-0.99 to -0.5) 88.0% Dist 88.0% Area        ● Mildly Wet (0 to 0.99) 0.0% Dist 0.0% Area        ● Moderately Wet (1 to 1.49) 8.5% Dist 7.6% Area        ● Severely Wet (1.5 to 1.99) 0.0% Dist 0.0% Area        ● Extremely Wet (&gt;=2) 1.4% Dist 1.3% Area</p>	<p style="text-align: center;">NOAA/VIIRS/BLENDED NDVI Composite ending on Week no 50 (09.12.2024 to 15.12.2024) over Agricultural region</p> <p style="text-align: center;"><b>Odisha</b></p>  <p style="font-size: x-small;">Legend for NDVI:        -9999 - 0: Bare soil, rocks clouds, rain, snow, etc.        0 - 0.05        0.05 - 0.1        0.1 - 0.15        0.15 - 0.2        0.2 - 0.25        0.25 - 0.3: Moderate        0.3 - 0.35        0.35 - 0.4        0.4 - 0.45        0.45 - 0.5        0.5 - 0.55        0.55 - 0.6        &gt; 0.6: Good        &gt; 0.6: Very good</p> <p style="text-align: center;">Agriculture vigour is good at few places of the state.</p>
<p style="background-color: #00FFFF; padding: 2px;"><b>As per SPI value (cumulative 4weeks -14 November to 11 December, 2024), mildly wet condition was there in Cuttack district.</b></p>	<p style="background-color: #90EE90; padding: 2px;"><b>According to NDVI map (09.12.2024 to 15.12.2024) the agriculture vigor was light to moderate over most places of Cuttack district.</b></p>

<b>LEGEND (Medium range block level rainfall forecast)</b>		
	No Rainfall	00 mm
	Very light rainfall	01-05 mm
	Light rainfall	06-10 mm
	Light to moderate rainfall	11-30 mm
	Moderate to heavy rainfall	31-65 mm
	Heavy to very heavy rainfall	66-115 mm
	Extremely heavy rainfall	>116 mm




















**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**  
 (A Unit of ICAR-National Rice Research Institute)  
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



**Bulletin No.-547**  
**Date- 24/12/2024**

**Weather (Past and Forecasted) based Disease and pest in Major Crops and vegetables**

Sl No	Crop	Stage and condition of the crop	Anticipated Disease and pest	
1	Pulses (Green gram and black gram)	Vegetative stage		Aphid
1	Mustard	Vegetative stage		Aphid
				Saw fly
2	Rabi Groundnut	Vegetative stage		Leaf Miner
3	Vegetables (Tomato,Chilli, Brinjal)	Vegetative to fruiting		Thrips
				Mites
				Chilli Aphid
4	Brinjal	Fruiting		Fruit and shoot borer
5	Tomato	Vegetative		Bacterial wilt
		Fruiting		Late blight
6	Potato	Sprouting/Vegetative stage		Leaf Blight
7	Cole crops (Cauliflower and cabbage)	Vegetative to fruiting		Leaf eating caterpillar
				Aphid
				Fruit borer
				DBM

Dr Sujata Sethy  
 (Nodal officer)



**Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area**



Debasish Jena  
 SMS (Agrometeorology)  
 Contact No-9861445080/7008453427



**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**  
 (A Unit of ICAR-National Rice Research Institute)  
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



**Bulletin No.-547**  
**Date- 24/12/2024**

**AGROMET ADVISORY BULLETIN FOR ATHAGAD**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 03mm and district received average 06mm rainfall during the last week. The mean daily maximum and minimum temperature was 25.14°C (-3.47) and 17°C (+2.83), respectively. The mean relative humidity was 72-79%.

**Forecasted period: Today(8.30IST) to 29<sup>th</sup> December(8.30IST)2024**

**Weather advisory:** *Moderate rainfall is forecasted for today and tomorrow and dry weather is forecasted for other 3 days in the block.* The day temperature will be 23-31°C and night temperature will be 17-19°C for upcoming 5 days. Maximum and minimum relative humidity will be 53-85% and 35-59% respectively. **There will be partly cloudy to overcast sky for next 5 days.**

Block	Athagad (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	24/12/2024	25/12/2024	26/12/2024	27/12/2024	28/12/2024
Rainfall (mm)	16	13	00	00	00
T-Max (°C)	23	24	29	31	29
T-Min (°C)	19	18	18	18	17
Cloud Cover (octa)	8	4	1	0	0
RH Max (%)	85	78	63	65	53
RH Min (%)	55	59	35	36	35
Wind Speed (kmph)	4	4	7	6	6
Wind Direction (deg)	68	113	30	75	73

**Agromet Advisory**

**General Advisory:** Stop irrigation, fertilizer and pesticide application on today and tomorrow to standing crops like Mustard, rabi groundnut and other rabi vegetables due to forecast of rain. After rain by utilization of rain water farmers should complete rabi oil seeds, pulses and vegetables.

**Pulses:** Forecasted Cloudy and foggy weather thereafter may aggravate the infestation of aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9% EC per litre water for controlling Powdery mildew.

**Rabi-Groundnut:** The early sown rabi groundnut crop is at 3 week stage. Go for weeding and hoeing. Apply 100 kg Gypsum by broadcasting or side dress 50 Kg of Gypsum. Irrigate the crop if required. To manage leaf miner spray Neem Based Pesticide (Azadirachtin) 300 PPM @ 1-litre /acre during primary stage of infestation. To manage chemically spray Profenophos 50% EC @ 400 ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 -litre of water.

**Mustard:** In view of past and forecasted weather condition farmers are advised to don't irrigate the mustard crop. Due to partly cloudy weather, there are chances of Aphid infestation in early sown mustard crop. To manage aphids at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water. There may be chances of white rust disease in Mustard, to control spray Metalaxyl 2g/liter of water.

**Vegetables:** Transplanting of mature seedlings of tomato, cauliflower, cabbage, knol khol and broccoli crop may be done in the raised bed.

**Potato:** After 3 weeks of first earthing up go for hoeing and hand weeding and apply again 26 Kg of Urea and 20 Kg of MOP in line on both side the plants and earthing up should be done to cover up the tubers properly. Apply chlorpyrifos @ 2.5 ml/lit of water against the attack of cut worm and leaf eating insect. Due to forecasted rain and foggy weather in upcoming days there are chances of Leaf Blight disease incidence in potato crop. To manage leaf blight in potato, spray Mancozeb 75% WP @ 600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre.

**Tomato:** Stop irrigation activities as rainfall is forecasted. Due to rain and foggy weather in upcoming days, chances of leaf blight likely to happen in tomato. To avoid blight precaution must be taken during the initial infection stage. Spray 2.5-gram Metalaxyl + Mancozeb/lit of water To manage Whitefly chemically sprays Thiamethoxam 25 % WG @ 40-gram/acre or Pyriproxifen 10% EC @ 200-ml/acre or Spiromesifen 22.9% SC @ 250-ml/acre by mixing in 200- litre of water. If Bacterial wilt will occur in brinjal and tomato. To control them, drench with copper oxychloride @ 3g+ plantomycin @ 1 g/ litre of water.

**Onion:** In present weather conditions, constant monitoring of timely sown onion crop against attack of thrips and infection of purple blotch should be done. Need based spraying of Diethane M -45 @ 3 grams per litre of water with sticky material are advised.

**Brinjal and Chilly:** Shoot and fruit borer and leaf eating caterpillar may observe in brinjal field, to control it spray Emamectin benzoate 5% SG @ 0.4 g/litre of water or Chlorantranilprole 18.5% SC @ 0.4 ml/litre of water. Spraying should be done during the early morning or evening hours.

**Cabbage and Cauliflower:** Browning or Brown rot is a common physiological disorder in cauliflower crop due to deficiency of Boron. This may be controlled by application of borax @ 4-kg/ acre as soil application during transplanting. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Due to slight increase in temperature Cole crops may get affected by diamond back moth. To control this apply Spinosad 45SC @ 0.5ml/l. There is a chance of occurrence of thrip, mites and Aphids in cole crops, for control spray imidachloprid 0.3 ml /liter or fipronil 2 ml /liter of water. Leaf eating caterpillar and fruit borer may infest the crops. Spray fipronil 5% SL 400ml or emamectinbenzoate 5% SG 80gm or indoxacarb 14.5% SC 200ml mixing in 200lit of water per acre crop.

**Animal husbandry:**

In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Give animals mineral mixture along with salt regularly and also wheat grains, jaggery etc. @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals. • In Poultry, keep the chicks warm by providing artificial light in the poultry sheds.

**Dr Sujata Sethy**  
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



**Debasish Jena**

**SMS (Agrometeorology)**

**Contact No-9861445080/7008453427**



**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**  
(A Unit of ICAR-National Rice Research Institute)  
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



**Bulletin No.-547**  
**Date- 24/12/2024**

**AGROMET ADVISORY BULLETIN FOR BADAMBA**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 02mm and district received average 06mm rainfall during the last week. The mean daily maximum and minimum temperature was 25.14°C (-3.47) and 17°C (+2.83), respectively. The mean relative humidity was 72-79%.

**Forecasted period: Today(8.30IST) to 29<sup>th</sup> December (8.30IST)2024**

**Weather advisory:** Moderate rainfall is forecasted for today and tomorrow and dry weather is forecasted for other 3 days in the block. The day temperature will be 22-30°C and night temperature will be 16-18°C for upcoming 5 days. Maximum and minimum relative humidity will be 62-89% and 35-66% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Badamba (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	24/12/2024	25/12/2024	26/12/2024	27/12/2024	28/12/2024
Rainfall(mm)	16	13	00	00	00
T-Max(°C)	22	22	29	30	29
T-Min(°C)	17	17	17	18	16
Cloud Cover(octa)	8	5	0	0	0
RH Max (%)	89	87	62	69	62
RH Min (%)	54	66	35	36	35
Wind Speed (Kmph)	3	3	6	5	5
Wind Direction (deg)	63	315	113	72	122

**Agromet Advisory**

**General Advisory:** Stop irrigation, fertilizer and pesticide application on today and tomorrow to standing crops like Mustard, rabi groundnut and other rabi vegetables due to forecast of rain. After rain by utilization of rain water farmers should complete rabi oil seeds, pulses and vegetables.

**Pulses:** Forecasted Cloudy and foggy weather thereafter may aggravate the infestation of aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9% EC per litre water for controlling Powdery mildew.

**Rabi-Groundnut:** The early sown rabi groundnut crop is at 3 week stage. Go for weeding and hoeing. Apply 100 kg Gypsum by broadcasting or side dress 50 Kg of Gypsum. Irrigate the crop if required. To manage leaf miner spray Neem Based Pesticide (Azadirachtin) 300 PPM @ 1-litre /acre during primary stage of infestation. To manage chemically spray Profenophos 50% EC @ 400 ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 -litre of water.

**Mustard:** In view of past and forecasted weather condition farmers are advised to don't irrigate the mustard crop. Due to partly cloudy weather, there are chances of Aphid infestation in early sown mustard crop. To manage aphids at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water. There may be chances of white rust disease in Mustard, to control spray Metalaxyl 2g/liter of water.

**Vegetables:**

**Transplanting of mature seedlings of tomato, cauliflower, cabbage, knol khol and broccoli crop may be done in the raised bed.**

**Potato:** After 3 weeks of first earthing up go for hoeing and hand weeding and apply again 26 Kg of Urea and 20 Kg of MOP in line on both side the plants and earthing up should be done to cover up the tubers properly. Apply chlorpyrifos @ 2.5 ml/lit of water against the attack of cut worm and leaf eating insect. Due to forecasted rain and foggy weather in upcoming days there are chances of Leaf Blight disease incidence in potato crop. To manage leaf blight in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre.

**Tomato:** Stop irrigation activities as rainfall is forecasted. Due to rain and foggy weather in upcoming days, chances of leaf blight likely to happen in tomato. To avoid blight precaution must be taken during the initial infection stage. Spray 2.5-gram Metalaxyl + Mancozeb/lit of water To manage Whitefly chemically sprays Thiamethoxam 25 % WG @ 40-gram/acre or Pyriproxifen 10% EC @ 200-ml/acre or Spiromesifen 22.9% SC @ 250-ml/acre by mixing in 200- litre of water. If Bacterial wilt will occur in brinjal and tomato. To control them, drench with copper oxychloride @ 3g+ plantomycin @1 g/ litre of water.

**Onion:** In present weather conditions, constant monitoring of timely sown onion crop against attack of thrips and infection of purple blotch should be done. Need based spraying of Diethane M -45 @ 3 grams per litre of water with sticky material are advised.

**Brinjal and Chilly:** Shoot and fruit borer and leaf eating caterpillar may observe in brinjal field, to control it spray Emamectin benzoate 5% SG @ 0.4 g/litre of water or Chlorantraniliprole 18.5% SC @ 0.4 ml/litre of water. Spraying should be done during the early morning or evening hours.

**Cabbage and Cauliflower:** Browning or Brown rot is a common physiological disorder in cauliflower crop due to deficiency of Boron. This may be controlled by application of borax @ 4-kg/ acre as soil application during transplanting. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Due to slight increase in temperature Cole crops may get affected by diamond back moth. To control this apply Spinosad 45SC @ 0.5ml/l. There is a chance of occurrence of thrip, mites and Aphids in cole crops, for control spray imidachloprid 0.3 ml /liter or fipronil 2 ml /liter of water. Leaf eating caterpillar and fruit borer may infest the crops. Spray fipronil 5% SL 400ml or emamectinbenzoate 5% SG 80gm or indoxacarb 14.5% SC 200ml mixing in 200lit of water per acre crop.

**Animal husbandry:**

In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to direct air. Give animals mineral mixture along with salt regularly and also wheat grains, jaggery etc. @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals. • In Poultry, keep the chicks warm by providing artificial light in the poultry sheds.

**Dr Sujata Sethy**  
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



**Debasish Jena**

**SMS (Agrometeorology)**

**Contact No-9861445080/7008453427**



**GRAMIN KRISHI SAMAKSHEPA SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**  
 (A Unit of ICAR-National Rice Research Institute)  
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



**Bulletin No.-547**  
**Date- 24/12/2024**

**AGROMET ADVISORY BULLETIN FOR BANKI**

**District Past week average weather condition**

The district received average 06mm rainfall during the last week. The mean daily maximum and minimum temperature was 25.14°C (-3.47) and 17°C (+2.83), respectively. The mean relative humidity was 72-79%.

**Forecasted period: Today(8.30IST) to 29<sup>th</sup> December (8.30IST)2024**

**Weather advisory:** Moderate rainfall is forecasted for today and tomorrow and dry weather is forecasted for other 3 days in the block. The day temperature will be 22-31°C and night temperature will be 17-18°C for upcoming 5 days. Maximum and minimum relative humidity will be 60-87% and 33-67% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Banki (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	24/12/2024	25/12/2024	26/12/2024	27/12/2024	28/12/2024
Rainfall(mm)	18	15	00	00	00
T-Max(°C)	22	23	29	31	29
T-Min(°C)	18	17	17	18	17
Cloud Cover(octa)	8	5	1	0	0
RH Max (%)	87	82	62	67	60
RH Min (%)	56	67	35	36	33
Wind Speed (Kmph)	4	4	7	6	6
Wind Direction (deg)	64	113	23	72	75

**Agromet Advisory**

**General Advisory:** Stop irrigation, fertilizer and pesticide application on today and tomorrow to standing crops like Mustard, rabi groundnut and other rabi vegetables due to forecast of rain. After rain by utilization of rain water farmers should complete rabi oil seeds, pulses and vegetables.

**Pulses:** Forecasted Cloudy and foggy weather thereafter may aggravate the infestation of aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9%EC per litre water for controlling Powdery mildew.

**Rabi-Groundnut:** The early sown rabi groundnut crop is at 3 week stage. Go for weeding and hoeing. Apply 100 kg Gypsum by broadcasting or side dress 50 Kg of Gypsum. Irrigate the crop if required. To manage leaf miner spray Neem Based Pesticide (Azadirachtin) 300 PPM @ 1-litre /acre during primary stage of infestation. To manage chemically spray Profenophos 50% EC @ 400 ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 -litre of water.

**Mustard:** In view of past and forecasted weather condition farmers are advised to don't irrigate the mustard crop. Due to partly cloudy weather, there are chances of Aphid infestation in early sown mustard crop. To manage aphids at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water. There may be chances of white rust disease in Mustard, to control spray Metalaxyl 2g/liter of water.

**Vegetables:**

**Transplanting of mature seedlings of tomato, cauliflower, cabbage, knol khol and broccoli crop may be done in the raised bed.**

**Potato:** After 3 weeks of first earthing up go for hoeing and hand weeding and apply again 26 Kg of Urea and 20 Kg of MOP in line on both side the plants and earthing up should be done to cover up the tubers properly. Apply chlorpyrifos @ 2.5 ml/lit of water against the attack of cut worm and leaf eating insect. Due to forecasted rain and foggy weather in upcoming days there are chances of Leaf Blight disease incidence in potato crop. To manage leaf blight in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre.

**Tomato:** Stop irrigation activities as rainfall is forecasted. Due to rain and foggy weather in upcoming days, chances of leaf blight likely to happen in tomato. To avoid blight precaution must be taken during the initial infection stage. Spray 2.5-gram Metalaxyl + Mancozeb/lit of water To manage Whitefly chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Pyriproxifen 10% EC @ 200-ml/acre or Spiromesifen 22.9% SC @ 250-ml/acre by mixing in 200- litre of water. If Bacterial wilt will occur in brinjal and tomato. To control them, drench with copper oxychloride @ 3g+ plantomycin @1 g/ litre of water.

**Onion:** In present weather conditions, constant monitoring of timely sown onion crop against attack of thrips and infection of purple blotch should be done. Need based spraying of Diethane M -45 @ 3 grams per litre of water with sticky material are advised.

**Brinjal and Chilly:** Shoot and fruit borer and leaf eating caterpillar may observe in brinjal field, to control it spray Emamectin benzoate 5% SG @ 0.4 g/litre of water or Chlorantranilprole 18.5% SC @ 0.4 ml/litre of water. Spraying should be done during the early morning or evening hours.

**Cabbage and Cauliflower:** Browning or Brown rot is a common physiological disorder in cauliflower crop due to deficiency of Boron. This may be controlled by application of borax @ 4-kg/ acre as soil application during transplanting. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Due to slight increase in temperature Cole crops may get affected by diamond back moth. To control this apply Spinosad 45SC @ 0.5ml/l. There is a chance of occurrence of thrip, mites and Aphids in cole crops, for control spray imidachloprid 0.3 ml /liter or fipronil 2 ml /liter of water. Leaf eating caterpillar and fruit borer may infest the crops. Spray fipronil 5% SL 400ml or emamectinbenzoate 5% SG 80gm or indoxacarb 14.5% SC 200ml mixing in 200lit of water per acre crop.

**Animal husbandry:**

In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Give animals mineral mixture along with salt regularly and also wheat grains, jaggery etc. @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals. • In Poultry, keep the chicks warm by providing artificial light in the poultry sheds.

**Dr Sujata Sethy**  
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



**Debasish Jena**

**SMS (Agrometeorology)**

**Contact No-9861445080/7008453427**



**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**  
 (A Unit of ICAR-National Rice Research Institute)  
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



**Bulletin No.-547**  
**Date- 24/12/2024**

**AGROMET ADVISORY BULLETIN FOR BANKI-DAMPADA**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 08mm and district received average 06mm rainfall during the last week. The mean daily maximum and minimum temperature was 25.14°C (-3.47) and 17°C (+2.83), respectively. The mean relative humidity was 72-79%.

**Forecasted period: Today(8.30IST) to 29<sup>th</sup> December (8.30IST)2024**

**Weather Advisory:** Moderate rainfall is forecasted for today and tomorrow and dry weather is forecasted for other 3 days in the block. The day temperature will be 22-31°C and night temperature will be 17-18°C for upcoming 5 days. Maximum and minimum relative humidity will be 59-87% and 33-67% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Banki-Dampada (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	24/12/2024	25/12/2024	26/12/2024	27/12/2024	28/12/2024
Rainfall(mm)	18	15	00	00	00
T-Max(°C)	22	23	29	31	29
T-Min(°C)	18	17	17	18	17
Cloud Cover(octa)	8	5	1	0	0
RH Max (%)	87	81	62	66	59
RH Min (%)	56	67	36	36	33
Wind Speed (Kmph)	4	4	7	6	6
Wind Direction (deg)	64	113	23	75	70

**Agromet Advisory**

**General Advisory:** Stop irrigation, fertilizer and pesticide application on today and tomorrow to standing crops like Mustard, rabi groundnut and other rabi vegetables due to forecast of rain. After rain by utilization of rain water farmers should complete rabi oil seeds, pulses and vegetables.

**Pulses:** Forecasted Cloudy and foggy weather thereafter may aggravate the infestation of aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9% EC per litre water for controlling Powdery mildew.

**Rabi-Groundnut:** The early sown rabi groundnut crop is at 3 week stage. Go for weeding and hoeing. Apply 100 kg Gypsum by broadcasting or side dress 50 Kg of Gypsum. Irrigate the crop if required. To manage leaf miner spray Neem Based Pesticide (Azadirachtin) 300 PPM @ 1-litre /acre during primary stage of infestation. To manage chemically spray Profenophos 50% EC @ 400 ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 -litre of water.

**Mustard:** In view of past and forecasted weather condition farmers are advised to don't irrigate the mustard crop. Due to partly cloudy weather, there are chances of Aphid infestation in early sown mustard crop. To manage aphids at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water. There may be chances of white rust disease in Mustard, to control spray Metalaxyl 2g/liter of water.

**Vegetables:**

**Transplanting of mature seedlings of tomato, cauliflower, cabbage, knol khol and broccoli crop may be done in the raised bed.**

**Potato:** After 3 weeks of first earthing up go for hoeing and hand weeding and apply again 26 Kg of Urea and 20 Kg of MOP in line on both side the plants and earthing up should be done to cover up the tubers properly. Apply chlorpyrifos @ 2.5 ml/lit of water against the attack of cut worm and leaf eating insect. Due to forecasted rain and foggy weather in upcoming days there are chances of Leaf Blight disease incidence in potato crop. To manage leaf blight in potato, spray Mancozeb 75% WP @ 600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre.

**Tomato:** Stop irrigation activities as rainfall is forecasted. Due to rain and foggy weather in upcoming days, chances of leaf blight likely to happen in tomato. To avoid blight precaution must be taken during the initial infection stage. Spray 2.5-gram Metalaxyl + Mancozeb/lit of water To manage Whitefly chemically sprays Thiamethoxam 25 % WG @ 40-gram/acre or Pyriproxifen 10% EC @ 200-ml/acre or Spiromesifen 22.9% SC @ 250-ml/acre by mixing in 200- litre of water. If Bacterial wilt will occur in brinjal and tomato. To control them, drench with copper oxychloride @ 3g+ plantomycin @1 g/ litre of water.

**Onion:** In present weather conditions, constant monitoring of timely sown onion crop against attack of thrips and infection of purple blotch should be done. Need based spraying of Diethane M -45 @ 3 grams per litre of water with sticky material are advised.

**Brinjal and Chilly:** Shoot and fruit borer and leaf eating caterpillar may observe in brinjal field, to control it spray Emamectin benzoate 5% SG @ 0.4 g/litre of water or Chlorantranilprole 18.5% SC @ 0.4 ml/litre of water. Spraying should be done during the early morning or evening hours.

**Cabbage and Cauliflower:** Browning or Brown rot is a common physiological disorder in cauliflower crop due to deficiency of Boron. This may be controlled by application of borax @ 4-kg/ acre as soil application during transplanting. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Due to slight increase in temperature Cole crops may get affected by diamond back moth. To control this apply Spinosad 45SC @ 0.5ml/l. There is a chance of occurrence of thrip, mites and Aphids in cole crops, for control spray imidachloprid 0.3 ml /liter or fipronil 2 ml /liter of water. Leaf eating caterpillar and fruit borer may infest the crops. Spray fipronil 5% SL 400ml or emamectinbenzoate 5% SG 80gm or indoxacarb 14.5% SC 200ml mixing in 200lit of water per acre crop.

**Animal husbandry:**

- In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Give animals mineral mixture along with salt regularly and also wheat grains, jaggery etc. @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals. • In Poultry, keep the chicks warm by providing artificial light in the poultry sheds.

Dr Sujata Sethy  
 (Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427



**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**  
(A Unit of ICAR-National Rice Research Institute)  
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



**Bulletin No.-547**  
**Date- 24/12/2024**

**AGROMET ADVISORY BULLETIN FOR BARANGA**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 07mm and district received average 06mm rainfall during the last week. The mean daily maximum and minimum temperature was 25.14°C (-3.47) and 17°C (+2.83), respectively. The mean relative humidity was 72-79%.

**Forecasted period: Today(8.30IST) to 29<sup>th</sup> December (8.30IST)2024**

**Weather advisory:** Moderate rainfall is forecasted for today and tomorrow and dry weather is forecasted for other 3 days in the block. The day temperature will be 22-31°C and night temperature will be 18-19°C for upcoming 5 days. Maximum and minimum relative humidity will be 51-85% and 34-64% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Baranga (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	24/12/2024	25/12/2024	26/12/2024	27/12/2024	28/12/2024
Rainfall(mm)	19	14	00	00	00
T-Max(°C)	22	23	28	31	30
T-Min(°C)	19	18	18	18	18
Cloud Cover	8	6	2	0	0
RH Max (%)	85	83	63	61	51
RH Min (%)	60	64	39	36	34
Wind Speed (Kmph)	6	5	8	8	8
Wind Direction (deg)	341	23	25	23	64

**Agromet Advisory**

**General Advisory:** Stop irrigation, fertilizer and pesticide application on today and tomorrow to standing crops like Mustard, rabi groundnut and other rabi vegetables due to forecast of rain. After rain by utilization of rain water farmers should complete rabi oil seeds, pulses and vegetables.

**Pulses:** Forecasted Cloudy and foggy weather thereafter may aggravate the infestation of aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9% EC per litre water for controlling Powdery mildew.

**Rabi-Groundnut:** The early sown rabi groundnut crop is at 3 week stage. Go for weeding and hoeing. Apply 100 kg Gypsum by broadcasting or side dress 50 Kg of Gypsum. Irrigate the crop if required. To manage leaf miner spray Neem Based Pesticide (Azadirachtin) 300 PPM @ 1-litre /acre during primary stage of infestation. To manage chemically spray Profenophos 50% EC @ 400 ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 -litre of water.

**Mustard:** In view of past and forecasted weather condition farmers are advised to don't irrigate the mustard crop. Due to partly cloudy weather, there are chances of Aphid infestation in early sown mustard crop. To manage aphids at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water. There may be chances of white rust disease in Mustard, to control spray Metalaxyl 2g/liter of water.

**Vegetables:**

**Transplanting of mature seedlings of tomato, cauliflower, cabbage, knol khol and broccoli crop may be done in the raised bed.**

**Potato:** After 3 weeks of first earthing up go for hoeing and hand weeding and apply again 26 Kg of Urea and 20 Kg of MOP in line on both side the plants and earthing up should be done to cover up the tubers properly. Apply chlorpyrifos @ 2.5 ml/lit of water against the attack of cut worm and leaf eating insect. Due to forecasted rain and foggy weather in upcoming days there are chances of Leaf Blight disease incidence in potato crop. To manage leaf blight in potato, spray Mancozeb 75% WP @ 600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre.

**Tomato:** Stop irrigation activities as rainfall is forecasted. Due to rain and foggy weather in upcoming days, chances of leaf blight likely to happen in tomato. To avoid blight precaution must be taken during the initial infection stage. Spray 2.5-gram Metalaxyl + Mancozeb/lit of water To manage Whitefly chemically sprays Thiamethoxam 25 % WG @ 40-gram/acre or Pyriproxifen 10% EC @ 200-ml/acre or Spiromesifen 22.9% SC @ 250-ml/acre by mixing in 200- litre of water. If Bacterial wilt will occur in brinjal and tomato. To control them, drench with copper oxychloride @ 3g+ plantomycin @1 g/ litre of water.

**Onion:** In present weather conditions, constant monitoring of timely sown onion crop against attack of thrips and infection of purple blotch should be done. Need based spraying of Diethane M -45 @ 3 grams per litre of water with sticky material are advised.

**Brinjal and Chilly:** Shoot and fruit borer and leaf eating caterpillar may observe in brinjal field, to control it spray Emamectin benzoate 5% SG @ 0.4 g/litre of water or Chlorantranilprole 18.5% SC @ 0.4 ml/litre of water. Spraying should be done during the early morning or evening hours.

**Cabbage and Cauliflower:** Browning or Brown rot is a common physiological disorder in cauliflower crop due to deficiency of Boron. This may be controlled by application of borax @ 4-kg/ acre as soil application during transplanting. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Due to slight increase in temperature Cole crops may get affected by diamond back moth. To control this apply Spinosad 45SC @ 0.5ml/l. There is a chance of occurrence of thrip, mites and Aphids in cole crops, for control spray imidachloprid 0.3 ml /liter or fipronil 2 ml /liter of water. Leaf eating caterpillar and fruit borer may infest the crops. Spray fipronil 5% SL 400ml or emamectinbenzoate 5% SG 80gm or indoxacarb 14.5% SC 200ml mixing in 200lit of water per acre crop.

**Animal husbandry:**

- In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Give animals mineral mixture along with salt regularly and also wheat grains, jaggery etc. @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals.
- In Poultry, keep the chicks warm by providing artificial light in the poultry sheds.

Dr Sujata Sethy  
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena

SMS (Agrometeorology)

Contact No-9861445080/7008453427





**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**  
**(A Unit of ICAR-National Rice Research Institute)**  
**Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com**



**Bulletin No.-547**  
**Date- 24/12/2024**

**AGROMET ADVISORY BULLETIN FOR CUTTACK SADAR**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 06mm and district received average 06mm rainfall during the last week. The mean daily maximum and minimum temperature was 25.14°C (-3.47) and 17°C (+2.83), respectively. The mean relative humidity was 72-79%.

**Forecasted period: Today(8.30IST) to 29<sup>th</sup> December (8.30IST)2024**

**Weather advisory:** Moderate rainfall is forecasted for today and tomorrow and dry weather is forecasted for other 3 days in the block. The day temperature will be 23-31°C and night temperature will be 17-19°C for upcoming 5 days. Maximum and minimum relative humidity will be 48-84% and 34-58% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Cuttack sadar (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	24/12/2024	25/12/2024	26/12/2024	27/12/2024	28/12/2024
Rainfall(mm)	15	12	00	00	00
T-Max(°C)	23	24	28	31	30
T-Min(°C)	19	18	19	18	17
Cloud Cover(octa)	8	4	1	0	0
RH Max (%)	84	80	62	62	48
RH Min (%)	57	58	37	36	34
Wind Speed (Kmph)	5	5	8	8	8
Wind Direction (deg)	341	15	30	75	68

**Agromet Advisory**

**General Advisory:** Stop irrigation, fertilizer and pesticide application on today and tomorrow to standing crops like Mustard, rabi groundnut and other rabi vegetables due to forecast of rain. After rain by utilization of rain water farmers should complete rabi oil seeds, pulses and vegetables.

**Pulses:** Forecasted Cloudy and foggy weather thereafter may aggravate the infestation of aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9% EC per litre water for controlling Powdery mildew.

**Rabi-Groundnut:** The early sown rabi groundnut crop is at 3 week stage. Go for weeding and hoeing. Apply 100 kg Gypsum by broadcasting or side dress 50 Kg of Gypsum. Irrigate the crop if required. To manage leaf miner spray Neem Based Pesticide (Azadirachtin) 300 PPM @ 1-litre /acre during primary stage of infestation. To manage chemically spray Profenophos 50% EC @ 400 ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 -litre of water.

**Mustard:** In view of past and forecasted weather condition farmers are advised to don't irrigate the mustard crop. Due to partly cloudy weather, there are chances of Aphid infestation in early sown mustard crop. To manage aphids at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water. There may be chances of white rust disease in Mustard, to control spray Metalaxyl 2g/liter of water.

**Vegetables:**

**Transplanting of mature seedlings of tomato, cauliflower, cabbage, knol khol and broccoli crop may be done in the raised bed.**

**Potato:** After 3 weeks of first earthing up go for hoeing and hand weeding and apply again 26 Kg of Urea and 20 Kg of MOP in line on both side the plants and earthing up should be done to cover up the tubers properly. Apply chlorpyrifos @ 2.5 ml/lit of water against the attack of cut worm and leaf eating insect. Due to forecasted rain and foggy weather in upcoming days there are chances of Leaf Blight disease incidence in potato crop. To manage leaf blight in potato, spray Mancozeb 75% WP @ 600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre.

**Tomato:** Stop irrigation activities as rainfall is forecasted. Due to rain and foggy weather in upcoming days, chances of leaf blight likely to happen in tomato. To avoid blight precaution must be taken during the initial infection stage. Spray 2.5-gram Metalaxyl + Mancozeb/lit of water To manage Whitefly chemically sprays Thiamethoxam 25 % WG @ 40-gram/acre or Pyriproxifen 10% EC @ 200-ml/acre or Spiromesifen 22.9% SC @ 250-ml/acre by mixing in 200- litre of water. If Bacterial wilt will occur in brinjal and tomato. To control them, drench with copper oxychloride @ 3g+ plantomycin @1 g/ litre of water.

**Onion:** In present weather conditions, constant monitoring of timely sown onion crop against attack of thrips and infection of purple blotch should be done. Need based spraying of Diethane M -45 @ 3 grams per litre of water with sticky material are advised.

**Brinjal and Chilly:** Shoot and fruit borer and leaf eating caterpillar may observe in brinjal field, to control it spray Emamectin benzoate 5% SG @ 0.4 g/litre of water or Chlorantranilprole 18.5% SC @ 0.4 ml/litre of water. Spraying should be done during the early morning or evening hours.

**Cabbage and Cauliflower:** Browning or Brown rot is a common physiological disorder in cauliflower crop due to deficiency of Boron. This may be controlled by application of borax @ 4-kg/ acre as soil application during transplanting. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Due to slight increase in temperature Cole crops may get affected by diamond back moth. To control this apply Spinosad 45SC @ 0.5ml/l. There is a chance of occurrence of thrip, mites and Aphids in cole crops, for control spray imidachloprid 0.3 ml /liter or fipronil 2 ml /liter of water. . Leaf eating caterpillar and fruit borer may infest the crops. Spray fipronil 5% SL 400ml or emamectinbenzoate 5% SG 80gm or indoxacarb 14.5% SC 200ml mixing in 200lit of water per acre crop.

**Animal husbandry:**

In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Give animals mineral mixture along with salt regularly and also wheat grains, jaggery etc. @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals. • In Poultry, keep the chicks warm by providing artificial light in the poultry sheds.

**Dr Sujata Sethy**  
**(Nodal officer)**



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



**Debasish Jena**

**SMS (Agrometeorology)**

**Contact No-9861445080/7008453427**



**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**  
 (A Unit of ICAR-National Rice Research Institute)  
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



**Bulletin No.-547**  
**Date- 24/12/2024**

**AGROMET ADVISORY BULLETIN FOR KANTAPADA**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 11mm and district received average 06mm rainfall during the last week. The mean daily maximum and minimum temperature was 25.14°C (-3.47) and 17°C (+2.83), respectively. The mean relative humidity was 72-79%.

**Forecasted period: Today(8.30IST) to 29<sup>th</sup> December (8.30IST)2024**

**Weather advisory:** Moderate rainfall is forecasted for today and tomorrow and dry weather is forecasted for other 3 days in the block. The day temperature will be 22-30°C and night temperature will be 18-19°C for upcoming 5 days. Maximum and minimum relative humidity will be 54-84% and 34-64% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Kantapada (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	24/12/2024	25/12/2024	26/12/2024	27/12/2024	28/12/2024
Rainfall(mm)	17	14	00	00	00
T-Max(°C)	23	22	27	30	30
T-Min(°C)	19	18	18	19	18
Cloud Cover(octa)	8	6	2	0	0
RH Max (%)	83	84	65	61	54
RH Min (%)	62	64	51	38	34
Wind Speed (Kmph)	8	6	8	9	9
Wind Direction (deg)	360	26	28	30	56

**Agromet Advisory**

**General Advisory:** Stop irrigation, fertilizer and pesticide application on today and tomorrow to standing crops like Mustard, rabi groundnut and other rabi vegetables due to forecast of rain. After rain by utilization of rain water farmers should complete rabi oil seeds, pulses and vegetables.

**Pulses:** Forecasted Cloudy and foggy weather thereafter may aggravate the infestation of aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9% EC per litre water for controlling Powdery mildew.

**Rabi-Groundnut:** The early sown rabi groundnut crop is at 3 week stage. Go for weeding and hoeing. Apply 100 kg Gypsum by broadcasting or side dress 50 Kg of Gypsum. Irrigate the crop if required. To manage leaf miner spray Neem Based Pesticide (Azadirachtin) 300 PPM @ 1-litre /acre during primary stage of infestation. To manage chemically spray Profenophos 50% EC @ 400 ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 -litre of water.

**Mustard:** In view of past and forecasted weather condition farmers are advised to don't irrigate the mustard crop. Due to partly cloudy weather, there are chances of Aphid infestation in early sown mustard crop. To manage aphids at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water. There may be chances of white rust disease in Mustard, to control spray Metalaxyl 2g/liter of water.

**Vegetables:**

**Transplanting of mature seedlings of tomato, cauliflower, cabbage, knol khol and broccoli crop may be done in the raised bed.**

**Potato:** After 3 weeks of first earthing up go for hoeing and hand weeding and apply again 26 Kg of Urea and 20 Kg of MOP in line on both side the plants and earthing up should be done to cover up the tubers properly. Apply chlorpyrifos @ 2.5 ml/lit of water against the attack of cut worm and leaf eating insect. Due to forecasted rain and foggy weather in upcoming days there are chances of Leaf Blight disease incidence in potato crop. To manage leaf blight in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre.

**Tomato:** Stop irrigation activities as rainfall is forecasted. Due to rain and foggy weather in upcoming days, chances of leaf blight likely to happen in tomato. To avoid blight precaution must be taken during the initial infection stage. Spray 2.5-gram Metalaxyl + Mancozeb/lit of water. To manage Whitefly chemically sprays Thiamethoxam 25 % WG @ 40-gram/acre or Pyriproxifen 10% EC @ 200-ml/acre or Spiromesifen 22.9% SC @ 250-ml/acre by mixing in 200- litre of water. If Bacterial wilt will occur in brinjal and tomato. To control them, drench with copper oxychloride @ 3g+ plantomycin @1 g/ litre of water.

**Onion:** In present weather conditions, constant monitoring of timely sown onion crop against attack of thrips and infection of purple blotch should be done. Need based spraying of Diethane M -45 @ 3 grams per litre of water with sticky material are advised.

**Brinjal and Chilly:** Shoot and fruit borer and leaf eating caterpillar may observe in brinjal field, to control it spray Emamectin benzoate 5% SG @ 0.4 g/litre of water or Chlorantraniliprole 18.5% SC @ 0.4 ml/litre of water. Spraying should be done during the early morning or evening hours.

**Cabbage and Cauliflower:** Browning or Brown rot is a common physiological disorder in cauliflower crop due to deficiency of Boron. This may be controlled by application of borax @ 4-kg/ acre as soil application during transplanting. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Due to slight increase in temperature Cole crops may get affected by diamond back moth. To control this apply Spinosad 45SC @ 0.5ml/l. There is a chance of occurrence of thrip, mites and Aphids in cole crops, for control spray imidachloprid 0.3 ml /liter or fipronil 2 ml /liter of water. Leaf eating caterpillar and fruit borer may infest the crops. Spray fipronil 5% SL 400ml or emamectinbenzoate 5% SG 80gm or indoxacarb 14.5% SC 200ml mixing in 200lit of water per acre crop.

**Animal husbandry:**

In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Give animals mineral mixture along with salt regularly and also wheat grains, jaggery etc. @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals. • In Poultry, keep the chicks warm by providing artificial light in the poultry sheds.

**Dr Sujata Sethy**  
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



**Debasish Jena**  
**SMS (Agrometeorology)**  
 Contact No-9861445080/7008453427



**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**  
**(A Unit of ICAR-National Rice Research Institute)**  
**Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com**



**Bulletin No.-547**  
**Date- 24/12/2024**

**AGROMET ADVISORY BULLETIN FOR MAHANGA**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 06mm and district received average 06mm rainfall during the last week. The mean daily maximum and minimum temperature was 25.14°C (-3.47) and 17°C (+2.83), respectively. The mean relative humidity was 72-79%.

**Forecasted period: Today(8.30IST) to 29<sup>th</sup> December (8.30IST)2024**

**Weather advisory:** Moderate rainfall is forecasted for today and tomorrow and dry weather is forecasted for other 3 days in the block. The day temperature will be and it will be 23-30°C and night temperature will be 17-19°C for upcoming 5 days. Maximum and minimum relative humidity will be 48-82% and 32-59% respectively. **There will be partly cloudy to overcast sky for next 5 days.**

Block	Mahanga (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	24/12/2024	25/12/2024	26/12/2024	27/12/2024	28/12/2024
Rainfall(mm)	14	12	00	00	00
T-Max(°C)	23	24	28	30	30
T-Min(°C)	19	18	18	18	17
Cloud Cover(octal)	8	4	1	0	0
RH Max (%)	82	72	63	64	48
RH Min (%)	59	54	39	37	32
Wind Speed (Kmph)	5	5	10	9	9
Wind Direction (deg)	30	23	30	15	30

**Agromet Advisory**

**General Advisory:** Stop irrigation, fertilizer and pesticide application on today and tomorrow to standing crops like Mustard, rabi groundnut and other rabi vegetables due to forecast of rain. After rain by utilization of rain water farmers should complete rabi oil seeds, pulses and vegetables.

**Pulses:** Forecasted Cloudy and foggy weather thereafter may aggravate the infestation of aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9% EC per litre water for controlling Powdery mildew.

**Rabi-Groundnut:** The early sown rabi groundnut crop is at 3 week stage. Go for weeding and hoeing. Apply 100 kg Gypsum by broadcasting or side dress 50 Kg of Gypsum. Irrigate the crop if required. To manage leaf miner spray Neem Based Pesticide (Azadirachtin) 300 PPM @ 1-litre /acre during primary stage of infestation. To manage chemically spray Profenophos 50% EC @ 400 ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 -litre of water.

**Mustard:** In view of past and forecasted weather condition farmers are advised to don't irrigate the mustard crop. Due to partly cloudy weather, there are chances of Aphid infestation in early sown mustard crop. To manage aphids at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water. There may be chances of white rust disease in Mustard, to control spray Metalaxyl 2g/liter of water.

**Vegetables:**

**Transplanting of mature seedlings of tomato, cauliflower, cabbage, knol khol and broccoli crop may be done in the raised bed.**

**Potato:** After 3 weeks of first earthing up go for hoeing and hand weeding and apply again 26 Kg of Urea and 20 Kg of MOP in line on both side the plants and earthing up should be done to cover up the tubers properly. Apply chlorpyrifos @ 2.5 ml/lit of water against the attack of cut worm and leaf eating insect. Due to forecasted rain and foggy weather in upcoming days there are chances of Leaf Blight disease incidence in potato crop. To manage leaf blight in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre.

**Tomato:** Stop irrigation activities as rainfall is forecasted. Due to rain and foggy weather in upcoming days, chances of leaf blight likely to happen in tomato. To avoid blight precaution must be taken during the initial infection stage. Spray 2.5-gram Metalaxyl + Mancozeb/lit of water. To manage Whitefly chemically sprays Thiamethoxam 25 % WG @ 40-gram/acre or Pyriproxifen 10% EC @ 200-ml/acre or Spiromesifen 22.9% SC @ 250-ml/acre by mixing in 200- litre of water. If Bacterial wilt will occur in brinjal and tomato. To control them, drench with copper oxychloride @ 3g+ plantomycin @1 g/ litre of water.

**Onion:** In present weather conditions, constant monitoring of timely sown onion crop against attack of thrips and infection of purple blotch should be done. Need based spraying of Diethane M -45 @ 3 grams per litre of water with sticky material are advised.

**Brinjal and Chilly:** Shoot and fruit borer and leaf eating caterpillar may observe in brinjal field, to control it spray Emamectin benzoate 5% SG @ 0.4 g/litre of water or Chlorantraniliprole 18.5% SC @ 0.4 ml/litre of water. Spraying should be done during the early morning or evening hours.

**Cabbage and Cauliflower:** Browning or Brown rot is a common physiological disorder in cauliflower crop due to deficiency of Boron. This may be controlled by application of borax @ 4-kg/ acre as soil application during transplanting. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Due to slight increase in temperature Cole crops may get affected by diamond back moth. To control this apply Spinosad 45SC @ 0.5ml/l. There is a chance of occurrence of thrip, mites and Aphids in cole crops, for control spray imidachloprid 0.3 ml /liter or fipronil 2 ml /liter of water. Leaf eating caterpillar and fruit borer may infest the crops. Spray fipronil 5% SL 400ml or emamectinbenzoate 5% SG 80gm or indoxacarb 14.5% SC 200ml mixing in 200lit of water per acre crop.

**Animal husbandry:**

In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Give animals mineral mixture along with salt regularly and also wheat grains, jaggery etc. @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals. • In Poultry, keep the chicks warm by providing artificial light in the poultry sheds.

**Dr Sujata Sethy**  
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



**Debasish Jena**

**SMS (Agrometeorology)**

**Contact No-9861445080/7008453427**



**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**  
 (A Unit of ICAR-National Rice Research Institute)  
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



**Bulletin No.-547**  
**Date- 24/12/2024**

**AGROMET ADVISORY BULLETIN FOR NARASINGHPUR**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 05mm and district received average 06mm rainfall during the last week. The mean daily maximum and minimum temperature was 25.14°C (-3.47) and 17°C (+2.83), respectively. The mean relative humidity was 72-79%.

**Forecasted period: Today(8.30IST) to 29<sup>th</sup> December (8.30IST)2024**

**Weather advisory:** Moderate rainfall is forecasted for today and tomorrow and dry weather is forecasted for other 3 days in the block. The day temperature will be 21-30°C and night temperature will be 16°C for upcoming 5 days. Maximum and minimum relative humidity will be 63-93% and 36-67% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Narasinghpur (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	24/12/2024	25/12/2024	26/12/2024	27/12/2024	28/12/2024
Rainfall(mm)	19	15	00	00	00
T-Max(°C)	22	21	28	30	29
T-Min(°C)	16	16	16	16	16
Cloud Cover(octa)	7	4	0	0	0
RH Max (%)	92	93	63	74	70
RH Min (%)	54	67	37	36	38
Wind Speed (Kmph)	2	3	5	4	5
Wind Direction (deg)	45	330	135	146	135

**Agromet Advisory**

**General Advisory:** Stop irrigation, fertilizer and pesticide application on today and tomorrow to standing crops like Mustard, rabi groundnut and other rabi vegetables due to forecast of rain. After rain by utilization of rain water farmers should complete rabi oil seeds, pulses and vegetables.

**Pulses:** Forecasted Cloudy and foggy weather thereafter may aggravate the infestation of aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9% EC per litre water for controlling Powdery mildew.

**Rabi-Groundnut:** The early sown rabi groundnut crop is at 3 week stage. Go for weeding and hoeing. Apply 100 kg Gypsum by broadcasting or side dress 50 Kg of Gypsum. Irrigate the crop if required. To manage leaf miner spray Neem Based Pesticide (Azadirachtin) 300 PPM @ 1-litre /acre during primary stage of infestation. To manage chemically spray Profenophos 50% EC @ 400 ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 -litre of water.

**Mustard:** In view of past and forecasted weather condition farmers are advised to don't irrigate the mustard crop. Due to partly cloudy weather, there are chances of Aphid infestation in early sown mustard crop. To manage aphids at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water. There may be chances of white rust disease in Mustard, to control spray Metalaxyl 2g/liter of water.

**Vegetables:**

**Transplanting of mature seedlings of tomato, cauliflower, cabbage, knol khol and broccoli crop may be done in the raised bed.**

**Potato:** After 3 weeks of first earthing up go for hoeing and hand weeding and apply again 26 Kg of Urea and 20 Kg of MOP in line on both side the plants and earthing up should be done to cover up the tubers properly. Apply chlorpyrifos @ 2.5 ml/lit of water against the attack of cut worm and leaf eating insect. Due to forecasted rain and foggy weather in upcoming days there are chances of Leaf Blight disease incidence in potato crop. To manage leaf blight in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre.

**Tomato:** Stop irrigation activities as rainfall is forecasted. Due to rain and foggy weather in upcoming days, chances of leaf blight likely to happen in tomato. To avoid blight precaution must be taken during the initial infection stage. Spray 2.5-gram Metalaxyl + Mancozeb/lit of water. To manage Whitefly chemically sprays Thiamethoxam 25 % WG @ 40-gram/acre or Pyriproxifen 10% EC @ 200-ml/acre or Spiromesifen 22.9% SC @ 250-ml/acre by mixing in 200- litre of water. If Bacterial wilt will occur in brinjal and tomato. To control them, drench with copper oxychloride @ 3g+ plantomycin @1 g/ litre of water.

**Onion:** In present weather conditions, constant monitoring of timely sown onion crop against attack of thrips and infection of purple blotch should be done. Need based spraying of Diethane M -45 @ 3 grams per litre of water with sticky material are advised.

**Brinjal and Chilly:** Shoot and fruit borer and leaf eating caterpillar may observe in brinjal field, to control it spray Emamectin benzoate 5% SG @ 0.4 g/litre of water or Chlorantraniliprole 18.5% SC @ 0.4 ml/litre of water. Spraying should be done during the early morning or evening hours.

**Cabbage and Cauliflower:** Browning or Brown rot is a common physiological disorder in cauliflower crop due to deficiency of Boron. This may be controlled by application of borax @ 4-kg/ acre as soil application during transplanting. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Due to slight increase in temperature Cole crops may get affected by diamond back moth. To control this apply Spinosad 45SC @ 0.5ml/l. There is a chance of occurrence of thrip, mites and Aphids in cole crops, for control spray imidachloprid 0.3 ml /liter or fipronil 2 ml /liter of water. Leaf eating caterpillar and fruit borer may infest the crops. Spray fipronil 5% SL 400ml or emamectinbenzoate 5% SG 80gm or indoxacarb 14.5% SC 200ml mixing in 200lit of water per acre crop.

**Animal husbandry:**

- In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Give animals mineral mixture along with salt regularly and also wheat grains, jaggery etc. @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals.
- In Poultry, keep the chicks warm by providing artificial light in the poultry sheds.

**Dr Sujata Sethy**  
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



**Debasish Jena**

**SMS (Agrometeorology)**

**Contact No-9861445080/7008453427**



**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**  
 (A Unit of ICAR-National Rice Research Institute)  
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



**Bulletin No.-547**  
**Date- 24/12/2024**

**AGROMET ADVISORY BULLETIN FOR NIALI**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 09mm and district received average 06mm rainfall during the last week. The mean daily maximum and minimum temperature was 25.14°C (-3.47) and 17°C (+2.83), respectively. The mean relative humidity was 72-79%.

**Forecasted period: Today(8.30IST) to 29<sup>th</sup> December (8.30IST)2024**

**Weather advisory:** Moderate rainfall is forecasted for today and tomorrow and dry weather is forecasted for other 3 days in the block. The day temperature will be 21-30°C and night temperature will be 17-19°C for upcoming 5 days. Maximum and minimum relative humidity will be 57-84% and 35-62% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Niali (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	24/12/2024	25/12/2024	26/12/2024	27/12/2024	28/12/2024
Rainfall(mm)	20	16	00	00	00
T-Max(°C)	24	21	27	29	30
T-Min(°C)	18	17	17	19	18
Cloud Cover(octa)	7	6	2	0	0
RH Max (%)	84	84	66	62	57
RH Min (%)	62	61	56	39	35
Wind Speed(Kmph)	9	7	9	10	10
Wind Direction(deg)	13	21	32	45	45

**Agromet Advisory**

**General Advisory:** Stop irrigation, fertilizer and pesticide application on today and tomorrow to standing crops like Mustard, rabi groundnut and other rabi vegetables due to forecast of rain. After rain by utilization of rain water farmers should complete rabi oil seeds, pulses and vegetables.

**Pulses:** Forecasted Cloudy and foggy weather thereafter may aggravate the infestation of aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9% EC per litre water for controlling Powdery mildew.

**Rabi-Groundnut:** The early sown rabi groundnut crop is at 3 week stage. Go for weeding and hoeing. Apply 100 kg Gypsum by broadcasting or side dress 50 Kg of Gypsum. Irrigate the crop if required. To manage leaf miner spray Neem Based Pesticide (Azadirachtin) 300 PPM @ 1-litre /acre during primary stage of infestation. To manage chemically spray Profenophos 50% EC @ 400 ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 -litre of water.

**Mustard:** In view of past and forecasted weather condition farmers are advised to don't irrigate the mustard crop. Due to partly cloudy weather, there are chances of Aphid infestation in early sown mustard crop. To manage aphids at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water. There may be chances of white rust disease in Mustard, to control spray Metalaxyl 2g/liter of water.

**Vegetables:**

**Transplanting of mature seedlings of tomato, cauliflower, cabbage, knol khol and broccoli crop may be done in the raised bed.**

**Potato:** After 3 weeks of first earthing up go for hoeing and hand weeding and apply again 26 Kg of Urea and 20 Kg of MOP in line on both side the plants and earthing up should be done to cover up the tubers properly. Apply chlorpyrifos @ 2.5 ml/lit of water against the attack of cut worm and leaf eating insect. Due to forecasted rain and foggy weather in upcoming days there are chances of Leaf Blight disease incidence in potato crop. To manage leaf blight in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre.

**Tomato:** Stop irrigation activities as rainfall is forecasted. Due to rain and foggy weather in upcoming days, chances of leaf blight likely to happen in tomato. To avoid blight precaution must be taken during the initial infection stage. Spray 2.5-gram Metalaxyl + Mancozeb/lit of water To manage Whitefly chemically sprays Thiamethoxam 25 % WG @ 40-gram/acre or Pyriproxifen 10% EC @ 200-ml/acre or Spiromesifen 22.9% SC @ 250-ml/acre by mixing in 200- litre of water. If Bacterial wilt will occur in brinjal and tomato. To control them, drench with copper oxychloride @ 3g+ plantomycin @1 g/ litre of water.

**Onion:** In present weather conditions, constant monitoring of timely sown onion crop against attack of thrips and infection of purple blotch should be done. Need based spraying of Diethane M -45 @ 3 grams per litre of water with sticky material are advised.

**Brinjal and Chilly:** Shoot and fruit borer and leaf eating caterpillar may observe in brinjal field, to control it spray Emamectin benzoate 5% SG @ 0.4 g/litre of water or Chlorantranilprole 18.5% SC @ 0.4 ml/litre of water. Spraying should be done during the early morning or evening hours.

**Cabbage and Cauliflower:** Browning or Brown rot is a common physiological disorder in cauliflower crop due to deficiency of Boron. This may be controlled by application of borax @ 4-kg/ acre as soil application during transplanting. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Due to slight increase in temperature Cole crops may get affected by diamond back moth. To control this apply Spinosad 45SC @ 0.5ml/l. There is a chance of occurrence of thrip, mites and Aphids in cole crops, for control spray imidachloprid 0.3 ml /liter or fipronil 2 ml /liter of water. . Leaf eating caterpillar and fruit borer may infest the crops. Spray fipronil 5% SL 400ml or emamectinbenzoate 5% SG 80gm or indoxacarb 14.5% SC 200ml mixing in 200lit of water per acre crop.

**Animal husbandry:**

- In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Give animals mineral mixture along with salt regularly and also wheat grains, jaggery etc. @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals. • In Poultry, keep the chicks warm by providing artificial light in the poultry sheds.

**Dr Sujata Sethy**  
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



**Debasish Jena**

**SMS (Agrometeorology)**

**Contact No-9861445080/7008453427**



**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**  
 (A Unit of ICAR-National Rice Research Institute)  
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



**Bulletin No.-547**  
**Date- 24/12/2024**

**AGROMET ADVISORY BULLETIN FOR NISCHINTAKOILI**  
**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 07mm and district received average 06mm rainfall during the last week. The mean daily maximum and minimum temperature was 25.14°C (-3.47) and 17°C (+2.83), respectively. The mean relative humidity was 72-79%.

**Forecasted period: Today(8.30IST) to 29<sup>th</sup> December (8.30IST)2024**

**Weather advisory:** Moderate rainfall is forecasted for today and tomorrow and dry weather is forecasted for other 3 days in the block. The day temperature will be 22-30°C and night temperature will be 18-19°C for upcoming 5 days. Maximum and minimum relative humidity will be 48-82% and 32-62% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Nischintakoili (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	24/12/2024	25/12/2024	26/12/2024	27/12/2024	28/12/2024
Rainfall(mm)	17	14	00	00	00
T-Max(°C)	22	23	28	30	30
T-Min(°C)	19	18	18	18	18
Cloud Cover(octa)	8	5	1	0	0
RH Max (%)	82	76	63	63	48
RH Min (%)	62	55	43	37	32
Wind Speed (Kmph)	6	6	10	9	10
Wind Direction (deg)	360	23	25	23	28

**Agromet Advisory**

**General Advisory:** Stop irrigation, fertilizer and pesticide application on today and tomorrow to standing crops like Mustard, rabi groundnut and other rabi vegetables due to forecast of rain. After rain by utilization of rain water farmers should complete rabi oil seeds, pulses and vegetables.

**Pulses:** Forecasted Cloudy and foggy weather thereafter may aggravate the infestation of aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9% EC per litre water for controlling Powdery mildew.

**Rabi-Groundnut:** The early sown rabi groundnut crop is at 3 week stage. Go for weeding and hoeing. Apply 100 kg Gypsum by broadcasting or side dress 50 Kg of Gypsum. Irrigate the crop if required. To manage leaf miner spray Neem Based Pesticide (Azadirachtin) 300 PPM @ 1-litre /acre during primary stage of infestation. To manage chemically spray Profenophos 50% EC @ 400 ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 -litre of water.

**Mustard:** In view of past and forecasted weather condition farmers are advised to don't irrigate the mustard crop. Due to partly cloudy weather, there are chances of Aphid infestation in early sown mustard crop. To manage aphids at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water. There may be chances of white rust disease in Mustard, to control spray Metalaxyl 2g/liter of water.

**Vegetables:**

**Transplanting of mature seedlings of tomato, cauliflower, cabbage, knol khol and broccoli crop may be done in the raised bed.**

**Potato:** After 3 weeks of first earthing up go for hoeing and hand weeding and apply again 26 Kg of Urea and 20 Kg of MOP in line on both side the plants and earthing up should be done to cover up the tubers properly. Apply chlorpyrifos @ 2.5 ml/lit of water against the attack of cut worm and leaf eating insect. Due to forecasted rain and foggy weather in upcoming days there are chances of Leaf Blight disease incidence in potato crop. To manage leaf blight in potato, spray Mancozeb 75% WP @ 600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre.

**Tomato:** Stop irrigation activities as rainfall is forecasted. Due to rain and foggy weather in upcoming days, chances of leaf blight likely to happen in tomato. To avoid blight precaution must be taken during the initial infection stage. Spray 2.5-gram Metalaxyl + Mancozeb/lit of water. To manage Whitefly chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Pyriproxifen 10% EC @ 200-ml/acre or Spiromesifen 22.9% SC @ 250-ml/acre by mixing in 200- litre of water. If Bacterial wilt will occur in brinjal and tomato. To control them, drench with copper oxychloride @ 3g+ plantomycin @1 g/ litre of water.

**Onion:** In present weather conditions, constant monitoring of timely sown onion crop against attack of thrips and infection of purple blotch should be done. Need based spraying of Diethane M -45 @ 3 grams per litre of water with sticky material are advised.

**Brinjal and Chilly:** Shoot and fruit borer and leaf eating caterpillar may observe in brinjal field, to control it spray Emamectin benzoate 5% SG @ 0.4 g/litre of water or Chlorantraniliprole 18.5% SC @ 0.4 ml/litre of water. Spraying should be done during the early morning or evening hours.

**Cabbage and Cauliflower:** Browning or Brown rot is a common physiological disorder in cauliflower crop due to deficiency of Boron. This may be controlled by application of borax @ 4-kg/ acre as soil application during transplanting. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Due to slight increase in temperature Cole crops may get affected by diamond back moth. To control this apply Spinosad 45SC @ 0.5ml/l. There is a chance of occurrence of thrip, mites and Aphids in cole crops, for control spray imidachloprid 0.3 ml /liter or fipronil 2 ml /liter of water. Leaf eating caterpillar and fruit borer may infest the crops. Spray fipronil 5% SL 400ml or emamectinbenzoate 5% SG 80gm or indoxacarb 14.5% SC 200ml mixing in 200lit of water per acre crop.

**Animal husbandry:**

- In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Give animals mineral mixture along with salt regularly and also wheat grains, jaggery etc. @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals.
- In Poultry, keep the chicks warm by providing artificial light in the poultry sheds.

**Dr Sujata Sethy**  
 (Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



**Debasish Jena**

**SMS (Agrometeorology)**

**Contact No-9861445080/7008453427**



**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**  
(A Unit of ICAR-National Rice Research Institute)  
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



**Bulletin No.-547**  
**Date- 24/12/2024**

**AGROMET ADVISORY BULLETIN FOR SALEPUR**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 06mm and district received average 06mm rainfall during the last week. The mean daily maximum and minimum temperature was 25.14°C (-3.47) and 17°C (+2.83), respectively. The mean relative humidity was 72-79%.

**Forecasted period: Today(8.30IST) to 29<sup>th</sup> December (8.30IST)2024**

**Weather advisory:** Moderate rainfall is forecasted for today and tomorrow and dry weather is forecasted for other 3 days in the block. The day temperature will be 22-30°C and night temperature will be 18-19°C for upcoming 5 days. Maximum and minimum relative humidity will be 48-82% and 32-61% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Salepur (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	24/12/2024	25/12/2024	26/12/2024	27/12/2024	28/12/2024
Rainfall(mm)	17	14	00	00	00
T-Max(°C)	22	24	28	30	30
T-Min(°C)	19	18	18	18	18
Cloud Cover(octa)	8	4	1	0	0
RH Max (%)	82	77	63	63	48
RH Min (%)	61	55	42	37	32
Wind Speed (Kmph)	6	6	9	9	9
Wind Direction (deg)	360	18	25	23	36

**Agromet Advisory**

**General Advisory:** Stop irrigation, fertilizer and pesticide application on today and tomorrow to standing crops like Mustard, rabi groundnut and other rabi vegetables due to forecast of rain. After rain by utilization of rain water farmers should complete rabi oil seeds, pulses and vegetables.

**Pulses:** Forecasted Cloudy and foggy weather thereafter may aggravate the infestation of aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9% EC per litre water for controlling Powdery mildew.

**Rabi-Groundnut:** The early sown rabi groundnut crop is at 3 week stage. Go for weeding and hoeing. Apply 100 kg Gypsum by broadcasting or side dress 50 Kg of Gypsum. Irrigate the crop if required. To manage leaf miner spray Neem Based Pesticide (Azadirachtin) 300 PPM @ 1-litre /acre during primary stage of infestation. To manage chemically spray Profenophos 50% EC @ 400 ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 -litre of water.

**Mustard:** In view of past and forecasted weather condition farmers are advised to don't irrigate the mustard crop. Due to partly cloudy weather, there are chances of Aphid infestation in early sown mustard crop. To manage aphids at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water. There may be chances of white rust disease in Mustard, to control spray Metalaxyl 2g/liter of water.

**Vegetables:**

**Transplanting of mature seedlings of tomato, cauliflower, cabbage, knol khol and broccoli crop may be done in the raised bed.**

**Potato:** After 3 weeks of first earthing up go for hoeing and hand weeding and apply again 26 Kg of Urea and 20 Kg of MOP in line on both side the plants and earthing up should be done to cover up the tubers properly. Apply chlorpyrifos @ 2.5 ml/lit of water against the attack of cut worm and leaf eating insect. Due to forecasted rain and foggy weather in upcoming days there are chances of Leaf Blight disease incidence in potato crop. To manage leaf blight in potato, spray Mancozeb 75% WP @ 600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre.

**Tomato:** Stop irrigation activities as rainfall is forecasted. Due to rain and foggy weather in upcoming days, chances of leaf blight likely to happen in tomato. To avoid blight precaution must be taken during the initial infection stage. Spray 2.5-gram Metalaxyl + Mancozeb/lit of water. To manage Whitefly chemically sprays Thiamethoxam 25 % WG @ 40-gram/acre or Pyriproxifen 10% EC @ 200-ml/acre or Spiromesifen 22.9% SC @ 250-ml/acre by mixing in 200- litre of water. If Bacterial wilt will occur in brinjal and tomato. To control them, drench with copper oxychloride @ 3g+ plantomycin @ 1 g/ litre of water.

**Onion:** In present weather conditions, constant monitoring of timely sown onion crop against attack of thrips and infection of purple blotch should be done. Need based spraying of Diethane M -45 @ 3 grams per litre of water with sticky material are advised.

**Brinjal and Chilly:** Shoot and fruit borer and leaf eating caterpillar may observe in brinjal field, to control it spray Emamectin benzoate 5% SG @ 0.4 g/litre of water or Chlorantraniliprole 18.5% SC @ 0.4 ml/litre of water. Spraying should be done during the early morning or evening hours.

**Cabbage and Cauliflower:** Browning or Brown rot is a common physiological disorder in cauliflower crop due to deficiency of Boron. This may be controlled by application of borax @ 4-kg/ acre as soil application during transplanting. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Due to slight increase in temperature Cole crops may get affected by diamond back moth. To control this apply Spinosad 45SC @ 0.5ml/l. There is a chance of occurrence of thrip, mites and Aphids in cole crops, for control spray imidachloprid 0.3 ml /liter or fipronil 2 ml /liter of water. Leaf eating caterpillar and fruit borer may infest the crops. Spray fipronil 5% SL 400ml or emamectinbenzoate 5% SG 80gm or indoxacarb 14.5% SC 200ml mixing in 200lit of water per acre crop.

**Animal husbandry:**

In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Give animals mineral mixture along with salt regularly and also wheat grains, jaggery etc. @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals. • In Poultry, keep the chicks warm by providing artificial light in the poultry sheds.

**Dr Sujata Sethy**  
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



**Debasish Jena**  
**SMS (Agrometeorology)**  
**Contact No-9861445080/7008453427**



**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**



(A Unit of ICAR-National Rice Research Institute)  
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com

Bulletin No.-547  
 Date- 24/12/2024

**AGROMET ADVISORY BULLETIN FOR TANGI-CHOUDWAR**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 04mm and district received average 06mm rainfall during the last week. The mean daily maximum and minimum temperature was 25.14°C (-3.47) and 17°C (+2.83), respectively. The mean relative humidity was 72-79%.

**Forecasted period: Today(8.30IST) to 29<sup>th</sup> December (8.30IST)2024**

**Weather advisory: Moderate rainfall is forecasted for today and tomorrow and dry weather is forecasted for other 3 days in the block.** The day temperature will be 22-30°C and night temperature will be 18-19°C for upcoming 5 days. Maximum and minimum relative humidity will be 52-79% and 38-61% respectively. **There will be partly cloudy to overcast sky for next 5 days.**

Block	Tangi-Choudwar (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	24/12/2024	25/12/2024	26/12/2024	27/12/2024	28/12/2024
Rainfall(mm)	16	13	00	00	00
T-Max(°C)	22	24	28	30	30
T-Min(°C)	19	18	18	18	18
Cloud Cover(octa)	7	4	1	0	0
RH Max (%)	79	72	67	67	52
RH Min (%)	61	57	47	46	38
Wind Speed (Kmph)	11	10	14	13	13
Wind Direction (deg)	21	21	32	45	45

**Agromet Advisory**

**General Advisory:** Stop irrigation, fertilizer and pesticide application on today and tomorrow to standing crops like Mustard, rabi groundnut and other rabi vegetables due to forecast of rain. After rain by utilization of rain water farmers should complete rabi oil seeds, pulses and vegetables.

**Pulses:** Forecasted Cloudy and foggy weather thereafter may aggravate the infestation of aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and spray 3 g Sulfex or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9% EC per litre water for controlling Powdery mildew.

**Rabi-Groundnut:** The early sown rabi groundnut crop is at 3 week stage. Go for weeding and hoeing. Apply 100 kg Gypsum by broadcasting or side dress 50 Kg of Gypsum. Irrigate the crop if required. To manage leaf miner spray Neem Based Pesticide (Azadirachtin) 300 PPM @ 1-litre /acre during primary stage of infestation. To manage chemically spray Profenophos 50% EC @ 400 ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 -litre of water.

**Mustard:** In view of past and forecasted weather condition farmers are advised to don't irrigate the mustard crop. Due to partly cloudy weather, there are chances of Aphid infestation in early sown mustard crop. To manage aphids at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water. There may be chances of white rust disease in Mustard, to control spray Metalaxyl 2g/liter of water.

**Vegetables:**

**Transplanting of mature seedlings of tomato, cauliflower, cabbage, knol khol and broccoli crop may be done in the raised bed.**

**Potato:** After 3 weeks of first earthing up go for hoeing and hand weeding and apply again 26 Kg of Urea and 20 Kg of MOP in line on both side the plants and earthing up should be done to cover up the tubers properly. Apply chlorpyrifos @ 2.5 ml/lit of water against the attack of cut worm and leaf eating insect. Due to forecasted rain and foggy weather in upcoming days there are chances of Leaf Blight disease incidence in potato crop. To manage leaf blight in potato, spray Mancozeb 75% WP @ 600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre.

**Tomato:** Stop irrigation activities as rainfall is forecasted. Due to rain and foggy weather in upcoming days, chances of leaf blight likely to happen in tomato. To avoid blight precaution must be taken during the initial infection stage. Spray 2.5-gram Metalaxyl + Mancozeb/lit of water. To manage Whitefly chemically sprays Thiamethoxam 25 % WG @ 40-gram/acre or Pyriproxifen 10% EC @ 200-ml/acre or Spiromesifen 22.9% SC @ 250-ml/acre by mixing in 200- litre of water. If Bacterial wilt will occur in brinjal and tomato. To control them, drench with copper oxychloride @ 3g+ plantomycin @ 1 g/ litre of water.

**Onion:** In present weather conditions, constant monitoring of timely sown onion crop against attack of thrips and infection of purple blotch should be done. Need based spraying of Diethane M -45 @ 3 grams per litre of water with sticky material are advised.

**Brinjal and Chilly:** Shoot and fruit borer and leaf eating caterpillar may observe in brinjal field, to control it spray Emamectin benzoate 5% SG @ 0.4 g/litre of water or Chlorantraniliprole 18.5% SC @ 0.4 ml/litre of water. Spraying should be done during the early morning or evening hours.

**Cabbage and Cauliflower:** Browning or Brown rot is a common physiological disorder in cauliflower crop due to deficiency of Boron. This may be controlled by application of borax @ 4-kg/ acre as soil application during transplanting. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Due to slight increase in temperature Cole crops may get affected by diamond back moth. To control this apply Spinosad 45SC @ 0.5ml/l. There is a chance of occurrence of thrip, mites and Aphids in cole crops, for control spray imidachloprid 0.3 ml /liter or fipronil 2 ml /liter of water. Leaf eating caterpillar and fruit borer may infest the crops. Spray fipronil 5% SL 400ml or emamectinbenzoate 5% SG 80gm or indoxacarb 14.5% SC 200ml mixing in 200lit of water per acre crop.

**Animal husbandry:**

In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Give animals mineral mixture along with salt regularly and also wheat grains, jaggery etc. @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals. • In Poultry, keep the chicks warm by providing artificial light in the poultry sheds.

Dr Sujata Sethy  
 (Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



Debasish Jena  
 SMS (Agrometeorology)  
 Contact No-9861445080/7008453427





**GRAMIN KRISHI APP MAASAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**  
 (A Unit of ICAR-National Rice Research Institute)  
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



**Bulletin No.-547**  
**Date- 24/12/2024**

**AGROMET ADVISORY BULLETIN FOR TIGIRIA**

**District Past week average weather condition and Block past week average Rainfall condition**

The block received average 01mm and district received average 06mm rainfall during the last week. The mean daily maximum and minimum temperature was 25.14°C (-3.47) and 17°C (+2.83), respectively. The mean relative humidity was 72-79%.

**Forecasted period: Today(8.30IST) to 29<sup>th</sup> December (8.30IST)2024**

**Weather advisory:** Moderate rainfall is forecasted for today and tomorrow and dry weather is forecasted for other 3 days in the block. The day temperature will be 23-30°C and night temperature will be 17-18°C for upcoming 5 days. Maximum and minimum relative humidity will be 57-87% and 35-63% respectively. There will be partly cloudy to overcast sky for next 5 days.

Block	Tigiria (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	24/12/2024	25/12/2024	26/12/2024	27/12/2024	28/12/2024
Rainfall(mm)	17	13	00	00	00
T-Max(°C)	23	24	29	30	29
T-Min(°C)	18	17	17	18	17
Cloud Cover(octa)	8	5	0	0	0
RH Max (%)	87	83	63	67	57
RH Min (%)	54	63	35	36	35
Wind Speed (Kmph)	3	4	7	6	5
Wind Direction (deg)	68	150	90	68	90

**Agro met Advisory**

**General Advisory:** Stop irrigation, fertilizer and pesticide application on today and tomorrow to standing crops like Mustard, rabi groundnut and other rabi vegetables due to forecast of rain. After rain by utilization of rain water farmers should complete rabi oil seeds, pulses and vegetables.

**Pulses:** Forecasted Cloudy and foggy weather thereafter may aggravate the infestation of aphid and powdery mildew in pulses. Spray 2 ml. Dimethoate or Metacystox per litre water to control aphid and spray 3 g Sulfox or 1 ml Propiconazole 10% EC or 1 ml Tebiconazole 25.9% EC per litre water for controlling Powdery mildew.

**Rabi-Groundnut:** The early sown rabi groundnut crop is at 3 week stage. Go for weeding and hoeing. Apply 100 kg Gypsum by broadcasting or side dress 50 Kg of Gypsum. Irrigate the crop if required. To manage leaf miner spray Neem Based Pesticide (Azadirachtin) 300 PPM @ 1-litre /acre during primary stage of infestation. To manage chemically spray Profenophos 50% EC @ 400 ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 -litre of water.

**Mustard:** In view of past and forecasted weather condition farmers are advised to don't irrigate the mustard crop. Due to partly cloudy weather, there are chances of Aphid infestation in early sown mustard crop. To manage aphids at early stage, put 8 number of yellow sticky traps per acre and to manage Aphids chemically spray Thiamethoxam 25 % WG @ 40-gram/acre or Acetamiprid 20 % SP @ 50-gram/acre in 200-liter of water. To manage Mustard sawfly spray Ethofenprox 10 % EC @ 200-ml/acre or Chlorpyrifos 50% + Cypermethrin 5 % E.C @ 400 ml/acre by mixing it in 200 litre of water. There may be chances of white rust disease in Mustard, to control spray Metalaxyl 2g/litre of water.

**Vegetables:** Transplanting of mature seedlings of tomato, cauliflower, cabbage, knol khol and broccoli crop may be done in the raised bed.

**Potato:** After 3 weeks of first earthing up go for hoeing and hand weeding and apply again 26 Kg of Urea and 20 Kg of MOP in line on both side the plants and earthing up should be done to cover up the tubers properly. Apply chlorpyrifos @ 2.5 ml/lit of water against the attack of cut worm and leaf eating insect. Due to forecasted rain and foggy weather in upcoming days there are chances of Leaf Blight disease incidence in potato crop. To manage leaf blight in potato, spray Mancozeb 75% WP @600 gram/acre or Carbendazim 12% + Mancozeb 63% W.P @ 400-gram/acre.

**Tomato:** Stop irrigation activities as rainfall is forecasted. Due to rain and foggy weather in upcoming days, chances of leaf blight likely to happen in tomato. To avoid blight precaution must be taken during the initial infection stage. Spray 2.5-gram Metalaxyl + Mancozeb/lit of water To manage Whitefly chemically sprays Thiamethoxam 25 % WG @ 40-gram/acre or Pyriproxifen 10% EC @ 200-ml/acre or Spiromesifen 22.9% SC @ 250-ml/acre by mixing in 200- litre of water. If Bacterial wilt will occur in brinjal and tomato. To control them, drench with copper oxychloride @ 3g+ plantomycin @1 g/ litre of water.

**Onion:** In present weather conditions, constant monitoring of timely sown onion crop against attack of thrips and infection of purple blotch should be done. Need based spraying of Diethane M -45 @ 3 grams per litre of water with sticky material are advised.

**Brinjal and Chilly:** Shoot and fruit borer and leaf eating caterpillar may observe in brinjal field, to control it spray Emamectin benzoate 5% SG @ 0.4 g/litre of water or Chlorantranilprole 18.5% SC @ 0.4 ml/litre of water. Spraying should be done during the early morning or evening hours.

**Cabbage and Cauliflower:** Browning or Brown rot is a common physiological disorder in cauliflower crop due to deficiency of Boron. This may be controlled by application of borax @ 4-kg/ acre as soil application during transplanting. At three week stage go for hoeing, weeding and apply 55kg urea per acre followed by irrigation. Due to slight increase in temperature Cole crops may get affected by diamond back moth. To control this apply Spinosad 45SC @ 0.5ml/l. There is a chance of occurrence of thrip, mites and Aphids in cole crops, for control spray imidachloprid 0.3 ml /liter or fipronil 2 ml /liter of water. Leaf eating caterpillar and fruit borer may infest the crops. Spray fipronil 5% SL 400ml or emamectinbenzoate 5% SG 80gm or indoxacarb 14.5% SC 200ml mixing in 200lit of water per acre crop.

**Animal husbandry:**

In winter during cold hours farmers should keep doors, windows and open spaces in sheds covered with polythene or gunny bag curtains to prevent direct contact to cold air. Give animals mineral mixture along with salt regularly and also wheat grains, jaggery etc. @ 10%-20% in the daily ration during winter season to meet the energy requirement of the animals. • In Poultry, keep the chicks warm by providing artificial light in the poultry sheds.

**Dr Sujata Sethy**  
(Nodal officer)



Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area



**Debasish Jena**

**SMS (Agrometeorology)**


**Contact No-9861445080/7008453427**





**GRAMIN KRISHI MAUSAM SEWA**  
**INDIA METEOROLOGICAL DEPARTMENT**  
**DISTRICT AGROMET UNIT**  
**KRISHI VIGYAN KENDRA CUTTACK**  
 (A Unit of ICAR-National Rice Research Institute)  
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-547  
Date- 24/12/2024

				
MEGHDOOT APP	DAMINI APP	FACEBOOK (Damu Cuttack)	BLOCK SPECIFIC WHAT'S APP GROUPS	@DAMUCUTTACK

- Kindly install and use **“Meghdoot mobile App”** for Cuttack district weather forecast, warnings and weather based agromet advisory services.  
<https://play.google.com/store/apps/details?id=com.aas.meghdoot>
- Kindly install and use **“Damini mobile App”** for location specific for lightening forecast and warnings.  
<https://play.google.com/store/apps/details?id=com.lightening.live.damini>
- Follow the **block specific whats app groups** and join in your block wise groups for getting weather based agromet advisory services twice in a week (Tuesday and Friday) and other weather warning based farm management advisories.
- Follow the **Official Website links** to get the weather based agromet advisory service bulletins (**English & Odia**) twice in a week (Tuesday and Friday).

1	ICAR-NRRI, Cuttack, Odisha	<a href="https://icar-nrri.in/agromate/">https://icar-nrri.in/agromate/</a>	
2	Agrimet division, IMD, Pune	<a href="https://imdagrmet.gov.in/imdproject/AGIndex.php">https://imdagrmet.gov.in/imdproject/AGIndex.php</a>	

Dr Sujata Sethy  
(Nodal officer)



**Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area**



Debasish Jena  
SMS (Agrometeorology)  
Contact No-9861445080/7008453427