



ICAR – Central Research Institute (An
ISO 9001: 2015 Certified Institute)
Cuttack-753 006, Odisha, India



Agro-Advisory Service

Strategies for First Fortnight of February 2025

1. Winter Rice

- ❖ Farmers are advised not to burn the rice straw in the field.
- ❖ Paddy grains after threshing need to be sun dried to 14% moisture for consumption purpose and for seed purpose it should be dried to 12% moisture for better shelf life. Pack grains of each variety separately without mixing for better price of the produce.
- ❖ Use 'Super Grain Bag' for safe storage of paddy/rice for longer period of time **or** store the harvested paddy in properly bagged and stacked with suitable cover.
- ❖ Soon after noticing the infestation of the stored grain insect pest, take up fumigation by using Aluminium phosphide (do not use in dwelling houses) tablets @ of 3 tablets/ ton grain (total 9 gm of tablets) in fairly air tight containers **or** by covering with thick tarpaulin leaving no gaps. The tablets should be wrapped in cotton pouches before placing them in the stacks. All the corners of plastic cover should be plastered with 6 inch thick layer of mud/ adhesive tapes to prevent leakage of gas. Minimum exposure period is for about 7-10 days.

2. Summer rice

2.1 Transplanted rice

- ❖ In nursery, where stem borer infestation is expected to notice, installation of pheromone traps with Scirpholure is recommended (at least 3 per nursery of 200 m²). When the number of male moths/trap reaches 4 or 5, spray Azadirachtin 0.15% neem seed kernel based EC formulation @ 800ml/acre **or** broadcast granular insecticide

Chlorantraniliprole 4% GR @ 4 kg/acre or Cartaphydrochloride 4G @ 10 kg/acre mixing with sand at 1:1 ratio or spray Chlorantraniliprole 18.5% SC @ 60 ml/acre in 200 litre of water.

- ❖ If infestation of Thrips is noticed in rice nursery, spray Azadirachtin 0.15% neem seed kernel based EC formulation @ 800 ml/acre or Lambda-cyhalothrin 5 % EC @ 200 ml/acre or Thiamethoxam 25 % WG @ 40g /acre.
- ❖ In-case of Leaf blast infestation, spray Tebuconazole 50% + Trifloxystrobin 25% WG @ 80 g/acre or Isoprothiolane 40EC @ 300 ml/acre in 200 litres of water. Repeat the spray at 7-10 days' interval.
- ❖ If infestation of seedling blight is noticed in rice nursery, spray Carbendazim @ 400 g/acre or Propiconazole @ 200 g/acre in 200 litre of water.
- ❖ Complete the transplanting of 3-5 week old seedlings in well puddled soil at spacing of 15cm x 15cm with 3-4 seedlings per hill at shallow depth.
- ❖ In bacterial blight endemic areas, dip these seedling roots in 0.1% (1g in one litre of water) Plantomycin solution before transplanting for 30 minutes.
- ❖ Apply (DAP 44kg + MOP 22kg) or (Urea 22kg + SSP 125kg + MOP 22 kg) as basal dose during final puddling.
- ❖ Farmers those have already completed transplanting, apply granular herbicide Bensulfuron-methyl + Pretilachlor @ 4kg/ acre at 5-8 days after transplanting. Mix the granular herbicide with 4 kg of sand/ acre and broadcast it uniformly in the field or spray Pyrazosulfuron-ethyl 10 WP @ 80g/acre in 140 litres of water at 3-5 days after transplanting or spray Bispyribac sodium @ 120ml/acre at 10-15 DAT or 2-3 leaf stage of weeds in 140 litres of water.
- ❖ In just transplanted field put 3 pheromone traps with 5mg lure/acre in the rice field for monitoring of the yellow stem borer and leaf folder. Whenever the number of male moths/trap reaches 4 or 5, spray Azadirachtin 0.15% neem seed kernel based EC formulation @ 800 ml/acre or broadcast granular insecticide Chlorantraniliprole 4% GR @ 4 kg/acre or Cartap hydrochloride 4G @ 10 kg/acre mixing with sand at 1:1 ratio or spray Chlorantraniliprole 18.5% SC @ 60 ml/acre in 200 litre of water.

- ❖ If infestation of Bakanae disease is noticed, spray Carbendazim 50WP @ 200 g per acre in 200 litre of water and repeat the spray at 7-10 days' interval.
- ❖ In established transplanted rice field apply urea @ 35 kg/acre as top dressing at 20-25 DAT (maximum tillering stage)

2.2 Wet direct seeded rice

- ❖ If pre-emergence herbicides were not applied, to control weeds apply early post-emergent ready mix Bensulfuron-methyl+ Pretilachlor granular herbicide @ 4 kg/acre mixed with 4 kg of sand at 5-10 DAS or spray Bispyribac sodium @ 120 ml/acre at 10-12 DAS or 2-3 leaf stage of weeds or, Penoxsulum 1.02% + Cyhalofop-butyl 5.1% @ 800ml/acre at 15-20 DAS in 140 litres of water.
- ❖ Topdress 24 kg of urea per acre after weeding at tillering stage (20-25 DAS).
- ❖ Monitoring should continue for stem borer. Whenever the number of male moths/trap reaches 4 or 5 spray Azadiractin 0.15% neem seed kernel based EC formulation @ 800 ml/acre or Chlorantraniliprole 4% GR @ 4 kg/acre may be applied mixing with sand at 1:1 ratio or spray Chlorantraniliprole 18.5% SC @ 60 ml/acre in 200 litres of water.
- ❖ If infestation of blast disease is noticed, spray Carbendazim 50WP @ 200g/acre in 200 litres of water or Trifloxystrobin 25% + Tebuconazole 50% @ 80g per acre in 200 litres of water or Edifenphos 50EC @ 200 ml/acre in 200 litres of water.
- ❖ Farmers are advised to download and use NRR I developed **riceXpert** mobile App (available in Google Play store) for getting information on all aspects of rice cultivation
