




Cuttack District Block Wise Agro-met Advisory Service Bulletin

*Tuesday, 25th February, 2025
 (For the period 25th February 01st
 March, 2025)*





*Issued by
 District Agromet Unit,
 KrishiVigyan Kendra Cuttack*





STANDARD PRECIPITATION INDEX AND NDVI MAP

Standard Precipitation Index (SPI)	Normalized difference vegetation Index (NDVI)
<p style="text-align: center; font-size: small;">MINISTRY OF EARTH SCIENCES INDIA METEOROLOGICAL DEPARTMENT HYDROMET SECTION, CRS PUNE</p>  <p style="text-align: center; font-size: x-small;">SPI VALUES</p> <ul style="list-style-type: none"> 0.0 to 0.99 1.0 to 1.99 (Moderately Dry) 0% 0.00 - 0% Above 2.00 to 2.99 (Moderately Dry) 0% 0.00 - 0% Above 3.00 to 3.99 (Moderately Dry) 0% 0.00 - 0% Above 4.00 to 4.99 (Moderately Dry) 0% 0.00 - 0% Above 5.00 to 5.99 (Moderately Dry) 0% 0.00 - 0% Above 6.00 to 6.99 (Moderately Dry) 0% 0.00 - 0% Above 7.00 to 7.99 (Moderately Dry) 0% 0.00 - 0% Above 8.00 to 8.99 (Moderately Dry) 0% 0.00 - 0% Above 9.00 to 9.99 (Moderately Dry) 0% 0.00 - 0% Above 10.00 to 10.99 (Moderately Dry) 0% 0.00 - 0% Above 11.00 to 11.99 (Moderately Dry) 0% 0.00 - 0% Above 12.00 to 12.99 (Moderately Dry) 0% 0.00 - 0% Above 13.00 to 13.99 (Moderately Dry) 0% 0.00 - 0% Above 14.00 to 14.99 (Moderately Dry) 0% 0.00 - 0% Above 15.00 to 15.99 (Moderately Dry) 0% 0.00 - 0% Above 16.00 to 16.99 (Moderately Dry) 0% 0.00 - 0% Above 17.00 to 17.99 (Moderately Dry) 0% 0.00 - 0% Above 18.00 to 18.99 (Moderately Dry) 0% 0.00 - 0% Above 19.00 to 19.99 (Moderately Dry) 0% 0.00 - 0% Above 20.00 to 20.99 (Moderately Dry) 0% 0.00 - 0% Above 21.00 to 21.99 (Moderately Dry) 0% 0.00 - 0% Above 22.00 to 22.99 (Moderately Dry) 0% 0.00 - 0% Above 23.00 to 23.99 (Moderately Dry) 0% 0.00 - 0% Above 24.00 to 24.99 (Moderately Dry) 0% 0.00 - 0% Above 25.00 to 25.99 (Moderately Dry) 0% 0.00 - 0% Above 26.00 to 26.99 (Moderately Dry) 0% 0.00 - 0% Above 27.00 to 27.99 (Moderately Dry) 0% 0.00 - 0% Above 28.00 to 28.99 (Moderately Dry) 0% 0.00 - 0% Above 29.00 to 29.99 (Moderately Dry) 0% 0.00 - 0% Above 30.00 to 30.99 (Moderately Dry) 0% 0.00 - 0% Above 31.00 to 31.99 (Moderately Dry) 0% 0.00 - 0% Above 32.00 to 32.99 (Moderately Dry) 0% 0.00 - 0% Above 33.00 to 33.99 (Moderately Dry) 0% 0.00 - 0% Above 34.00 to 34.99 (Moderately Dry) 0% 0.00 - 0% Above 35.00 to 35.99 (Moderately Dry) 0% 0.00 - 0% Above 36.00 to 36.99 (Moderately Dry) 0% 0.00 - 0% Above 37.00 to 37.99 (Moderately Dry) 0% 0.00 - 0% Above 38.00 to 38.99 (Moderately Dry) 0% 0.00 - 0% Above 39.00 to 39.99 (Moderately Dry) 0% 0.00 - 0% Above 40.00 to 40.99 (Moderately Dry) 0% 0.00 - 0% Above 41.00 to 41.99 (Moderately Dry) 0% 0.00 - 0% Above 42.00 to 42.99 (Moderately Dry) 0% 0.00 - 0% Above 43.00 to 43.99 (Moderately Dry) 0% 0.00 - 0% Above 44.00 to 44.99 (Moderately Dry) 0% 0.00 - 0% Above 45.00 to 45.99 (Moderately Dry) 0% 0.00 - 0% Above 46.00 to 46.99 (Moderately Dry) 0% 0.00 - 0% Above 47.00 to 47.99 (Moderately Dry) 0% 0.00 - 0% Above 48.00 to 48.99 (Moderately Dry) 0% 0.00 - 0% Above 49.00 to 49.99 (Moderately Dry) 0% 0.00 - 0% Above 50.00 to 50.99 (Moderately Dry) 0% 0.00 - 0% Above 51.00 to 51.99 (Moderately Dry) 0% 0.00 - 0% Above 52.00 to 52.99 (Moderately Dry) 0% 0.00 - 0% Above 53.00 to 53.99 (Moderately Dry) 0% 0.00 - 0% Above 54.00 to 54.99 (Moderately Dry) 0% 0.00 - 0% Above 55.00 to 55.99 (Moderately Dry) 0% 0.00 - 0% Above 56.00 to 56.99 (Moderately Dry) 0% 0.00 - 0% Above 57.00 to 57.99 (Moderately Dry) 0% 0.00 - 0% Above 58.00 to 58.99 (Moderately Dry) 0% 0.00 - 0% Above 59.00 to 59.99 (Moderately Dry) 0% 0.00 - 0% Above 60.00 to 60.99 (Moderately Dry) 0% 0.00 - 0% Above 61.00 to 61.99 (Moderately Dry) 0% 0.00 - 0% Above 62.00 to 62.99 (Moderately Dry) 0% 0.00 - 0% Above 63.00 to 63.99 (Moderately Dry) 0% 0.00 - 0% Above 64.00 to 64.99 (Moderately Dry) 0% 0.00 - 0% Above 65.00 to 65.99 (Moderately Dry) 0% 0.00 - 0% Above 66.00 to 66.99 (Moderately Dry) 0% 0.00 - 0% Above 67.00 to 67.99 (Moderately Dry) 0% 0.00 - 0% Above 68.00 to 68.99 (Moderately Dry) 0% 0.00 - 0% Above 69.00 to 69.99 (Moderately Dry) 0% 0.00 - 0% Above 70.00 to 70.99 (Moderately Dry) 0% 0.00 - 0% Above 71.00 to 71.99 (Moderately Dry) 0% 0.00 - 0% Above 72.00 to 72.99 (Moderately Dry) 0% 0.00 - 0% Above 73.00 to 73.99 (Moderately Dry) 0% 0.00 - 0% Above 74.00 to 74.99 (Moderately Dry) 0% 0.00 - 0% Above 75.00 to 75.99 (Moderately Dry) 0% 0.00 - 0% Above 76.00 to 76.99 (Moderately Dry) 0% 0.00 - 0% Above 77.00 to 77.99 (Moderately Dry) 0% 0.00 - 0% Above 78.00 to 78.99 (Moderately Dry) 0% 0.00 - 0% Above 79.00 to 79.99 (Moderately Dry) 0% 0.00 - 0% Above 80.00 to 80.99 (Moderately Dry) 0% 0.00 - 0% Above 81.00 to 81.99 (Moderately Dry) 0% 0.00 - 0% Above 82.00 to 82.99 (Moderately Dry) 0% 0.00 - 0% Above 83.00 to 83.99 (Moderately Dry) 0% 0.00 - 0% Above 84.00 to 84.99 (Moderately Dry) 0% 0.00 - 0% Above 85.00 to 85.99 (Moderately Dry) 0% 0.00 - 0% Above 86.00 to 86.99 (Moderately Dry) 0% 0.00 - 0% Above 87.00 to 87.99 (Moderately Dry) 0% 0.00 - 0% Above 88.00 to 88.99 (Moderately Dry) 0% 0.00 - 0% Above 89.00 to 89.99 (Moderately Dry) 0% 0.00 - 0% Above 90.00 to 90.99 (Moderately Dry) 0% 0.00 - 0% Above 91.00 to 91.99 (Moderately Dry) 0% 0.00 - 0% Above 92.00 to 92.99 (Moderately Dry) 0% 0.00 - 0% Above 93.00 to 93.99 (Moderately Dry) 0% 0.00 - 0% Above 94.00 to 94.99 (Moderately Dry) 0% 0.00 - 0% Above 95.00 to 95.99 (Moderately Dry) 0% 0.00 - 0% Above 96.00 to 96.99 (Moderately Dry) 0% 0.00 - 0% Above 97.00 to 97.99 (Moderately Dry) 0% 0.00 - 0% Above 98.00 to 98.99 (Moderately Dry) 0% 0.00 - 0% Above 99.00 to 99.99 (Moderately Dry) 0% 0.00 - 0% Above 100.00 to 100.99 (Moderately Dry) 0% 0.00 - 0% Above 	<p style="text-align: center;">NOAA/VIIRS/BLENDED NDVI Composite ending on Week no 7 (12.02.2025 to 18.02.2025) over Agricultural region</p> <p style="text-align: center; font-size: large;">Odisha</p>  <ul style="list-style-type: none"> -0.9999 - 0 0 - 0.05 0.05 - 0.1 0.1 - 0.15 0.15 - 0.2 0.2 - 0.25 0.25 - 0.3 0.3 - 0.35 0.35 - 0.4 0.4 - 0.45 0.45 - 0.5 0.5 - 0.55 0.55 - 0.6 > 0.6 <p style="text-align: center; font-size: small;">Bare soil, rocks clouds, rain, snow, etc.</p> <p style="text-align: center; font-size: small;">Moderate</p> <p style="text-align: center; font-size: small;">Good</p> <p style="text-align: center; font-size: small;">Very good</p> <p style="text-align: center; font-size: small;">Agriculture vigour is moderate at isolated places of the state.</p>
<p>As per SPI value (cumulative 4weeks -26 December, 2024 to 22 January, 2025), moderately wet condition was there in Cuttack district.</p>	<p>According to NDVI map (12.02.2025 to 18.02.2025) the agriculture vigor was moderate over Narasinghpur, Badamba, Tigiria, Banki and Banki Dampada and light over other places of Cuttack district.</p>

LEGEND (Medium range block level rainfall forecast)		
	No Rainfall	00 mm
	Very light rainfall	01-05 mm
	Light rainfall	06-10 mm
	Light to moderate rainfall	11-30 mm
	Moderate to heavy rainfall	31-65 mm
	Heavy to very heavy rainfall	66-115 mm
	Extremely heavy rainfall	>116 mm

DrRanjan Kumar Mohanta
 (Nodal officer)

Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightning alert in your area


















GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-555
Date- 25/02/2025

Weather (Past and Forecasted) based Disease and pest in Major Crops and vegetables

Sl No	Crop	Stage and condition of the crop	Anticipated Disease and pest	
1	Pulses (Green gram and black gram)	Flowering to pod development		Anthracnose
				Cercospera leaf spot
				YMV
2	Mustard	Flowering to maturity		Saw Fly
				Leaf webber
3	Rabi Groundnut	Flowering to maturity		Spodotera/Tobacco caterpillar
6	Tomato	Flowering to Fruting		Late blight
				Bacterial Canker
				Sun burn
				Flower drop
				Bacterial wilt
7	Potato	Flowering to fruiting stage		Potato tuber moth
				Cut worm

DrRanjan Kumar Mohanta
 (Nodal officer)

Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area









GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-555
Date- 25/02/2025

8	Onion	Flowering		Purple blight
				Stemphylium Blight
10	Mango	Flowering		Mango hooper
				Thrips



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-555
Date- 25/02/2025

AGROMET ADVISORY BULLETIN FOR ATHAGAD

District Past week average weather condition and Block past week average Rainfall condition

The block received average 08 mm and district received average 03 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.42°C (-1.33) and 20.57°C (+1.79), respectively. The mean relative humidity was 63-87%.

Forecasted period: Today(8.30IST) to 02nd March (8.30IST)2025

Weather advisory: Dry weather is forecasted for next 5days in the block. The day temperature will be 32-36°C and night temperature will be 21-22°C for upcoming 5 days. Maximum and minimum relative humidity will be 58-85% and 13-23% respectively. **There will be clear to partly cloudy sky for next 5days.**

Block	Athagad (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	25/02/2025	26/02/2025	27/02/2025	28/02/2025	01/03/2025
Rainfall (mm)	0	0	0	0	0
T-Max (°C)	33	32	33	34	36
T-Min (°C)	21	21	21	22	22
Cloud Cover (octa)	1	0	0	1	1
RH Max (%)	85	83	82	65	58
RH Min (%)	23	22	17	16	13
Wind Speed (kmph)	6	8	8	10	14
Wind Direction (deg)	158	189	198	193	206

Agromet Advisory

General advisory:

Farmers should undertake field preparation for cultivation of different summer vegetables like sponge gourd, bottle gourd, pumpkin, water melon, musk melon etc. If seedlings are already raised in polythene bags; transplant in main field @ 2-3 seedlings per pit. Irrigation, fertilizer and pesticide application should be resumed as there will clear sky for upcoming 5 days.

Pulses:

November to December sown crop is at pod formation to seed development (nine to twelve week) stage.

In case of **YMV infestation** uproots and destroys the infected plant and spray imidacloprid 50ml/acre mixing with 200 lit of water. There may be chances of infestation of **Anthraxnose and Cercosera leaf spot** in Pulses. To manage these spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre in 200 ltr water.

Rabi-Groundnut:

Do not disturb the soil after flower initiation in groundnut. Do irrigation in the field based upon soil moisture. Due to abrupt rise and fall in temperature, there is possibility of attack of **Tobacco caterpillar** (Spodoptera litura). The forecasted weather is also favourable for its proliferation. This is a polyphagous pest. So following integrated approach should be started. Use Pheromone trap using Spodo lure to monitor adult population of the insect. In case of appearance of the caterpillar, prepare poison bait with 1 kg rice bran+ 200g jaggery/gur+20-25g Cartap Hydrochloride +Water. Keep the bait at 20-25 different places in the affected field.

Mustard: present weather may favour the infestation of **saw fly, leaf Webber**. To control saw fly, spray 400ml Endosulphan/acre, to control leaf webber spray 200 ml Ithofenprox or Deltamethrin per acre in 200 litre water.

Vegetables:

Tomato: Transplanted seedlings are at 6-8 week stage in main field. **Due deficiency of boron due to hormonal disbalance flower drop occurs.**

Control: Apply timely and required amount of irrigation. Spray Boron 20% 2 gram/lit during early flowering and 15 days later. Due to increase in temperature there may be chances of **bacterial canker and Sun burn** disease in tomato at fruiting stage. To control bacterial canker Farmers should spray Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water. They should use shade net to protect the crop from sun burn. **Bacterial wilt** may be reported in tomato field, to control it drenching with copper oxychloride @ 3g+ plantomycin @1 g/ litre of water.

• **Potato:** There are chances of **Potato Tuber Moth** infestation in potato. To manage this pest chemically spray Spinosad 45 % SC @ 75-ml/acre or Emamectin Benzoate 5% SG @ 80-gram/ acre or Chlorantraniliprole 18.5 % SC @80- ml/acre. To manage **cut worm** in potato crop spray Chlorpyrifos 50% + Cypermethrin 5 % E.C (Lethal-505/Super D) @ 400 ml/acre by mixing it in 200 litre of water.

• **Onion:** There are chances of **purple blotch and Stemphyllium blight** disease incidence in onion crop. To manage purple blotch and Stemphyllium leaf blight disease in onion spray Mancozeb 75% WP @ 600gram/acre or Difenconazole 25% E.C @ 200ml/acre or Tebuconazole 25.9% E.C @ 100ml/acre by mixing in 200 litre of water at 15 days interval.

• **Cucutbits:** The current weather is favorable for attack of **red pumpkin beetle** in Cucumber & Bitter gourd. To control it spray Neem based pesticide 300ppm@5ml/litre of water at initial stages and at later stages spray Acephate@1g/litre of water.

Fruits:

• **Mango:** Present weather conditions are favourable for the incidence of **Mango hoppers**. To control go for spraying of phosphamidon @ 0.5ml/l (or) dimethoate 2ml/l (or) chlorpyrifos @2.5ml/l. during initial stages of flowering. Spray Alpha Naphthyl Acetic Acid 4.5% SL @ 4.5 ml in 10 litre of water when tender fruits are in pea size. With the help of ring method of irrigation apply 25 kg FYM, 500 gram of urea and 1 kg of potash and phosphate at each tree. For **thrips** control, spraying of Acephate @1.5 g /litre of water is advocated.

• **Cashew:** Prevailing conditions are suitable for the incidence of Tea mosquito bug. Farmers are advised to go for spraying of Neem oil @ 5ml/L + Monocrotophos @ 1.6ml/L at new blush stage and Chlorpyrifos @ 2ml/L at flowering stage.

Mushroom: Initiate cultivation of paddy straw mushroom. As dry weather condition is prevailed in Cuttack district, Spray water if the mushrooms are at fruiting stage. Cover the Mushroom unit with water soaked Gunny bags to maintain the inside temperature.



DrRanjan Kumar Mohanta
 (Nodal officer)

Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightning alert in your area





INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-555
Date- 25/02/2025

AGROMET ADVISORY BULLETIN FOR BADAMBA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 01 mm and district received average 03 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.42°C (-1.33) and 20.57°C (+1.79), respectively. The mean relative humidity was 63-87%.

Forecasted period: Today(8.30IST) to 02nd March (8.30IST)2025

Weather advisory: Dry weather is forecasted for next 5 days in the block. The day temperature will be 33-36°C and night temperature will be 20-21°C for upcoming 5 days. Maximum and minimum relative humidity will be 51-87% and 13-22% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Badamba (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	25/02/2025	26/02/2025	27/02/2025	28/02/2025	01/03/2025
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	33	33	33	34	36
T-Min(°C)	20	20	20	21	21
Cloud Cover(octa)	0	1	0	1	2
RH Max (%)	87	78	76	57	51
RH Min (%)	21	22	16	16	13
Wind Speed (Kmph)	6	8	7	10	13
Wind Direction (deg)	158	171	203	193	206

Agromet Advisory

General advisory:

Farmers should undertake field preparation for cultivation of different summer vegetables like sponge gourd, bottle gourd, pumpkin, water melon, musk melon etc. If seedlings are already raised in polythene bags; transplant in main field @ 2-3 seedlings per pit. Irrigation, fertilizer and pesticide application should be resumed as there will clear sky for upcoming 5 days.

Pulses:

November to December sown crop is at pod formation to seed development (nine to twelve week) stage.

In case of **YMV infestation** uproots and destroys the infected plant and spray imidacloprid 50ml/acre mixing with 200 lit of water. There may be chances of infestation of **Anthraxnose and Cercopersa leaf spot** in Pulses. To manage these spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre in 200 ltr water.

Rabi-Groundnut:

Do not disturb the soil after flower initiation in groundnut. Do irrigation in the field based upon soil moisture. Due to abrupt rise and fall in temperature, there is possibility of attack of **Tobacco caterpillar** (Spodoptera litura). The forecasted weather is also favourable for its proliferation. This is a polyphagous pest. So following integrated approach should be started. Use Pheromone trap using Spodo lure to monitor adult population of the insect. In case of appearance of the caterpillar, prepare poison bait with 1 kg rice bran+ 200g jaggery/gur+20-25g Cartap Hydrochloride +Water. Keep the bait at 20-25 different places in the affected field.

Mustard: present weather may favour the infestation of **saw fly, leaf Webber**. To control saw fly, spray 400ml Endosulphan/acre, to control leaf webber spray 200 ml Ithofenprox or Deltamethrin per acre in 200 litr water.

Vegetables:

Tomato: Transplanted seedlings are at 6-8 week stage in main field. **Due deficiency of boron due to hormonal disbalance flower drop occurs.** Control: Apply timely and required amount of irrigation. Spray Boron 20% 2 gram/lit during early flowering and 15 days later. Due to increase in temperature there may be chances of **bacterial canker and Sun burn** disease in tomato at fruiting stage. To control bacterial canker Farmers should spray Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water. They should use shade net to protect the crop from sun burn. **Bacterial wilt** may be reported in tomato field, to control it drenching with copper oxychloride @ 3g+ plantomycin @1 g/ litre of water.

• **Potato:** There are chances of **Potato Tuber Moth** infestation in potato. To manage this pest chemically spray Spinosad 45 % SC @ 75-ml/acre or Emamectin Benzoate 5% SG @ 80-gram/ acre or Chlorantraniliprole 18.5 % SC @80- ml/acre. To manage **cut worm** in potato crop spray Chlorpyrifos 50% + Cypermethrin 5 % E.C (Lethal-505/Super D) @ 400 ml/acre by mixing it in 200 litre of water.

• **Onion:** There are chances of **purple blotch and Stemphyllium blight** disease incidence in onion crop. To manage purple blotch and Stemphyllium leaf blight disease in onion spray Mancozeb 75% WP @ 600gram/acre or Difenoconazole 25% E.C @ 200ml/acre or Tebuconazole 25.9% E.C @ 100ml/acre by mixing in 200 litre of water at 15 days interval.

• **Cucutbits:** The current weather is favorable for attack of **red pumpkin beetle** in Cucumber & Bitter gourd. To control it spray Neem based pesticide 300ppm@5ml/litre of water at initial stages and at later stages spray Acephate@1g/litre of water.

Fruits:

• **Mango:** Present weather conditions are favourable for the incidence of **Mango hoppers**. To control go for spraying of phosphamidon @ 0.5ml/l (or) dimethoate 2ml/l (or) chloropyrifos @2.5ml/l. during initial stages of flowering. Spray Alpha Naphthyl Acetic Acid 4.5% SL @ 4.5 ml in 10 litre of water when tender fruits are in pea size. With the help of ring method of irrigation apply 25 kg FYM, 500 gram of urea and 1 kg of potash and phosphate at each tree. For **thrips** control, spraying of Acephate @1.5 g /litre of water is advocated.

• **Cashew:** Prevailing conditions are suitable for the incidence of Tea mosquito bug. Farmers are advised to go for spraying of Neem oil @ 5ml/L + Monocrotophos @ 1.6ml/L at new blush stage and Chlorpyriqos @ 2ml/L at flowering stage.

Mushroom: Initiate cultivation of paddy straw mushroom. As dry weather condition is prevailed in Cuttack district, Spray water if the mushrooms are at fruiting stage. Cover the Mushroom unit with water soaked Gunny bags to maintain the inside temperature.



DrRanjan Kumar Mohanta
 (Nodal officer)

Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-555
Date- 25/02/2025

AGROMET ADVISORY BULLETIN FOR BANKI

District Past week average weather condition

The district received average 03 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.42°C (-1.33) and 20.57°C (+1.79), respectively. The mean relative humidity was 63-87%.

Forecasted period: Today(8.30IST) to 02nd March (8.30IST)2025

Weather advisory; Dry weather is forecasted for next 5 days in the block. The day temperature will be 32-36°C and night temperature will be 20-22°C for upcoming 5 days. Maximum and minimum relative humidity will be 57-87% and 13-23% respectively. **There will be clear to partly cloudy sky for next 5 days.**

Block	Banki (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	25/02/2025	26/02/2025	27/02/2025	28/02/2025	01/03/2025
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	33	32	33	34	36
T-Min(°C)	21	21	20	21	22
Cloud Cover(octa)	0	0	0	1	1
RH Max (%)	87	82	80	63	57
RH Min (%)	22	23	18	17	13
Wind Speed (Kmph)	6	9	8	11	15
Wind Direction (deg)	158	180	203	198	206

Agromet Advisory

General advisory:

Farmers should undertake field preparation for cultivation of different summer vegetables like sponge gourd, bottle gourd, pumpkin, water melon, musk melon etc. If seedlings are already raised in polythene bags; transplant in main field @ 2-3 seedlings per pit. Irrigation, fertilizer and pesticide application should be resumed as there will clear sky for upcoming 5 days.

Pulses:

November to December sown crop is at pod formation to seed development (nine to twelve week) stage.

In case of **YMV infestation** uproots and destroys the infected plant and spray imidacloprid 50ml/acre mixing with 200 lit of water. There may be chances of infestation of **Anthraxnose and Cercopsera leaf spot** in Pulses. To manage these spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre in 200 ltr water.

Rabi-Groundnut:

Do not disturb the soil after flower initiation in groundnut. Do irrigation in the field based upon soil moisture. Due to abrupt rise and fall in temperature, there is possibility of attack of **Tobacco caterpillar** (Spodoptera litura). The forecasted weather is also favourable for its proliferation. This is a polyphagous pest. So following integrated approach should be started. Use Pheromone trap using Spodo lure to monitor adult population of the insect. In case of appearance of the caterpillar, prepare poison bait with 1 kg rice bran+ 200g jaggery/gur+20-25g Cartap Hydrochloride +Water. Keep the bait at 20-25 different places in the affected field.

Mustard: present weather may favour the infestation of **saw fly, leaf Webber**. To control saw fly, spray 400ml Endosulphan/acre, to control leaf webber spray 200 ml Ithofenprox or Deltamethrin per acre in 200 litre water.

Vegetables:

Tomato: Transplanted seedlings are at 6-8 week stage in main field. **Due deficiency of boron due to hormonal disbalance flower drop occurs.** Control: Apply timely and required amount of irrigation. Spray Boron 20% 2 gram/lit during early flowering and 15 days later. Due to increase in temperature there may be chances of **bacterial canker and Sun burn** disease in tomato at fruiting stage. To control bacterial canker farmers should spray Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water. They should use shade net to protect the crop from sun burn. **Bacterial wilt** may be reported in tomato field, to control it drenching with copper oxychloride @ 3g+ plantomycin @1 g/ litre of water.

• **Potato:** There are chances of **Potato Tuber Moth** infestation in potato. To manage this pest chemically spray Spinosad 45 % SC @ 75-ml/acre or Emamectin Benzoate 5% SG @ 80-gram/ acre or Chlorantraniliprole 18.5 % SC @80- ml/acre. To manage **cut worm** in potato crop spray Chlorpyrifos 50% + Cypermethrin 5 % E.C (Lethal-505/Super D) @ 400 ml/acre by mixing it in 200 litre of water.

• **Onion:** There are chances of **purple blotch and Stemphyllium blight** disease incidence in onion crop. To manage purple blotch and Stemphyllium leaf blight disease in onion spray Mancozeb 75% WP @ 600gram/acre or Difenconazole 25% E.C @ 200ml/acre or Tebuconazole 25.9% E.C @ 100ml/acre by mixing in 200 litre of water at 15 days interval.

• **Cucurbits:** The current weather is favorable for attack of **red pumpkin beetle** in Cucumber & Bitter gourd. To control it spray Neem based pesticide 300ppm@5ml/litre of water at initial stages and at later stages spray Acephate@1g/litre of water.

Fruits:

• **Mango:** Present weather conditions are favourable for the incidence of **Mango hoppers**. To control go for spraying of phosphamidon @ 0.5ml/l (or) dimethoate 2ml/l (or) chlorpyrifos @2.5ml/l. during initial stages of flowering. Spray Alpha Naphthyl Acetic Acid 4.5% SL @ 4.5 ml in 10 litre of water when tender fruits are in pea size. With the help of ring method of irrigation apply 25 kg FYM, 500 gram of urea and 1 kg of potash and phosphate at each tree. For **thrips** control, spraying of Acephate @1.5 g /litre of water is advocated.

• **Cashew:** Prevailing conditions are suitable for the incidence of Tea mosquito bug. Farmers are advised to go for spraying of Neem oil @ 5ml/L + Monocrotophos @ 1.6ml/L at new blush stage and Chlorpyrifos @ 2ml/L at flowering stage.

Mushroom: Initiate cultivation of paddy straw mushroom. As dry weather condition is prevailed in Cuttack district, Spray water if the mushrooms are at fruiting stage. Cover the Mushroom unit with water soaked Gunny bags to maintain the inside temperature.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area





**GRAMIN KRISHI MAUSAM SAUKHYA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com**



**Bulletin No.-555
Date- 25/02/2025**

AGROMET ADVISORY BULLETIN FOR BANKI-DAMPADA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 03 mm and district received average 03 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.42°C (-1.33) and 20.57°C (+1.79), respectively. The mean relative humidity was 63-87%.

Forecasted period: Today(8.30IST) to 02nd March (8.30IST)2025

Weather Advisory: Dry weather is forecasted for next 5 days in the block. The day temperature will be 32-36°C and night temperature will be 20-22°C for upcoming 5 days. Maximum and minimum relative humidity will be 58-87% and 14-23% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Banki-Dampada (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	25/02/2025	26/02/2025	27/02/2025	28/02/2025	01/03/2025
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	33	32	34	34	36
T-Min(°C)	21	21	20	21	22
Cloud Cover(octa)	0	0	0	1	1
RH Max (%)	87	83	81	64	58
RH Min (%)	23	23	18	17	14
Wind Speed (Kmph)	6	9	8	11	15
Wind Direction (deg)	158	180	203	201	206

Agromet Advisory

General advisory:

Farmers should undertake field preparation for cultivation of different summer vegetables like sponge gourd, bottle gourd, pumpkin, water melon, musk melon etc. If seedlings are already raised in polythene bags; transplant in main field @ 2-3 seedlings per pit. Irrigation, fertilizer and pesticide application should be resumed as there will clear sky for upcoming 5 days.

Pulses:

November to December sown crop is at pod formation to seed development (nine to twelve week) stage.

In case of **YMV infestation** uproots and destroys the infected plant and spray imidacloprid 50ml/acre mixing with 200 lit of water. There may be chances of infestation of **Anthraxnose and Cercopersa leaf spot** in Pulses. To manage these spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre in 200 ltr water.

Rabi-Groundnut:

Do not disturb the soil after flower initiation in groundnut. Do irrigation in the field based upon soil moisture. Due to abrupt rise and fall in temperature, there is possibility of attack of **Tobacco caterpillar** (Spodoptera litura). The forecasted weather is also favourable for its proliferation. This is a polyphagous pest. So following integrated approach should be started. Use Pheromone trap using Spodo lure to monitor adult population of the insect. In case of appearance of the caterpillar, prepare poison bait with 1 kg rice bran+ 200g jaggery/gur+20-25g Cartap Hydrochloride +Water. Keep the bait at 20-25 different places in the affected field.

Mustard: present weather may favour the infestation of **saw fly, leaf Webber**. To control saw fly, spray 400ml Endosulphan/acre, to control leaf webber spray 200 ml lthofenprox or Deltamethrin per acre in 200 litr water.

Vegetables:

Tomato: Transplanted seedlings are at 6-8 week stage in main field. **Due deficiency of boron or due to hormonal disbalance flower drop occurs.**

Control: Apply timely and required amount of irrigation. Spray Boron 20% 2 gram/lit during early flowering and 15 days later. Due to increase in temperature there may be chances of **bacterial canker and Sun burn** disease in tomato at fruiting stage. To control bacterial canker farmers should spray Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water. They should use shade net to protect the crop from sun burn. **Bacterial wilt** may be reported in tomato field, to control it drenching with copper oxychloride @ 3g+ plantomycin @ 1 g/ litre of water.

• **Potato:** There are chances of **Potato Tuber Moth** infestation in potato. To manage this pest chemically spray Spinosad 45 % SC @ 75-ml/acre or Emamectin Benzoate 5% SG @ 80-gram/ acre or Chlorantraniliprole 18.5 % SC @ 80- ml/acre. To manage **cut worm** in potato crop spray Chlorpyrifos 50% + Cypermethrin 5 % E.C (Lethal-505/Super D) @ 400 ml/acre by mixing it in 200 litre of water.

• **Onion:** There are chances of **purple blotch and Stemphyllium blight** disease incidence in onion crop. To manage purple blotch and Stemphyllium leaf blight disease in onion spray Mancozeb 75% WP @ 600gram/acre or Difenconazole 25% E.C @ 200ml/acre or Tebuconazole 25.9% E.C @ 100ml/acre by mixing in 200 litre of water at 15 days interval.

• **Cucutbits:** The current weather is favorable for attack of **red pumpkin beetle** in Cucumber & Bitter gourd. To control it spray Neem based pesticide 300ppm@5ml/litre of water at initial stages and at later stages spray Acephate@1g/litre of water.

Fruits:

• **Mango:** Present weather conditions are favourable for the incidence of **Mango hoppers**. To control go for spraying of phosphamidon @ 0.5ml/l (or) dimethoate 2ml/l (or) chlorpyrifos @ 2.5ml/l. during initial stages of flowering. Spray Alpha Naphthyl Acetic Acid 4.5% SL @ 4.5 ml in 10 litre of water when tender fruits are in pea size. With the help of ring method of irrigation apply 25 kg FYM, 500 gram of urea and 1 kg of potash and phosphate at each tree. For **thrips** control, spraying of Acephate @ 1.5 g /litre of water is advocated.

• **Cashew:** Prevailing conditions are suitable for the incidence of Tea mosquito bug. Farmers are advised to go for spraying of Neem oil @ 5ml/L + Monocrotophos @ 1.6ml/L at new blush stage and Chlorpyrifos @ 2ml/L at flowering stage.

• **Mushroom:** Initiate cultivation of paddy straw mushroom. As dry weather condition is prevailed in Cuttack district, Spray water if the mushrooms are at fruiting stage. Cover the Mushroom unit with water soaked Gunny bags to maintain the inside temperature.



**DrRanjan Kumar Mohanta
(Nodal officer)**

**Follow Meghdoot Mobile APP for Agromet
advisories in local language and DAMINI
APP for Lightning alert in your area**





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-555
Date- 25/02/2025

AGROMET ADVISORY BULLETIN FOR BARANGA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 04 mm and district received average 03 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.42°C (-1.33) and 20.57°C (+1.79), respectively. The mean relative humidity was 63-87%.

Forecasted period: Today(8.30IST) to 02nd March (8.30IST)2025

Weather advisory: *Dry weather is forecasted for next 5 days in the block.* The day temperature will be 31-36°C and night temperature will be 22-23°C for upcoming 5 days. Maximum and minimum relative humidity will be 66-85% and 16-29% respectively. **There will be clear to partly cloudy sky for next 5 days.**

Block	Baranga (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	25/02/2025	26/02/2025	27/02/2025	28/02/2025	01/03/2025
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	32	31	33	34	36
T-Min(°C)	22	22	22	22	23
Cloud Cover	1	0	0	1	1
RH Max (%)	85	85	85	77	66
RH Min (%)	29	25	22	20	16
Wind Speed (Kmph)	7	10	10	13	17
Wind Direction (deg)	150	195	198	203	207

Agromet Advisory

General advisory:

Farmers should undertake field preparation for cultivation of different summer vegetables like sponge gourd, bottle gourd, pumpkin, water melon, musk melon etc. If seedlings are already raised in polythene bags; transplant in main field @ 2-3 seedlings per pit. Irrigation, fertilizer and pesticide application should be resumed as there will clear sky for upcoming 5 days.

Pulses:

November to December sown crop is at pod formation to seed development (nine to twelve week) stage.

In case of **YMV infestation** uproots and destroys the infected plant and spray imidacloprid 50ml/acre mixing with 200 lit of water. There may be chances of infestation of **Anthraxnose and Cercopsera leaf spot** in Pulses. To manage these spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre in 200 ltr water.

Rabi-Groundnut:

Do not disturb the soil after flower initiation in groundnut. Do irrigation in the field based upon soil moisture. Due to abrupt rise and fall in temperature, there is possibility of attack of **Tobacco caterpillar** (Spodoptera litura). The forecasted weather is also favourable for its proliferation. This is a polyphagous pest. So following integrated approach should be started. Use Pheromone trap using Spodo lure to monitor adult population of the insect. In case of appearance of the caterpillar, prepare poison bait with 1 kg rice bran+ 200g jaggery/gur+20-25g Cartap Hydrochloride +Water. Keep the bait at 20-25 different places in the affected field.

Mustard: present weather may favour the infestation of **saw fly, leaf Webber**. To control saw fly, spray 400ml Endosulphan/acre, to control leaf webber spray 200 ml Ithofenprox or Deltamethrin per acre in 200 litre water.

Vegetables:

Tomato: Transplanted seedlings are at 6-8 week stage in main field. **Due deficiency of boron due to hormonal disbalance flower drop occurs.** Control: Apply timely and required amount of irrigation. Spray Boron 20% 2 gram/lit during early flowering and 15 days later. Due to increase in temperature there may be chances of **bacterial canker and Sun burn** disease in tomato at fruiting stage. To control bacterial canker farmers should spray Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water. They should use shade net to protect the crop from sun burn. **Bacterial wilt** may be reported in tomato field, to control it drenching with copper oxychloride @ 3g+ plantomycin @1 g/ litre of water.

• **Potato:** There are chances of **Potato Tuber Moth** infestation in potato. To manage this pest chemically spray Spinosad 45 % SC @ 75-ml/acre or Emamectin Benzoate 5% SG @ 80-gram/ acre or Chlorantraniliprole 18.5 % SC @80- ml/acre. To manage **cut worm** in potato crop spray Chlorpyrifos 50% + Cypermethrin 5 % E.C (Lethal-505/Super D) @ 400 ml/acre by mixing it in 200 litre of water.

• **Onion:** There are chances of **purple blotch and Stemphylium blight** disease incidence in onion crop. To manage purple blotch and Stemphylium leaf blight disease in onion spray Mancozeb 75% WP @ 600gram/acre or Difenconazole 25% E.C @ 200ml/acre or Tebuconazole 25.9% E.C @ 100ml/acre by mixing in 200 litre of water at 15 days interval.

• **Cucutbits:** The current weather is favorable for attack of **red pumpkin beetle** in Cucumber & Bitter gourd. To control it spray Neem based pesticide 300ppm@5ml/litre of water at initial stages and at later stages spray Acephate@1g/litre of water.

Fruits:

• **Mango:** Present weather conditions are favourable for the incidence of **Mango hoppers**. To control go for spraying of phosphamidon @ 0.5ml/l (or) dimethoate 2ml/l (or) chlorpyrifos @2.5ml/l. during initial stages of flowering. Spray Alpha Naphthyl Acetic Acid 4.5% SL @ 4.5 ml in 10 litre of water when tender fruits are in pea size. With the help of ring method of irrigation apply 25 kg FYM, 500 gram of urea and 1 kg of potash and phosphate at each tree. For **thrips** control, spraying of Acephate @1.5 g /litre of water is advocated.

• **Cashew:** Prevailing conditions are suitable for the incidence of Tea mosquito bug. Farmers are advised to go for spraying of Neem oil @ 5ml/L + Monocrotophos @ 1.6ml/L at new blush stage and Chlorpyrifos @ 2ml/L at flowering stage.

• **Mushroom:** Initiate cultivation of paddy straw mushroom. As dry weather condition is prevailed in Cuttack district, Spray water if the mushrooms are at fruiting stage. Cover the Mushroom unit with water soaked Gunny bags to maintain the inside temperature.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-555
Date- 25/02/2025

AGROMET ADVISORY BULLETIN FOR CUTTACK SADAR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 03 mm and district received average 03 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.42°C (-1.33) and 20.57°C (+1.79), respectively. The mean relative humidity was 63-87%.

Forecasted period: Today(8.30IST) to 02nd March (8.30IST)2025

Weather advisory: Dry weather is forecasted for next 5 days in the block. The day temperature will be 32-36°C and night temperature will be 22-23°C for upcoming 5 days. Maximum and minimum relative humidity will be 64-85% and 15-27% respectively. **There will be clear to partly cloudy sky for next 5 days.**

Block	Cuttack sadar (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	25/02/2025	26/02/2025	27/02/2025	28/02/2025	01/03/2025
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	32	32	33	34	36
T-Min(°C)	22	22	22	22	23
Cloud Cover(octa)	1	0	0	1	1
RH Max (%)	84	84	85	74	64
RH Min (%)	27	24	20	18	15
Wind Speed (Kmph)	7	9	9	11	16
Wind Direction (deg)	180	195	203	203	207

Agromet Advisory

General advisory:

Farmers should undertake field preparation for cultivation of different summer vegetables like sponge gourd, bottle gourd, pumpkin, water melon, musk melon etc. If seedlings are already raised in polythene bags; transplant in main field @ 2-3 seedlings per pit. Irrigation, fertilizer and pesticide application should be resumed as there will clear sky for upcoming 5 days.

Pulses:

November to December sown crop is at pod formation to seed development (nine to twelve week) stage.

In case of **YMV infestation** uproots and destroys the infected plant and spray imidacloprid 50ml/acre mixing with 200 lit of water. There may be chances of infestation of **Anthraxnose and Cercopsera leaf spot** in Pulses. To manage these spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre in 200 ltr water.

Rabi-Groundnut:

Do not disturb the soil after flower initiation in groundnut. Do irrigation in the field based upon soil moisture. Due to abrupt rise and fall in temperature, there is possibility of attack of **Tobacco caterpillar** (Spodoptera litura). The forecasted weather is also favourable for its proliferation. This is a polyphagous pest. So following integrated approach should be started. Use Pheromone trap using Spodo lure to monitor adult population of the insect. In case of appearance of the caterpillar, prepare poison bait with 1 kg rice bran+ 200g jaggery/gur+20-25g Cartap Hydrochloride +Water. Keep the bait at 20-25 different places in the affected field.

Mustard: present weather may favour the infestation of **saw fly, leaf Webber**. To control saw fly, spray 400ml Endosulphan/acre, to control leaf webber spray 200 ml Ithofenprox or Deltamethrin per acre in 200 litre water.

Vegetables:

Tomato: Transplanted seedlings are at 6-8 week stage in main field. **Due deficiency of boron due to hormonal disbalance flower drop occurs.** Control: Apply timely and required amount of irrigation. Spray Boron 20% 2 gram/lit during early flowering and 15 days later. Due to increase in temperature there may be chances of **bacterial canker and Sun burn** disease in tomato at fruiting stage. To control bacterial canker farmers should spray Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water. They should use shade net to protect the crop from sun burn. **Bacterial wilt** may be reported in tomato field, to control it drenching with copper oxychloride @ 3g+ plantomycin @1 g/ litre of water.

• **Potato:** There are chances of **Potato Tuber Moth** infestation in potato. To manage this pest chemically spray Spinosad 45 % SC @ 75-ml/acre or Emamectin Benzoate 5% SG @ 80-gram/ acre or Chlorantraniliprole 18.5 % SC @80- ml/acre. To manage **cut worm** in potato crop spray Chlorpyrifos 50% + Cypermethrin 5 % E.C (Lethal-505/Super D) @ 400 ml/acre by mixing it in 200 litre of water.

• **Onion:** There are chances of **purple blotch and Stemphylium blight** disease incidence in onion crop. To manage purple blotch and Stemphylium leaf blight disease in onion spray Mancozeb 75% WP @ 600gram/acre or Difenoconazole 25% E.C @ 200ml/acre or Tebuconazole 25.9% E.C @ 100ml/acre by mixing in 200 litre of water at 15 days interval.

• **Cucutbits:** The current weather is favorable for attack of **red pumpkin beetle** in Cucumber & Bitter gourd. To control it spray Neem based pesticide 300ppm@5ml/litre of water at initial stages and at later stages spray Acephate@1g/litre of water.

Fruits:

• **Mango:** Present weather conditions are favourable for the incidence of **Mango hoppers**. To control go for spraying of phosphamidon @ 0.5ml/l (or) dimethoate 2ml/l (or) chlorpyrifos @2.5ml/l. during initial stages of flowering. Spray Alpha Naphthyl Acetic Acid 4.5% SL @ 4.5 ml in 10 litre of water when tender fruits are in pea size. With the help of ring method of irrigation apply 25 kg FYM, 500 gram of urea and 1 kg of potash and phosphate at each tree. For **thrips** control, spraying of Acephate @1.5 g /litre of water is advocated.

• **Cashew:** Prevailing conditions are suitable for the incidence of Tea mosquito bug. Farmers are advised to go for spraying of Neem oil @ 5ml/L + Monocrotophos @ 1.6ml/L at new blush stage and Chlorpyrifos @ 2ml/L at flowering stage.

Mushroom: Initiate cultivation of paddy straw mushroom. As dry weather condition is prevailed in Cuttack district, Spray water if the mushrooms are at fruiting stage. Cover the Mushroom unit with water soaked Gunny bags to maintain the inside temperature.



DrRanjan Kumar Mohanta
 (Nodal officer)

Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-555
Date- 25/02/2025

AGROMET ADVISORY BULLETIN FOR KANTAPADA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 01 mm and district received average 03 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.42°C (-1.33) and 20.57°C (+1.79), respectively. The mean relative humidity was 63-87%.

Forecasted period: Today(8.30IST) to 02nd March (8.30IST)2025

Weather advisory: Dry weather is forecasted for next 5 days in the block. The day temperature will be 31-35°C and night temperature will be 22-24°C for upcoming 5 days. Maximum and minimum relative humidity will be 71-86% and 19-33% respectively. **There will be clear to partly cloudy sky for next 5 days.**

Block	Kantapada (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	25/02/2025	26/02/2025	27/02/2025	28/02/2025	01/03/2025
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	32	31	33	34	35
T-Min(°C)	22	22	22	22	24
Cloud Cover(octa)	1	0	0	1	1
RH Max (%)	85	86	86	82	71
RH Min (%)	33	27	26	24	19
Wind Speed (Kmph)	7	10	11	13	17
Wind Direction (deg)	135	135	203	203	204

Agromet Advisory

General advisory:

Farmers should undertake field preparation for cultivation of different summer vegetables like sponge gourd, bottle gourd, pumpkin, water melon, musk melon etc. If seedlings are already raised in polythene bags; transplant in main field @ 2-3 seedlings per pit. Irrigation, fertilizer and pesticide application should be resumed as there will clear sky for upcoming 5 days.

Pulses:

November to December sown crop is at pod formation to seed development (nine to twelve week) stage.

In case of **YMV infestation** uproots and destroys the infected plant and spray imidacloprid 50ml/acre mixing with 200 lit of water. There may be chances of infestation of **Anthraxnose and Cercopsora leaf spot** in Pulses. To manage these spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre in 200 ltr water.

Rabi-Groundnut:

Do not disturb the soil after flower initiation in groundnut. Do irrigation in the field based upon soil moisture. Due to abrupt rise and fall in temperature, there is possibility of attack of **Tobacco caterpillar** (Spodoptera litura). The forecasted weather is also favourable for its proliferation. This is a polyphagous pest. So following integrated approach should be started. Use Pheromone trap using Spodo lure to monitor adult population of the insect. In case of appearance of the caterpillar, prepare poison bait with 1 kg rice bran+ 200g jaggery/gur+20-25g Cartap Hydrochloride +Water. Keep the bait at 20-25 different places in the affected field.

Mustard: present weather may favour the infestation of **saw fly, leaf Webber**. To control saw fly, spray 400ml Endosulphan/acre, to control leaf webber spray 200 ml Ithofenprox or Deltamethrin per acre in 200 litr water.

Vegetables:

Tomato: Transplanted seedlings are at 6-8 week stage in main field. **Due deficiency of boron due to hormonal disbalance flower drop occurs.**

Control: Apply timely and required amount of irrigation. Spray Boron 20% 2 gram/lit during early flowering and 15 days later. Due to increase in temperature there may be chances of **bacterial canker and Sun burn** disease in tomato at fruiting stage. To control bacterial canker Farmers should spray Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water. They should use shade net to protect the crop from sun burn. **Bacterial wilt** may be reported in tomato field, to control it drenching with copper oxychloride @ 3g+ plantomycin @1 g/ litre of water.

• **Potato:** There are chances of **Potato Tuber Moth** infestation in potato. To manage this pest chemically spray Spinosad 45 % SC @ 75-ml/acre or Emamectin Benzoate 5% SG @ 80-gram/ acre or Chlorantraniliprole 18.5 % SC @80- ml/acre. To manage **cut worm** in potato crop spray Chlorpyrifos 50% + Cypermethrin 5 % E.C (Lethal-505/Super D) @ 400 ml/acre by mixing it in 200 litre of water.

• **Onion:** There are chances of **purple blotch and Stemphylium blight** disease incidence in onion crop. To manage purple blotch and Stemphylium leaf blight disease in onion spray Mancozeb 75% WP @ 600gram/acre or Difenoconazole 25% E.C @ 200ml/acre or Tebuconazole 25.9% E.C @ 100ml/acre by mixing in 200 litre of water at 15 days interval.

• **Cucutbits:** The current weather is favorable for attack of **red pumpkin beetle** in Cucumber & Bitter gourd. To control it spray Neem based pesticide 300ppm@5ml/litre of water at initial stages and at later stages spray Acephate@1g/litre of water.

Fruits:

• **Mango:** Present weather conditions are favourable for the incidence of **Mango hoppers**. To control go for spraying of phosphamidon @ 0.5ml/l (or) dimethoate 2ml/l (or) chlorpyrifos @2.5ml/l. during initial stages of flowering. Spray Alpha Naphthyl Acetic Acid 4.5% SL @ 4.5 ml in 10 litre of water when tender fruits are in pea size. With the help of ring method of irrigation apply 25 kg FYM, 500 gram of urea and 1 kg of potash and phosphate at each tree. For **thrips** control, spraying of Acephate @ 1.5 g /litre of water is advocated.

• **Cashew:** Prevailing conditions are suitable for the incidence of Tea mosquito bug. Farmers are advised to go for spraying of Neem oil @ 5ml/L + Monocrotophos @ 1.6ml/L at new blush stage and Chlorpyrifos @ 2ml/L at flowering stage.

Mushroom: Initiate cultivation of paddy straw mushroom. As dry weather condition is prevailed in Cuttack district, Spray water if the mushrooms are at fruiting stage. Cover the Mushroom unit with water soaked Gunny bags to maintain the inside temperature.



DrRanjan Kumar Mohanta
 (Nodal officer)

Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
(A Unit of ICAR-National Rice Research Institute)
Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-555
Date- 25/02/2025

AGROMET ADVISORY BULLETIN FOR MAHANGA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 01 mm and district received average 03 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.42°C (-1.33) and 20.57°C (+1.79), respectively. The mean relative humidity was 63-87%.

Forecasted period: Today(8.30IST) to 02nd March (8.30IST)2025

Weather advisory: Dry weather is forecasted for next 5 days in the block. The day temperature will be and it will be 31-36°C and night temperature will be 22-23°C for upcoming 5 days. Maximum and minimum relative humidity will be 63-84% and 19-32% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Mahanga (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	25/02/2025	26/02/2025	27/02/2025	28/02/2025	01/03/2025
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	31	31	33	34	36
T-Min(°C)	22	22	22	22	23
Cloud Cover(octal)	1	0	0	1	1
RH Max (%)	83	83	84	78	63
RH Min (%)	32	26	23	21	19
Wind Speed (Kmph)	7	8	9	11	18
Wind Direction (deg)	135	117	240	201	207

Agromet Advisory

General advisory:

Farmers should undertake field preparation for cultivation of different summer vegetables like sponge gourd, bottle gourd, pumpkin, water melon, musk melon etc. If seedlings are already raised in polythene bags; transplant in main field @ 2-3 seedlings per pit. Irrigation, fertilizer and pesticide application should be resumed as there will clear sky for upcoming 5 days.

Pulses:

November to December sown crop is at pod formation to seed development (nine to twelve week) stage.

In case of **YMV infestation** uproots and destroys the infected plant and spray imidacloprid 50ml/acre mixing with 200 lit of water. There may be chances of infestation of **Anthraxnose and Cercopsera leaf spot** in Pulses. To manage these spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre in 200 ltr water.

Rabi-Groundnut:

Do not disturb the soil after flower initiation in groundnut. Do irrigation in the field based upon soil moisture. Due to abrupt rise and fall in temperature, there is possibility of attack of **Tobacco caterpillar** (Spodoptera litura). The forecasted weather is also favourable for its proliferation. This is a polyphagous pest. So following integrated approach should be started. Use Pheromone trap using Spodo lure to monitor adult population of the insect. In case of appearance of the caterpillar, prepare poison bait with 1 kg rice bran+ 200g jaggery/gur+20-25g Cartap Hydrochloride +Water. Keep the bait at 20-25 different places in the affected field.

Mustard: present weather may favour the infestation of **saw fly, leaf Webber**. To control saw fly, spray 400ml Endosulphan/acre, to control leaf webber spray 200 ml Ithofenprox or Deltamethrin per acre in 200 litr water.

Vegetables:

Tomato: Transplanted seedlings are at 6-8 week stage in main field. **Due deficiency of boron due to hormonal disbalance flower drop occurs.**

Control: Apply timely and required amount of irrigation. Spray Boron 20% 2 gram/lit during early flowering and 15 days later. Due to increase in temperature there may be chances of **bacterial canker and Sun burn** disease in tomato at fruiting stage. To control bacterial canker Farmers should spray Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water. They should use shade net to protect the crop from sun burn. **Bacterial wilt** may be reported in tomato field, to control it drenching with copper oxychloride @ 3g+ plantomycin @1 g/ litre of water.

• **Potato:** There are chances of **Potato Tuber Moth** infestation in potato. To manage this pest chemically spray Spinosad 45 % SC @ 75-ml/acre or Emamectin Benzoate 5% SG @ 80-gram/ acre or Chlorantraniliprole 18.5 % SC @80- ml/acre. To manage **cut worm** in potato crop spray Chlorpyrifos 50% + Cypermethrin 5 % E.C (Lethal-505/Super D) @ 400 ml/acre by mixing it in 200 litre of water.

• **Onion:** There are chances of **purple blotch and Stemphylium blight** disease incidence in onion crop. To manage purple blotch and Stemphylium leaf blight disease in onion spray Mancozeb 75% WP @ 600gram/acre or Difenoconazole 25% E.C @ 200ml/acre or Tebuconazole 25.9% E.C @ 100ml/acre by mixing in 200 litre of water at 15 days interval.

• **Cucutbits:** The current weather is favorable for attack of **red pumpkin beetle** in Cucumber & Bitter gourd. To control it spray Neem based pesticide 300ppm@5ml/litre of water at initial stages and at later stages spray Acephate@1g/litre of water.

• **Fruits:**

• **Mango:** Present weather conditions are favourable for the incidence of **Mango hoppers**. To control go for spraying of phosphamidon @ 0.5ml/l (or) dimethoate 2ml/l (or) chlorpyrifos @2.5ml/l. during initial stages of flowering. Spray Alpha Naphthyl Acetic Acid 4.5% SL @ 4.5 ml in 10 litre of water when tender fruits are in pea size. With the help of ring method of irrigation apply 25 kg FYM, 500 gram of urea and 1 kg of potash and phosphate at each tree. For **thrips** control, spraying of Acephate @1.5 g /litre of water is advocated.

• **Cashew:** Prevailing conditions are suitable for the incidence of Tea mosquito bug. Farmers are advised to go for spraying of Neem oil @ 5ml/L + Monocrotophos @ 1.6ml/L at new blush stage and Chlorpyrifos @ 2ml/L at flowering stage.

Mushroom: Initiate cultivation of paddy straw mushroom. As dry weather condition is prevailed in Cuttack district, Spray water if the mushrooms are at fruiting stage. Cover the Mushroom unit with water soaked Gunny bags to maintain the inside temperature.



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SAKSHI
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-555
Date- 25/02/2025

AGROMET ADVISORY BULLETIN FOR NARASINGHPUR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 06 mm and district received average 03 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.42°C (-1.33) and 20.57°C (+1.79), respectively. The mean relative humidity was 63-87%.

Forecasted period: Today(8.30IST) to 02nd March (8.30IST)2025

Weather advisory: Dry weather is forecasted for next 5 days in the block. The day temperature will be 33-36°C and night temperature will be 19-20°C for upcoming 5 days. Maximum and minimum relative humidity will be 45-90% and 12-22% respectively. **There will be clear to partly cloudy sky for next 5 days.**

Block	Narasinghpur (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	25/02/2025	26/02/2025	27/02/2025	28/02/2025	01/03/2025
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	34	33	34	34	36
T-Min(°C)	19	19	19	20	20
Cloud Cover(/octa)	0	1	0	1	3
RH Max (%)	90	72	72	51	45
RH Min (%)	19	22	16	15	12
Wind Speed (Kmph)	5	6	6	7	9
Wind Direction (deg)	158	171	195	191	200

Agromet Advisory

General advisory:

Farmers should undertake field preparation for cultivation of different summer vegetables like sponge gourd, bottle gourd, pumpkin, water melon, musk melon etc. If seedlings are already raised in polythene bags; transplant in main field @ 2-3 seedlings per pit. Irrigation, fertilizer and pesticide application should be resumed as there will clear sky for upcoming 5 days.

Pulses:

November to December sown crop is at pod formation to seed development (nine to twelve week) stage.

In case of **YMV infestation** uproots and destroys the infected plant and spray imidacloprid 50ml/acre mixing with 200 lit of water. There may be chances of infestation of **Anthracoze and Cercopsera leaf spot** in Pulses. To manage these spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre in 200 ltr water.

Rabi-Groundnut:

Do not disturb the soil after flower initiation in groundnut. Do irrigation in the field based upon soil moisture. Due to abrupt rise and fall in temperature, there is possibility of attack of **Tobacco caterpillar** (Spodoptera litura). The forecasted weather is also favourable for its proliferation. This is a polyphagous pest. So following integrated approach should be started. Use Pheromone trap using Spodo lure to monitor adult population of the insect. In case of appearance of the caterpillar, prepare poison bait with 1 kg rice bran+ 200g jaggery/gur+20-25g Cartap Hydrochloride +Water. Keep the bait at 20-25 different places in the affected field.

Mustard: present weather may favour the infestation of **saw fly, leaf Webber**. To control saw fly, spray 400ml Endosulphan/acre, to control leaf webber spray 200 ml Ithofenprox or Deltamethrin per acre in 200 litr water.

Vegetables:

Tomato: Transplanted seedlings are at 6-8 week stage in main field. **Due deficiency of boron or due to hormonal disbalance flower drop occurs.**

Control: Apply timely and required amount of irrigation. Spray Boron 20% 2 gram/lit during early flowering and 15 days later. Due to increase in temperature there may be chances of **bacterial canker and Sun burn** disease in tomato at fruiting stage. To control bacterial canker farmers should spray Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water. They should use shade net to protect the crop from sun burn. **Bacterial wilt** may be reported in tomato field, to control it drenching with copper oxychloride @ 3g+ plantomycin @ 1 g/ litre of water.

- **Potato:** There are chances of **Potato Tuber Moth** infestation in potato. To manage this pest chemically spray Spinosad 45 % SC @ 75-ml/acre or Emamectin Benzoate 5% SG @ 80-gram/ acre or Chlorantraniliprole 18.5 % SC @ 80- ml/acre. To manage **cut worm** in potato crop spray Chlorpyrifos 50% + Cypermethrin 5 % E.C (Lethal-505/Super D) @ 400 ml/acre by mixing it in 200 litre of water.
- **Onion:** There are chances of **purple blotch and Stemphyllium blight** disease incidence in onion crop. To manage purple blotch and Stemphyllium leaf blight disease in onion spray Mancozeb 75% WP @ 600gram/acre or Difenconazole 25% E.C @ 200ml/acre or Tebuconazole 25.9% E.C @ 100ml/acre by mixing in 200 litre of water at 15 days interval.
- **Cucurbits:** The current weather is favorable for attack of **red pumpkin beetle** in Cucumber & Bitter gourd. To control it spray Neem based pesticide 300ppm@5ml/litre of water at initial stages and at later stages spray Acephate@1g/litre of water.
- **Fruits:**
- **Mango:** Present weather conditions are favourable for the incidence of **Mango hoppers**. To control go for spraying of phosphamidon @ 0.5ml/l (or) dimethoate 2ml/l (or) chlorpyrifos @ 2.5ml/l. during initial stages of flowering. Spray Alpha Naphthyl Acetic Acid 4.5% SL @ 4.5 ml in 10 litre of water when tender fruits are in pea size. With the help of ring method of irrigation apply 25 kg FYM, 500 gram of urea and 1 kg of potash and phosphate at each tree. For **thrips** control, spraying of Acephate @ 1.5 g /litre of water is advocated.
- **Cashew:** Prevailing conditions are suitable for the incidence of Tea mosquito bug. Farmers are advised to go for spraying of Neem oil @ 5ml/L + Monocrotophos @ 1.6ml/L at new blush stage and Chlorpyrifos @ 2ml/L at flowering stage.
- **Mushroom:** Initiate cultivation of paddy straw mushroom. As dry weather condition is prevailed in Cuttack district, Spray water if the mushrooms are at fruiting stage. Cover the Mushroom unit with water soaked Gunny bags to maintain the inside temperature.



Dr Ranjan Kumar Mohanta
 (Nodal officer)

Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightning alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-555
Date- 25/02/2025

AGROMET ADVISORY BULLETIN FOR NIALI

District Past week average weather condition and Block past week average Rainfall condition

The block received average 01 mm and district received average 03 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.42°C (-1.33) and 20.57°C (+1.79), respectively. The mean relative humidity was 63-87%.

Forecasted period: Today(8.30IST) to 02nd March (8.30IST)2025

Weather advisory: Dry weather is forecasted for next 5 days in the block. The day temperature will be 32-35°C and night temperature will be 22-24°C for upcoming 5 days. Maximum and minimum relative humidity will be 73-85% and 21-36% respectively. **There will be clear to partly cloudy sky for next 5 days.**

Block	Niali (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	25/02/2025	26/02/2025	27/02/2025	28/02/2025	01/03/2025
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	32	32	33	34	35
T-Min(°C)	22	22	22	23	24
Cloud Cover(octa)	1	0	0	1	1
RH Max (%)	84	85	85	83	73
RH Min (%)	36	29	29	27	21
Wind Speed(Kmph)	8	10	11	14	18
Wind Direction(deg)	120	165	203	203	204

Agromet Advisory

General advisory:

Farmers should undertake field preparation for cultivation of different summer vegetables like sponge gourd, bottle gourd, pumpkin, water melon, musk melon etc. If seedlings are already raised in polythene bags; transplant in main field @ 2-3 seedlings per pit. Irrigation, fertilizer and pesticide application should be resumed as there will clear sky for upcoming 5 days.

Pulses: November to December sown crop is at pod formation to seed development (nine to twelve week) stage.

In case of **YMV infestation** uproots and destroys the infected plant and spray imidacloprid 50ml/acre mixing with 200 lit of water. There may be chances of infestation of **Anthraxnose and Cercopsera leaf spot** in Pulses. To manage these spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre in 200 ltr water.

Rabi-Groundnut:

Do not disturb the soil after flower initiation in groundnut. Do irrigation in the field based upon soil moisture. Due to abrupt rise and fall in temperature, there is possibility of attack of **Tobacco caterpillar** (Spodoptera litura). The forecasted weather is also favourable for its proliferation. This is a polyphagous pest. So following integrated approach should be started. Use Pheromone trap using Spodo lure to monitor adult population of the insect. In case of appearance of the caterpillar, prepare poison bait with 1 kg rice bran+ 200g jaggery/gur+20-25g Cartap Hydrochloride +Water. Keep the bait at 20-25 different places in the affected field.

Mustard: present weather may favour the infestation of **saw fly, leaf Webber**. To control saw fly, spray 400ml Endosulphan/acre, to control leaf webber spray 200 ml Ithofenprox or Deltamethrin per acre in 200 litre water.

Vegetables:

Tomato: Transplanted seedlings are at 6-8 week stage in main field. **Due deficiency of boron due to hormonal disbalance flower drop occurs.**

Control: Apply timely and required amount of irrigation. Spray Boron 20% 2 gram/litre during early flowering and 15 days later. Due to increase in temperature there may be chances of **bacterial canker and Sun burn** disease in tomato at fruiting stage. To control bacterial canker Farmers should spray Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water. They should use shade net to protect the crop from sun burn. **Bacterial wilt** may be reported in tomato field, to control it drenching with copper oxychloride @ 3g+ plantomycin @1 g/ litre of water.

• **Potato:** There are chances of **Potato Tuber Moth** infestation in potato. To manage this pest chemically spray Spinosad 45 % SC @ 75-ml/acre or Emamectin Benzoate 5% SG @ 80-gram/ acre or Chlorantraniliprole 18.5 % SC @80- ml/acre. To manage **cut worm** in potato crop spray Chlorpyrifos 50% + Cypermethrin 5 % E.C (Lethal-505/Super D) @ 400 ml/acre by mixing it in 200 litre of water.

• **Onion:** There are chances of **purple blotch and Stemphyllium blight** disease incidence in onion crop. To manage purple blotch and Stemphyllium leaf blight disease in onion spray Mancozeb 75% WP @ 600gram/acre or Difenconazole 25% E.C @ 200ml/acre or Tebuconazole 25.9% E.C @ 100ml/acre by mixing in 200 litre of water at 15 days interval.

• **Cucurbits:** The current weather is favorable for attack of **red pumpkin beetle** in Cucumber & Bitter gourd. To control it spray Neem based pesticide 300ppm@5ml/litre of water at initial stages and at later stages spray Acephate@1g/litre of water.

• **Fruits:**

• **Mango:** Present weather conditions are favourable for the incidence of **Mango hoppers**. To control go for spraying of phosphamidon @ 0.5ml/l (or) dimethoate 2ml/l (or) chlorpyrifos @2.5ml/l. during initial stages of flowering. Spray Alpha Naphthyl Acetic Acid 4.5% SL @ 4.5 ml in 10 litre of water when tender fruits are in pea size. With the help of ring method of irrigation apply 25 kg FYM, 500 gram of urea and 1 kg of potash and phosphate at each tree. For **thrips** control, spraying of Acephate @1.5 g /litre of water is advocated.

• **Cashew:** Prevailing conditions are suitable for the incidence of Tea mosquito bug. Farmers are advised to go for spraying of Neem oil @ 5ml/L + Monocrotophos @ 1.6ml/L at new blush stage and Chlorpyrifos @ 2ml/L at flowering stage.

• **Mushroom:** Initiate cultivation of paddy straw mushroom. As dry weather condition is prevailed in Cuttack district, Spray water if the mushrooms are at fruiting stage. Cover the Mushroom unit with water soaked Gunny bags to maintain the inside temperature.



DrRanjan Kumar Mohanta
 (Nodal officer)

Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightning alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-555
Date- 25/02/2025

AGROMET ADVISORY BULLETIN FOR NISCHINTAKOILI
District Past week average weather condition and Block past week average Rainfall condition

The block received average 01 mm and district received average 03 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.42°C (-1.33) and 20.57°C (+1.79), respectively. The mean relative humidity was 63-87%.

Forecasted period: Today(8.30IST) to 02nd March (8.30IST)2025

Weather advisory: Dry weather is forecasted for next 5 days in the block. The day temperature will be 31-35°C and night temperature will be 22-23°C for upcoming 5 days. Maximum and minimum relative humidity will be 65-85% and 19-33% respectively. **There will be clear to partly cloudy sky for next 5 days.**

Block	Nischintakoili (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	25/02/2025	26/02/2025	27/02/2025	28/02/2025	01/03/2025
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	32	31	33	34	35
T-Min(°C)	22	22	22	22	23
Cloud Cover(octa)	1	0	0	1	1
RH Max (%)	84	84	85	81	65
RH Min (%)	33	26	24	22	19
Wind Speed (Kmph)	7	9	10	12	18
Wind Direction (deg)	124	243	203	203	207

Agromet Advisory

General advisory:

Farmers should undertake field preparation for cultivation of different summer vegetables like sponge gourd, bottle gourd, pumpkin, water melon, musk melon etc. If seedlings are already raised in polythene bags; transplant in main field @ 2-3 seedlings per pit. Irrigation, fertilizer and pesticide application should be resumed as there will clear sky for upcoming 5 days.

Pulses:

November to December sown crop is at pod formation to seed development (nine to twelve week) stage.

In case of **YMV infestation** uproots and destroys the infected plant and spray imidacloprid 50ml/acre mixing with 200 lit of water. There may be chances of infestation of **Anthraxnose and Cercopsera leaf spot** in Pulses. To manage these spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre in 200 ltr water.

Rabi-Groundnut:

Do not disturb the soil after flower initiation in groundnut. Do irrigation in the field based upon soil moisture. Due to abrupt rise and fall in temperature, there is possibility of attack of **Tobacco caterpillar** (Spodoptera litura). The forecasted weather is also favourable for its proliferation. This is a polyphagous pest. So following integrated approach should be started. Use Pheromone trap using Spodolure to monitor adult population of the insect. In case of appearance of the caterpillar, prepare poison bait with 1 kg rice bran+ 200g jaggery/gur+20-25g Cartap Hydrochloride +Water. Keep the bait at 20-25 different places in the affected field.

Mustard: present weather may favour the infestation of **saw fly, leaf Webber**. To control saw fly, spray 400ml Endosulphan/acre, to control leaf webber spray 200 ml Ithofenprox or Deltamethrin per acre in 200 litre water.

Vegetables:

Tomato: Transplanted seedlings are at 6-8 week stage in main field. **Due deficiency of boron due to hormonal disbalance flower drop occurs.** Control: Apply timely and required amount of irrigation. Spray Boron 20% 2 gram/lit during early flowering and 15 days later. Due to increase in temperature there may be chances of **bacterial canker and Sun burn** disease in tomato at fruiting stage. To control bacterial canker farmers should spray Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water. They should use shade net to protect the crop from sun burn. **Bacterial wilt** may be reported in tomato field, to control it drenching with copper oxychloride @ 3g+ plantomycin @1 g/ litre of water.

• **Potato:** There are chances of **Potato Tuber Moth** infestation in potato. To manage this pest chemically spray Spinosad 45 % SC @ 75-ml/acre or Emamectin Benzoate 5% SG @ 80-gram/ acre or Chlorantraniliprole 18.5 % SC @80- ml/acre. To manage **cut worm** in potato crop spray Chlorpyrifos 50% + Cypermethrin 5 % E.C (Lethal-505/Super D) @ 400 ml/acre by mixing it in 200 litre of water.

• **Onion:** There are chances of **purple blotch and Stemphylium blight** disease incidence in onion crop. To manage purple blotch and Stemphylium leaf blight disease in onion spray Mancozeb 75% WP @ 600gram/acre or Difenoconazole 25% E.C @ 200ml/acre or Tebuconazole 25.9% E.C @ 100ml/acre by mixing in 200 litre of water at 15 days interval.

• **Cucurbits:** The current weather is favorable for attack of **red pumpkin beetle** in Cucumber & Bitter gourd. To control it spray Neem based pesticide 300ppm@5ml/litre of water at initial stages and at later stages spray Acephate@1g/litre of water.

Fruits:

• **Mango:** Present weather conditions are favourable for the incidence of **Mango hoppers**. To control go for spraying of phosphamidon @ 0.5ml/l (or) dimethoate 2ml/l (or) chlorpyrifos @2.5ml/l. during initial stages of flowering. Spray Alpha Naphthyl Acetic Acid 4.5% SL @ 4.5 ml in 10 litre of water when tender fruits are in pea size. With the help of ring method of irrigation apply 25 kg FYM, 500 gram of urea and 1 kg of potash and phosphate at each tree. For **thrips** control, spraying of Acephate @1.5 g /litre of water is advocated.

• **Cashew:** Prevailing conditions are suitable for the incidence of Tea mosquito bug. Farmers are advised to go for spraying of Neem oil @ 5ml/L + Monocrotophos @ 1.6ml/L at new blush stage and Chlorpyrifos @ 2ml/L at flowering stage.

• **Mushroom:** Initiate cultivation of paddy straw mushroom. As dry weather condition is prevailed in Cuttack district, Spray water if the mushrooms are at fruiting stage. Cover the Mushroom unit with water soaked Gunny bags to maintain the inside temperature.



DrRanjan Kumar Mohanta
 (Nodal officer)

Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightning alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-555
Date- 25/02/2025

AGROMET ADVISORY BULLETIN FOR SALEPUR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 03 mm and district received average 03 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.42°C (-1.33) and 20.57°C (+1.79), respectively. The mean relative humidity was 63-87%.

Forecasted period: Today(8.30IST) to 02nd March (8.30IST)2025

Weather advisory: Dry weather is forecasted for next 5days in the block. The day temperature will be 31-35°C and night temperature will be 22-23°C for upcoming 5 days. Maximum and minimum relative humidity will be 65-85% and 19-32% respectively. **There will be clear to partly cloudy sky for next 5days.**

Block	Salepur (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	25/02/2025	26/02/2025	27/02/2025	28/02/2025	01/03/2025
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	32	31	33	34	35
T-Min(°C)	22	22	22	22	23
Cloud Cover(octa)	1	0	0	1	1
RH Max (%)	83	84	85	80	65
RH Min (%)	32	26	23	21	19
Wind Speed (Kmph)	7	9	10	12	18
Wind Direction (deg)	124	243	203	201	207

Agromet Advisory

General advisory:

Farmers should undertake field preparation for cultivation of different summer vegetables like sponge gourd, bottle gourd, pumpkin, water melon, musk melon etc. If seedlings are already raised in polythene bags; transplant in main field @ 2-3 seedlings per pit. Irrigation, fertilizer and pesticide application should be resumed as there will clear sky for upcoming 5 days.

Pulses:

November to December sown crop is at pod formation to seed development (nine to twelve week) stage.

In case of **YMV infestation** uproots and destroys the infected plant and spray imidacloprid 50ml/acre mixing with 200 lit of water. There may be chances of infestation of **Anthraxnose and Cercopersa leaf spot** in Pulses. To manage these spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre in 200 ltr water.

Rabi-Groundnut:

Do not disturb the soil after flower initiation in groundnut. Do irrigation in the field based upon soil moisture. Due to abrupt rise and fall in temperature, there is possibility of attack of **Tobacco caterpillar** (Spodoptera litura). The forecasted weather is also favourable for its proliferation. This is a polyphagous pest. So following integrated approach should be started. Use Pheromone trap using Spodo lure to monitor adult population of the insect. In case of appearance of the caterpillar, prepare poison bait with 1 kg rice bran+ 200g jaggery/gur+20-25g Cartap Hydrochloride +Water. Keep the bait at 20-25 different places in the affected field.

Mustard: present weather may favour the infestation of **saw fly, leaf Webber**. To control saw fly, spray 400ml Endosulphan/acre, to control leaf webber spray 200 ml Ithofenprox or Deltamethrin per acre in 200 litr water.

Vegetables:

Tomato: Transplanted seedlings are at 6-8 week stage in main field. **Due deficiency of boron or due to hormonal disbalance flower drop occurs.**

Control: Apply timely and required amount of irrigation. Spray Boron 20% 2 gram/lit during early flowering and 15 days later. Due to increase in temperature there may be chances of **bacterial canker and Sun burn** disease in tomato at fruiting stage. To control bacterial canker Farmers should spray Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water. They should use shade net to protect the crop from sun burn. **Bacterial wilt** may be reported in tomato field, to control it drenching with copper oxychloride @ 3g+ plantomycin @1 g/ litre of water.

• **Potato:** There are chances of **Potato Tuber Moth** infestation in potato. To manage this pest chemically spray Spinosad 45 % SC @ 75-ml/acre or Emamectin Benzoate 5% SG @ 80-gram/ acre or Chlorantraniliprole 18.5 % SC @80- ml/acre. To manage **cut worm** in potato crop spray Chlorpyrifos 50% + Cypermethrin 5 % E.C (Lethal-505/Super D) @ 400 ml/acre by mixing it in 200 litre of water.

• **Onion:** There are chances of **purple blotch and Stemphyllium blight** disease incidence in onion crop. To manage purple blotch and Stemphyllium leaf blight disease in onion spray Mancozeb 75% WP @ 600gram/acre or Difenoconazole 25% E.C @ 200ml/acre or Tebuconazole 25.9% E.C @ 100ml/acre by mixing in 200 litre of water at 15 days interval.

• **Cucutbits:** The current weather is favorable for attack of **red pumpkin beetle** in Cucumber & Bitter gourd. To control it spray Neem based pesticide 300ppm @ 5ml/litre of water at initial stages and at later stages spray Acephate @ 1g/litre of water.

Fruits:

• **Mango:** Present weather conditions are favourable for the incidence of **Mango hoppers**. To control go for spraying of phosphamidon @ 0.5ml/l (or) dimethoate 2ml/l (or) chlorpyrifos @ 2.5ml/l. during initial stages of flowering. Spray Alpha Naphthyl Acetic Acid 4.5% SL @ 4.5 ml in 10 litre of water when tender fruits are in pea size. With the help of ring method of irrigation apply 25 kg FYM, 500 gram of urea and 1 kg of potash and phosphate at each tree. For **thrips** control, spraying of Acephate @ 1.5 g /litre of water is advocated.

• **Cashew:** Prevailing conditions are suitable for the incidence of Tea mosquito bug. Farmers are advised to go for spraying of Neem oil @ 5ml/L + Monocrotophos @ 1.6ml/L at new blush stage and Chlorpyrifos @ 2ml/L at flowering stage.

Mushroom: Initiate cultivation of paddy straw mushroom. As dry weather condition is prevailed in Cuttack district, Spray water if the mushrooms are at fruiting stage. Cover the Mushroom unit with water soaked Gunny bags to maintain the inside temperature.



DrRanjan Kumar Mohanta
 (Nodal officer)

Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-555
Date- 25/02/2025

AGROMET ADVISORY BULLETIN FOR TANGI-CHOUDWAR

District Past week average weather condition and Block past week average Rainfall condition

The block received average 02 mm and district received average 03 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.42°C (-1.33) and 20.57°C (+1.79), respectively. The mean relative humidity was 63-87%.

Forecasted period: Today(8.30IST) to 02nd March (8.30IST)2025

Weather advisory: Dry weather is forecasted for next 5 days in the block. The day temperature will be 32-35°C and night temperature will be 23-25°C for upcoming 5 days. Maximum and minimum relative humidity will be 71-82% and 31-42% respectively. **There will be clear to partly cloudy sky for next 5 days.**

Block	Tangi-Choudwar (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
Date	25/02/2025	26/02/2025	27/02/2025	28/02/2025	01/03/2025
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	32	32	33	34	35
T-Min(°C)	23	23	23	24	25
Cloud Cover(octa)	0	0	0	1	1
RH Max (%)	81	82	80	80	71
RH Min (%)	42	36	38	38	31
Wind Speed (Kmph)	11	11	13	15	20
Wind Direction (deg)	120	146	169	203	212

Agromet Advisory

General advisory:

Farmers should undertake field preparation for cultivation of different summer vegetables like sponge gourd, bottle gourd, pumpkin, water melon, musk melon etc. If seedlings are already raised in polythene bags; transplant in main field @ 2-3 seedlings per pit. Irrigation, fertilizer and pesticide application should be resumed as there will clear sky for upcoming 5 days.

Pulses:

November to December sown crop is at pod formation to seed development (nine to twelve week) stage.

In case of **YMV infestation** uproots and destroys the infected plant and spray imidacloprid 50ml/acre mixing with 200 lit of water. There may be chances of infestation of **Anthraxnose and Cercopsera leaf spot** in Pulses. To manage these spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre in 200 ltr water.

Rabi-Groundnut:

Do not disturb the soil after flower initiation in groundnut. Do irrigation in the field based upon soil moisture. Due to abrupt rise and fall in temperature, there is possibility of attack of **Tobacco caterpillar** (Spodoptera litura). The forecasted weather is also favourable for its proliferation. This is a polyphagous pest. So following integrated approach should be started. Use Pheromone trap using Spodo lure to monitor adult population of the insect. In case of appearance of the caterpillar, prepare poison bait with 1 kg rice bran+ 200g jaggery/gur+20-25g Cartap Hydrochloride +Water. Keep the bait at 20-25 different places in the affected field.

Mustard: present weather may favour the infestation of **saw fly, leaf Webber**. To control saw fly, spray 400ml Endosulphan/acre, to control leaf webber spray 200 ml Ithofenprox or Deltamethrin per acre in 200 litr water.

Vegetables:

Tomato: Transplanted seedlings are at 6-8 week stage in main field. **Due deficiency of boron or due to hormonal disbalance flower drop occurs.**

Control: Apply timely and required amount of irrigation. Spray Boron 20% 2 gram/lit during early flowering and 15 days later. Due to increase in temperature there may be chances of **bacterial canker and Sun burn** disease in tomato at fruiting stage. To control bacterial canker farmers should spray Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water. They should use shade net to protect the crop from sun burn. **Bacterial wilt** may be reported in tomato field, to control it drenching with copper oxychloride @ 3g+ plantomycin @1 g/ litre of water.

• **Potato:** There are chances of **Potato Tuber Moth** infestation in potato. To manage this pest chemically spray Spinosad 45 % SC @ 75-ml/acre or Emamectin Benzoate 5% SG @ 80-gram/ acre or Chlorantraniliprole 18.5 % SC @80- ml/acre. To manage **cut worm** in potato crop spray Chlorpyrifos 50% + Cypermethrin 5 % E.C (Lethal-505/Super D) @ 400 ml/acre by mixing it in 200 litre of water.

• **Onion:** There are chances of **purple blotch and Stemphyllium blight** disease incidence in onion crop. To manage purple blotch and Stemphyllium leaf blight disease in onion spray Mancozeb 75% WP @ 600gram/acre or Difenoconazole 25% E.C @ 200ml/acre or Tebuconazole 25.9% E.C @ 100ml/acre by mixing in 200 litre of water at 15 days interval.

• **Cucutbits:** The current weather is favorable for attack of **red pumpkin beetle** in Cucumber & Bitter gourd. To control it spray Neem based pesticide 300ppm@5ml/litre of water at initial stages and at later stages spray Acephate@1g/litre of water.

Fruits:

• **Mango:** Present weather conditions are favourable for the incidence of **Mango hoppers**. To control go for spraying of phosphamidon @ 0.5ml/l (or) dimethoate 2ml/l (or) chloropyrifos @2.5ml/l. during initial stages of flowering. Spray Alpha Naphthyl Acetic Acid 4.5% SL @ 4.5 ml in 10 litre of water when tender fruits are in pea size. With the help of ring method of irrigation apply 25 kg FYM, 500 gram of urea and 1 kg of potash and phosphate at each tree. For **thrips** control, spraying of Acephate @1.5 g /litre of water is advocated.

• **Cashew:** Prevailing conditions are suitable for the incidence of Tea mosquito bug. Farmers are advised to go for spraying of Neem oil @ 5ml/L + Monocrotophos @ 1.6ml/L at new blush stage and Chlorpyriqos @ 2ml/L at flowering stage.

Mushroom: Initiate cultivation of paddy straw mushroom. As dry weather condition is prevailed in Cuttack district, Spray water if the mushrooms are at fruiting stage. Cover the Mushroom unit with water soaked Gunny bags to maintain the inside temperature.



DrRanjan Kumar Mohanta
 (Nodal officer)

Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightening alert in your area





GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-555
Date- 25/02/2025

AGROMET ADVISORY BULLETIN FOR TIGIRIA

District Past week average weather condition and Block past week average Rainfall condition

The block received average 03 mm and district received average 03 mm rainfall during the last week. The mean daily maximum and minimum temperature was 31.42°C (-1.33) and 20.57°C (+1.79), respectively. The mean relative humidity was 63-87%.

Forecasted period: Today(8.30IST) to 02nd March (8.30IST)2025

Weather advisory: Dry weather is forecasted for next 5 days in the block. The day temperature will be 33-36°C and night temperature will be 20-22°C for upcoming 5 days. Maximum and minimum relative humidity will be 53-84% and 13-23% respectively. There will be clear to partly cloudy sky for next 5 days.

Block	Tigiria (VALID TILL 08:30 IST OF THE NEXT 5 DAYS)				
	25/02/2025	26/02/2025	27/02/2025	28/02/2025	01/03/2025
Rainfall(mm)	0	0	0	0	0
T-Max(°C)	33	33	33	34	36
T-Min(°C)	21	21	20	22	22
Cloud Cover(octa)	0	0	0	1	2
RH Max (%)	84	79	78	59	53
RH Min (%)	23	22	17	16	13
Wind Speed (Kmph)	7	8	8	9	13
Wind Direction (deg)	153	167	203	180	206

Agro met Advisory

General advisory:

Farmers should undertake field preparation for cultivation of different summer vegetables like sponge gourd, bottle gourd, pumpkin, water melon, musk melon etc. If seedlings are already raised in polythene bags; transplant in main field @ 2-3 seedlings per pit. Irrigation, fertilizer and pesticide application should be resumed as there will clear sky for upcoming 5 days.

Pulses:

November to December sown crop is at pod formation to seed development (nine to twelve week) stage.

In case of **YMV infestation** uproots and destroys the infected plant and spray imidacloprid 50ml/acre mixing with 200 lit of water. There may be chances of infestation of **Anthraxnose and Cercopsera leaf spot** in Pulses. To manage these spray Metalaxyl 8% + Mancozeb 64% W.P @ 400- gram/acre in 200 ltr water.

Rabi-Groundnut:

Do not disturb the soil after flower initiation in groundnut. Do irrigation in the field based upon soil moisture. Due to abrupt rise and fall in temperature, there is possibility of attack of **Tobacco caterpillar** (Spodoptera litura). The forecasted weather is also favourable for its proliferation. This is a polyphagous pest. So following integrated approach should be started. Use Pheromone trap using Spodolure to monitor adult population of the insect. In case of appearance of the caterpillar, prepare poison bait with 1 kg rice bran+ 200g jaggery/gur+20-25g Cartap Hydrochloride +Water. Keep the bait at 20-25 different places in the affected field.

Mustard: present weather may favour the infestation of **saw fly, leaf Webber**. To control saw fly, spray 400ml Endosulphan/acre, to control leaf webber spray 200 ml Ithofenprox or Deltamethrin per acre in 200 litre water.

Vegetables:

Tomato: Transplanted seedlings are at 6-8 week stage in main field. **Due deficiency of boron due to hormonal disbalance flower drop occurs.** Control: Apply timely and required amount of irrigation. Spray Boron 20% 2 gram/lit during early flowering and 15 days later. Due to increase in temperature there may be chances of **bacterial canker and Sun burn** disease in tomato at fruiting stage. To control bacterial canker farmers should spray Copper oxychloride 50% WP @ 3 g + Plantomycin @ 1 g/litre of water. They should use shade net to protect the crop from sun burn. **Bacterial wilt** may be reported in tomato field, to control it drenching with copper oxychloride @ 3g+ plantomycin @1 g/ litre of water.

• **Potato:** There are chances of **Potato Tuber Moth** infestation in potato. To manage this pest chemically spray Spinosad 45 % SC @ 75-ml/acre or Emamectin Benzoate 5% SG @ 80-gram/ acre or Chlorantraniliprole 18.5 % SC @80- ml/acre. To manage **cut worm** in potato crop spray Chlorpyrifos 50% + Cypermethrin 5 % E.C (Lethal-505/Super D) @ 400 ml/acre by mixing it in 200 litre of water.

• **Onion:** There are chances of **purple blotch and Stemphylium blight** disease incidence in onion crop. To manage purple blotch and Stemphylium leaf blight disease in onion spray Mancozeb 75% WP @ 600gram/acre or Difenconazole 25% E.C @ 200ml/acre or Tebuconazole 25.9% E.C @ 100ml/acre by mixing in 200 litre of water at 15 days interval.

• **Cucurbits:** The current weather is favorable for attack of **red pumpkin beetle** in Cucumber & Bitter gourd. To control it spray Neem based pesticide 300ppm@5ml/litre of water at initial stages and at later stages spray Acephate@1g/litre of water.

Fruits:

• **Mango:** Present weather conditions are favourable for the incidence of **Mango hoppers**. To control go for spraying of phosphamidon @ 0.5ml/l (or) dimethoate 2ml/l (or) chlorpyrifos @2.5ml/l. during initial stages of flowering. Spray Alpha Naphthyl Acetic Acid 4.5% SL @ 4.5 ml in 10 litre of water when tender fruits are in pea size. With the help of ring method of irrigation apply 25 kg FYM, 500 gram of urea and 1 kg of potash and phosphate at each tree. For **thrips** control, spraying of Acephate @1.5 g /litre of water is advocated.

• **Cashew:** Prevailing conditions are suitable for the incidence of Tea mosquito bug. Farmers are advised to go for spraying of Neem oil @ 5ml/L + Monocrotophos @ 1.6ml/L at new blush stage and Chlorpyrifos @ 2ml/L at flowering stage.

Mushroom: Initiate cultivation of paddy straw mushroom. As dry weather condition is prevailed in Cuttack district, Spray water if the mushrooms are at fruiting stage. Cover the Mushroom unit with water soaked Gunny bags to maintain the inside temperature.



DrRanjan Kumar Mohanta
 (Nodal officer)

Follow Meghdoot Mobile APP for Agromet
 advisories in local language and DAMINI
 APP for Lightning alert in your area







GRAMIN KRISHI MAUSAM SEWA
INDIA METEOROLOGICAL DEPARTMENT
DISTRICT AGROMET UNIT
KRISHI VIGYAN KENDRA CUTTACK
 (A Unit of ICAR-National Rice Research Institute)
 Email: kvkcuttack@gmail.com/damukvkcuttack@gmail.com



Bulletin No.-555
Date- 25/02/2025

				
MEGHDOOT APP	DAMINI APP	FACEBOOK (Damu Cuttack)	BLOCK SPECIFIC WHAT'S APP GROUPS	@DAMUCUTTACK

- Kindly install and use **“Meghdoot mobile App”** for Cuttack district weather forecast, warnings and weather based agromet advisory services.
<https://play.google.com/store/apps/details?id=com.aas.meghdoot>
- Kindly install and use **“Damini mobile App”** for location specific for lightening forecast and warnings.
<https://play.google.com/store/apps/details?id=com.lightening.live.damini>
- Follow the **block specific whats app groups** and join in your block wise groups for getting weather based agromet advisory services twice in a week (Tuesday and Friday) and other weather warning based farm management advisories.
- Follow the **Official Website links** to get the weather based agromet advisory service bulletins (**English & Odia**) twice in a week (Tuesday and Friday).

1	ICAR-NRRI, Cuttack, Odisha	https://icar-nrri.in/agromate/	
2	Agrimet division, IMD, Pune	https://imdagrmet.gov.in/imdproject/AGIndex.php	



DrRanjan Kumar Mohanta
(Nodal officer)

Follow Meghdoot Mobile APP for Agromet advisories in local language and DAMINI APP for Lightening alert in your area

